

IMPACT OF MYCORRHIZAL FUNGI AND RHIZOSPHERE MICROORGANISMS ON MAIZE GRAIN YIELD AND CHEMICAL COMPOSITION

C. Tripaldi¹, M. Novero², S. Di Giovanni¹, P.M. Chiarabaglio³, P. Lorenzoni⁴, D. Meo Zilio^{1,*}, G. Palocci¹, C. Balconi⁵ and R. Aleandri⁴

¹Council for Agricultural Research and Economics, Research Centre for Animal Production and Aquaculture, Via Salaria 31, 00015 Monterotondo (RM), Italy; ²University of Torino, Department of Life Sciences and Systems Biology, Viale Mattioli 25, 10125 Torino, Italy; ³Council for Agricultural Research and Economics, Research Centre for Forestry and Wood, Strada per Frassineto 35, 15033 Casale Monferrato (AL), Italy; ⁴Council for Agricultural Research and Economics, Via Po 14, 00198 Roma, Italy; ⁵Council for Agricultural Research and Economics, Research Centre for Cereal and Industrial Crops, Via Stezzano 24, 24126 Bergamo, Italy.
*Corresponding author's email: david.meozilio@crea.gov.it

Yield and grain characteristics of maize plants grown in open-field conditions were evaluated after inoculation with Micosat F[®] on two different soils under dry and watered conditions. The mycorrhizal frequency and intensity were higher in inoculated maize (87.8 vs 80.3% and 26.8 vs 17.5%, respectively). The abundance of arbuscules in the root system was also higher in inoculated plants (9.7 vs 5.8%). The treatment did not affect grain yield. Positive effect of Typic Eutrudept soil on grain yield was observed. The irrigation effect on grain yield was evident only under draught conditions. Chemical characteristics of grain did not change substantially according to the experimental treatments; nevertheless, the NIRs indicated some physical differences among mycorrhized and not mycorrhized samples.

Keywords: Maize grain, root colonization, yield, composition, NIR, electronic nose.

INTRODUCTION

In most of plant species, about 80%, roots can live in symbiosis with arbuscular mycorrhizal (AM) fungi (Smith and Read, 2008), wherein the AM fungi obtain carbon from the plant partner while they transfer mineral nutrients (mostly phosphate) from the soil to the plant root. Advantages derived from AM colonization include enhanced resistance to foliar-feeding insects and soil pathogens, improved resistance to drought (Celebi *et al.*, 2010; Zoppellari *et al.*, 2014), tolerance of salinity and heavy metals and maintenance of soil aggregate stability other than an improved plant nutrition (Gosling *et al.*, 2006). These aspects make the efficient use of AM fungi very appealing to improve plants nutrition. Indeed, it is generally known that the efficiency of phosphorus fertilization is quite low, ranging from 10 to 30% (Gilani, 1983; Isherword, 1998; Gyaneshwar *et al.*, 2002; Hao *et al.*, 2002). Phosphorus deficiency decreases agricultural productivity on more than two billion hectares worldwide (Oberson *et al.*, 2001; Krey *et al.*, 2013).

Besides, owing to environmental issues, there is a strong interest to reduce the use of agrochemicals and to save water. Therefore, research must be directed to a sustainable crop yield improvement. The targeted use of AM fungi on cropping may allow the attainment of acceptable yield with minimum fertilizer dose, while reducing production costs and

environmental pollution risk (Covacevich *et al.*, 2007). Several studies have stressed the positive effect of AM crop inoculation on growth, nutrient uptake and yield for distinct species such as tomato (Candido *et al.*, 2013), potatoes (Douds, 2007) and many other cereal and vegetable crops under field conditions (Sharm *et al.*, 2007; Hamel and Plenchette, 2007).

Plant growth promoting rhizobacteria species (PGPR) are also very effective in promoting plant growth and yield through direct and indirect mechanisms (Günes *et al.*, 2014) and they have been proposed as a sustainable component of nutrient management (Richardson and Simpson, 2011; Krey *et al.*, 2013). Bacteria can release growth stimulating hormones, protect against soil-borne pathogens (Vassilev *et al.*, 2006; Krey *et al.*, 2013), improve mineral nutrition (*e.g.* by increasing plant availability of phosphorus in soil) and they are capable to promote mycorrhiza colonization of maize root as well as maize growth (Krey *et al.*, 2013). Although combined application of PGPR and fungi could be a meaningful approach for sustainable agriculture, there are still certain aspects that need further investigations for obtaining maximum benefits in terms of improved plant growth, particularly under stress conditions (Nadeem *et al.*, 2014). Owing to the multifactorial origin of the interaction between plant and microorganisms, one important aspect is the evaluation of this approach under field conditions.

Besides, most field studies evaluated the plant response to single AM fungal strain, while only few reports showed the effects of mixed exotic inoculum (Pellegrino *et al.*, 2011). Responses of plant to exotic inocula are supposed to depend on some factors including: physical and chemical soil characteristics (Davis *et al.*, 1983; George, 2000), native mycorrhizal density, host plant (Pellegrino *et al.*, 2011). Moreover, the effect of inocula on seed characteristics of major crops has been little focused (Berta *et al.*, 2014).

More efforts are needed to clarify what strains of fungi, PGPR as well as their combination are most effective in plant helping and benefit and to evaluate what is the effectiveness of co-inoculation in a multi-stressed natural environment for commercialization of microbial inocula (Nadeem *et al.*, 2014).

The Micosat F®, patented by an Italian company, contains symbiotic fungi, rhizosphere bacteria, saprophytic fungi and yeast. The aim of our work was to study the effect of Micosat F® inoculation on maize grown under field conditions. Yield and grain characteristics of mycorrhizal maize plants cultivated in two different soils and under dry and watered conditions were evaluated.

MATERIALS AND METHODS

The trial was carried out during two cropping years (2011-12, at the experimental farm of Colle S. Pastore (Rieti, Italy).

Experimental fields: The experimental design included the preparation of eight parcels to study the effects; mycorrhization (present to absent), soil (fine texture to coarse texture) and irrigation (dry to irrigated).

Eight parcels delimited by drains, according to the natural allotment, were prepared. The average size of parcels was 0.8 ha. The two soil types utilized were classified according to Soil Taxonomy (Soil Survey Staff, 2003) one Typic Eutrudept with silty loam texture, located in Piedifiume (PF) and the other Vertic Eutrudept with silty clay texture with tendency to surface fissuring (vertic characteristics) on dry season, located in Casabianca (CB) (Raglione *et al.*, 2011). The PF soil characteristics were with pH 8.1, cation exchange capacity 17.9 cmol (+) kg⁻¹, organic C 9.80 g kg⁻¹, organic matter 16.80 g kg⁻¹, N 1.5 g kg⁻¹, P₂O₅ 30.10 mg kg⁻¹. The CB soil had pH 7.7, cation exchange capacity 35.7 cmol (+) kg⁻¹, organic C 14.5 g kg⁻¹, organic matter 24.9 g kg⁻¹, N 2.1 g kg⁻¹, P₂O₅ 36.7 mg kg⁻¹ (Raglione *et al.*, 2011). Plots have been subjected to maize monoculture for the last 10 years at least.

Agronomic practices: Ploughing, grubbing and harrowing operations were performed before seeding. Fertilizer was applied at seeding with 92 kg/ha of P₂O₅ and 36 kg ha⁻¹ of N. An additional dose of 138 kg ha⁻¹ of N as urea was applied at leaf stage. 75000 plants/ha of NK Famoso cv class 500 were sown. Soil geodisinfestation was not performed.

MICOSAT F® was applied at sowing (May, 3, 2012) on the four selected fields, according to producer recommendations

(15 kg ha⁻¹). MICOSAT F® label composition was three AM species (Glomus caledonium GM24, Glomus intraradices GG31, Glomus coronatum GU53, in form of spores, hyphae and root fragments), and three Plant Growth Promoting Rhizobacteria species (Pseudomonas fluorescens PA28, Pseudo -monas borealis PA29, Bacillus subtilis BA41) with a total concentration of 10⁶ cells g⁻¹. The inoculum was applied in the same plots during both years. The herbicide Primagram Gold (metolachlor e terbutilazina) was used in pre-emergence, while in post-emergence, Glitter (nicosulfuron) and Maicol (dicamba) according to the producer recommended doses were administered. Four fields were assigned to dry cropping. For all the others only an emergency sprinkler irrigation was provided at the bloom phase.

Sampling and analyses of roots: Root samples were analyzed to evaluate AM colonization. These were collected only in the second cropping year. The samples were collected in two PF dry parcels (inoculated and uninoculated), at beginning of August. Five strips were obtained from each parcel. Five root systems were sampled from each strip. Three microscope slides per root were prepared. Each slide contained 20 pieces one cm long. In total, 150 slides and 3000 pieces of root were analyzed. Root fragments were stained according to the method described in Novero *et al.* (2002). Four traits considered were (1) F % (frequency of mycorrhization), reporting the percentage of segments showing internal colonization; (2) M % (intensity of mycorrhization) indicating the average percentage of colonization of root segments; (3) a % (percentage of arbuscules), quantifying the average presence of arbuscules within the infected areas; (4) A % (percentage of arbuscules in the root system, quantifying the presence of arbuscules in the whole root system (Trouvelot *et al.*, 1986).

Grain yield and characteristics: At the time of maturity, yield data were recorded. For this purpose, plants of three random parcels of 15 m² (10 x 1.5 m), not close to the margins of the field, were collected for each treatment. After a manual separation of seed from the cob, grain was weighed and a subset of 500 g was stored in oven at 105°C for moisture content analysis. Grain yield was then corrected for the standard moisture (14%).

The chemical analyses of grain were carried out on samples from the eight different fields (a subset from ten replicates/field). Moisture, crude protein, ether extract, ash, crude fiber, neutral detergent fiber (NDF), acid detergent fiber (ADF), acid detergent lignin (ADL), non-structural carbohydrates (NSC), were performed according to AOAC methods (2002). NSC were calculated as a difference.

NIR and electronic nose analyses: In the present paper, radiometric analyses and olfactometry tests were carried out on 80 maize grain samples collected from the eight different fields (ten replicates/field) in the second cropping year.

Radiometric tests were performed by an ADS Field Spec HH PRO (Analytical Spectral Devices, Boulder, USA)

spectroradiometer equipped to collect spectra from 350 to 2500 nm (Siesler, 2008). A probe (ASD A122300 high-intensity reflectance hand-probe), served by external light source (2900 K color temperature quartz halogen light) to illuminate the object of interest, was used. The probe allows to collect reflectance spectra over an area 20 mm in diameter. For each sample, the spectrum was acquired directly from the Nallophan bag containing the maize grain. A panel covered by the Nallophan was used as white referencing. The spectra were collected with the Lab Spec Pro software 'sample spectrum count'.

Olfactometric tests were performed by a ten-MOS electronic nose Air Sense PEN2 (AIRSENSE, Analytics GmbH, Schwerin, Germany). This system intercepts the aromatic and volatile molecules released from the matrix. The air flux method was used in this trial. Data were acquired using a needle inserted into the Nallophan bag containing the grain. The sample run lasted 20 sec followed by a 40 sec flushing. Each measurement was checked and acquired by Win Munster v.1.6 software.

Mycotoxin analyses: Mycotoxin analyses were carried out on samples from the eight different fields (ten replicates/field) of the second cropping year. Mycotoxins (fumonisin, aflatoxin B1, zearalenone, deoxynivalenol) were determined by ELISA test through the automatic system Chemwell (Awereness Technology, Inc., Palm City; USA). The kits used were Ridascreen® Fumonisin (R 3401 R-Biopharm, Darmstadt, Germany); Ridascreen® Aflatoxin B1 30/15 (R 1211 R-Biopharm, Darmstadt, Germany); Ridascreen® Zearaleon (R 1401 R-Biopharm, Darmstadt, Germany); Ridascreen® DON (R 5906 R-Biopharm, Darmstadt, Germany).

Statistical analyses:

Root colonization: After having assessed normality by the Shapiro-Wilk test, differences between treatments were tested using the Kruskal-Wallis test within the NPAR1WAY procedure of SAS software (SAS 9.3 for Windows 2011), at statistical significance of 0.05.

Yield: Main effects (treatment, soil and irrigation) and their interactions were analyzed within the two years of experimentation and tested by F-test. The analysis of variance was performed using the GLM procedure of SAS software (2011).

Chemical characteristics: The effect of mycorrhization on chemical composition of grain was evaluated over the two different cropping years and mean were separated at 5% by t-student test (GLM procedure of SAS 9.3 for Windows 2011).

NIR data: All the spectra, consisting of 2151 points in the Vis-NIR radiation, were first submitted to a mathematical pre-treatment (Standard Normal Variate with Detrend, SNDV) and then the first derivative with gap 4 and smoothing 4 (1-4-4) was calculated. The statistics used for the equation development and evaluation were the standard error of cross validation (SECV), and the coefficient of determination in cross validation (1-VR). Analysis was performed using

WinISI II software, version 1.04, supplied by Infrasoft International (ISI, State College, PA, USA) according to Shenk and Westerhaus (1996).

E-nose data: The olfactometric measurement of the 10 MOS sensors registered during the first 20 sec were processed as contiguous vectors, into a kind of 200 points 'spectrum', also pre-treated with SNVD and 1-4-4. The same data analysis performed on NIR data was adopted.

RESULTS

Mycorrhizal root colonization: The results on mycorrhizal index are shown in Figure 1. The rate of mycorrhizal colonization was very high for both the treatments. F% and M% of the mycorrhizal colonization were higher in the maize inoculated respect to the control plants, 87.8 vs 80.3% and 26.8 vs 17.5%, respectively, ($P < 0.05$). A% was also higher in inoculated plants, 9.7 vs 5.8%, ($P < 0.05$). Assuming the effect of mycorrhizae were of interest in terms of hydro-geological severity, the comparison was performed under the most unfavorable condition (non-irrigated). PF soil was selected owing to its lighter and coarser texture than CB soil. Indeed, those characteristics are related to a lower water holding capacity.




Figure 1. Frequency of mycorrhiza (F%), intensity of colonization (M%), abundance of arbuscules in a root system (A%) and abundance of arbuscules in mycorrhized roots (a%) of maize inoculated with MICOSAT F® (M) and uninoculated (C) produced under field conditions in PF soil non-irrigated plots.

Arbuscules and vesicles observed in maize root fragments are shown in Figure 2. The arbuscules are highly branched hyphal structures filling the root cortical cells and represent the key structures of the symbiosis where most of nutrient exchange between fungi and plant occur (Sawers *et al.*, 2008). The vesicles are lipid storage organs.




Figure 2. Arbuscules and vesicles observed on maize roots produced under field conditions in PF soil and non-irrigated plots.

The root system of experimental maize plants is shown in Figure 3a and 3b. The higher mycorrhizal colonization found in inoculated maize seems not to have clearly affected the development of the root system that are not different at glance.




Figure 3a. Root systems of non-inoculated (C) maize produced under field conditions in PF soil and non-irrigated plots.



Figure 3b. Root systems of inoculated (M) maize produced under field conditions in PF soil and non-irrigated plots.

Grain yield and chemical characteristics: Table 1 shows maize grain yield. Mycorrhizal treatment did not affect yield in terms of maize grain, but yield of maize cultivated on CB soil plots was lower than PF soil over both the years ($P < 0.05$). No significant interaction was found between soil and mycorrhizal treatment. The effect of irrigation was not univocal, depending on the year. Therefore, in the first year no difference came out according to the water treatment while in the second year was higher under wet conditions ($P < 0.01$). Again, no meaningful interaction was found between irrigation and treatment and between soil and irrigation.

Table 1. Grain yield (t/ha) of maize inoculated (M) and uninoculated (C) produced under field conditions in two soils, and in dry and irrigated plots in two cropping years.

Year		2011	2012
Treatment	M	10.3	8.8
	C	10.1	9.3
	CB	8.7 <i>b</i>	6.7 <i>B</i>
Soil	PF	11.7 <i>a</i>	11.4 <i>A</i>
	no	10.8	8.1 <i>B</i>
Irrigation	yes	9.6	10.0 <i>A</i>

a, b: $P \leq 0.05$ A, B: $P \leq 0.01$

Table 2. Chemical characteristics on dry matter of maize inoculated (M) and uninoculated (C) produced under field conditions in two cropping years.

Year	Treatment	Dry matter (%)	Crude protein (%)	Ether Extract (%)	Ashes (%)	Crude fiber (%)	NDF (%)	ADF (%)	ADL (%)	NSC (%)
2011	M	95.37	7.73	3.93A	1.75a	3.86	16.79	4.56 a	1.43	69.80
	C	95.32	7.37	3.35B	1.62b	3.79	17.62	4.09 b	1.62	70.05
2012	M	97.54	7.63	2.41	1.08	4.06	14.75	4.59	1.14	74.13
	C	97.31	7.79	2.54	1.06	4.25	15.17	4.88	1.03	73.70

a, b: $P \leq 0.05$ A, B: $P \leq 0.01$

Table 2 shows the chemical composition of maize grain. Mycorrhizal treatment did not affect chemical characteristics of maize grain except for lipid ($P < 0.01$), ash ($P < 0.05$), and part of the fibrous fractions (ADF, $P < 0.05$), that resulted significantly higher in inoculated maize grain for the first cropping year. In both years NDF content, the highest fibrous component, tended to be lower in modulus in mycorrhizal maize grain, even though this difference was never statistically meaningful.

NIR and electronic nose: For NIRs data (Tab. 3), the mycorrhization factor gave a high coefficient of determination in cross validation (1-VR=0.86) whilst the influence of irrigation and soil factors were slightly lower (0.79).

Table 3. NIR cross-validation results of maize samples produced under field conditions.

Constituent	N	Mean	SD	SEC	RSQ	SECV	1-VR
Mycorrhization	72	1.51	0.50	0.08	0.98	0.19	0.86
Soil	72	1.49	0.50	0.09	0.97	0.23	0.79
Irrigation	74	1.49	0.50	0.11	0.95	0.23	0.70

On the contrary the analysis of the e-nose data (Tab. 4) showed a strong influence of the soil (0.75) rather than mycorrhization (0.63) but the influence of irrigation was very low (0.33).

Table 4. E-nose cross-validation results of maize samples produced under field conditions.

Constituent	N	Mean	SD	SEC	RSQ	SECV	1-VR
Mycorrhization	70	1.49	0.50	0.22	0.80	0.30	0.63
Soil	69	1.52	0.50	0.20	0.85	0.25	0.75
Irrigation	71	1.46	0.50	0.34	0.55	0.41	0.30

Mycotoxins: Preliminary analyses about the content of main mycotoxins in maize grain were performed (data not shown; Dr. Locatelli personal communication). The values of *Fumonisin* content did not change in mycorrhized samples when compared to controls. *Aflatoxin B1*, *Deoxynivalenol* (DON) and *Zearalenone* (ZEA) contents were close to the limits of detection in all samples, so comments about the content of these mycotoxins according to treatments cannot be rationally formulated.

DISCUSSION

Mycorrhizal root colonization: The commercial inoculum applied in the trial increased the AM colonization of maize roots although a considerable abundance of mycorrhizae in uninoculated plants was also found.

As a matter of fact, mycorrhizal index (F%) values registered in uninoculated maize were very high (80.3%), when compared with related studies. The AM colonization of uninoculated maize produced under field conditions varied according to different Authors: 39.7% in maize as following crop after *Trifolium alexandrinum* (Pellegrino *et al.*, 2011), 25.1-61.2% in two consecutive years (Sousa *et al.*, 2012), 45% (Cheeke *et al.*, 2013) and 55.3% (Berta *et al.*, 2014).

Autochthonous AM fungi can colonize roots of most plant species, including maize (Smith and Read, 2008), but it was clearly demonstrated that the soils exposed to intensive cultivation have extremely low number of indigenous AM fungal propagules (Li and Zhao, 2005; Subramanian *et al.*, 2009). Despite intensive agricultural practices have been applied along many years in the farm where this experimental trial was carried out, the frequency of mycorrhizae found in uninoculated plants remained high.

In general, intensive agricultural practices include monoculture, fertilizer, pesticide and herbicide applications and conventional tillage. The latter, considered a disturbing practice, is supposed to reduce the function of AM symbiosis through the breakdown of their hyphal network in the soil (Jasper *et al.*, 1989; Menendez *et al.*, 2001). However, as results from meta-analysis of numerous studies (Lekberg and Koide, 2005), conducted to evaluate the effects of different agricultural practices on mycorrhizal colonization, the control of disturbance slightly contributes to increase mycorrhizal colonization. Disturbance is supposed to mostly affect hyphae (Evans and Miller, 1988; Evans and Miller, 1990), whereas spores and colonized root pieces may still result infective.

The monoculture, as reported by Lekberg and Koide (2005), did not affect subsequent mycorrhizal colonization when compared with crop rotations including mycorrhizal plants.

A further factor to consider for mycorrhizal colonization is the concentration of available P in the soil. The experimental plots were well supplied with P content (Raglione *et al.*, 2011)

Under such conditions, the colonization of root by AM fungi is not favored and is often suppressed (Jensen and Jakobsen, 1980; Al-Karaki and Clark, 1999; Kahiluoto *et al.*, 2001).

M fungi from fertilized soil produced fewer hyphae and arbuscules and consequently supplied their host with fewer inorganic nutrients from the soil (Johnson, 1993).

Concerning the mycorrhizal colonization of inoculated plants of this trial, F (87.8%) was lower than the expected based on some similar available studies in field conditions (Berta *et al.*, 2014).

The significant, but not remarkable differences between the rates of root colonization in inoculated plants respect to uninoculated ones may be explicated considering the difficulty the AM inoculum has to face in soil colonization. In fact, the survival of the introduced microorganisms, as well as their activity on the host plant, are restricted by the competition with the resident microflora. (Hazarika *et al.*, 1999; Zhang *et al.*, 2011).

Root development, grain yield and chemical characteristics of maize: The root systems of inoculated and uninoculated maize showed a comparable size, as reported in the pictures (Fig. 3a and 3b). These results disagree with those of other Authors (Berta *et al.*, 2014; Subramanian *et al.*, 2014), who found a higher length and volume of inoculated maize roots. The MICOSAT F® inocula did not affect maize grain yield for the two years. Even though mycorrhizal frequency is not always a good predictor of plant productivity, results of meta-analysis (Lekberg and Koide, 2005) showed that the increase of mycorrhizal colonization is related with some benefits for plants. The lack of effect of inoculum on grain yield could be due to: a) small difference in mycorrhizal colonization between inoculated and uninoculated maize; b) high soil available P content; c) inefficiency of the markers used to evaluate the advantage of MICOSAT F® inoculum in specific soil and climate conditions. A recent study (Sabia *et al.*, 2015) revealed a significant effect of AM seed inoculation on yield of forage maize cultivated within a low input system, including low P input.

Regarding the different soils used in this trial, grain yield of PF resulted higher in both the years respect to that of CB soil. The characteristics of the CB soil, *i.e.* high clay content, compactness and tendency to surface fissuring, make it less suitable for maize cropping (Passioura *et al.*, 1991).

Regarding the irrigation, the different weather conditions over the two years may justify the results: no significant difference between dry and irrigated cropping in 2011 and significantly higher yield for irrigated crop in 2012. The period between June and August 2011 was characterized by abundant average rainfall (41.6 mm) and mild average temperature (24.1°C). In the same period of 2012 only 4.7 mm of rain fell and the average temperature was 25.9°C. The lack of irrigation effect on grain yield of maize cultivated in 2011 could have been due to the rainy and damp weather conditions over that cropping period. On the other hand, 2012 was much drier and

although only an emergency sprinkling irrigation had been performed, the effect was evident.

The absence of significant interaction between mycorrhizal colonization and soil and between mycorrhizal colonization and irrigation suggests that the used inoculum did not improve the maize productivity in soil with different texture and different attitude to drought.

Concerning the grain chemical characteristics, a different response was obtained according to the year. In 2011 the higher ether extract, ADF and ash content registered in grain of mycorrhized maize is consistent with data presented by other Authors (Subramanian *et al.*, 2014; Al-Karaki *et al.*, 1998).

We should stress a scarce impact of MICOSAT F and the major effect of the cropping year on maize production and quality. Perhaps, owing to the drier weather in 2012 when compared to 2011, the balance of the competition between plants and between plant and free-living microorganisms (Rennenberg *et al.*, 2009) may result modified, leading to a reduction of efficiency of MICOSAT F in promoting plant's absorption of specific minerals.

Nevertheless, the NIRs analysis of grain of the second cropping year allowed to clearly separate between mycorrhized and not mycorrhized even though, in this case, no differences were highlighted by the chemical analysis of proximate composition. That last event confirms multifactor approaches, and particularly NIRs and multivariate analysis, as strong tool for discriminating complex matrices. As a matter of fact, NIRs, thanks to the detection of the overtones and combination bands that are specific for different functional groups, besides the fundamental transition from the ground to the excited state of the molecules (harmonic vibration model), can give a great deal of additional information. Therefore, NIR spectra are very powerful tool for predictive as well as discriminant analyses.

As for the electronic nose, soil factor affected the volatile compound of maize more than mycorrhizal inoculum. An electronic nose is designed to recognize different species in the headspace and its selectivity is limited. The response is often subjected to noise, drift, and signals pertaining to other factors and pre- and post-treatments are required in order to minimize those effects (Sheebha and Subadra, 2014; Al-Maskari *et al.*, 2014; Hines *et al.*, 1999). The stronger and/or confusing effects of other factors such as soil and water regimen may have veiled the effect of VAM inoculation. Since mycorrhizae are supposed to change volatile compounds production in plants (Fontana *et al.*, 2009; Leitner *et al.*, 2010), differences could be isolated and appreciated under strictly controlled conditions at greenhouse or pots, where the only involved factor is inoculation.

Conclusions: The inoculum (MICOSAT F®) increased the AM colonization of maize roots even though it probably underwent to a strong competitive pressure by the

autochthonic microflora that limited its performances and vitality. Indeed, a considerable number of mycorrhizae was detected in uninoculated plants as well. The NIRs indicated some physical differences between inoculated and uninoculated samples and resulted efficient for characterization of mycorrhized maize grain. On the other hand, e-nose's response is more affected by soil conditions than by mycorrhizal colonization. Further in-depth studies on AM fungi physiology, their function and the interactions with crops and environmental conditions are required to thoroughly explain these results. Compounds involved in the expression of inoculation's effect as well as each specific role must be characterized. Eventually, plants should be tested in different soils and conditions to clarify the effect of MICOSAT F® on maize yield and composition for food and feedstuff production.

Acknowledgment: This study was funded by the Italian Ministry of Agriculture within the project "Sustainable livestock models" (SOSZOOT).

REFERENCES

- Al-Karaki, G.N., A. Al-Raddad and R.B. Clark. 1998. Water stress and mycorrhizal isolate effect on growth and nutrient acquisition of wheat. *J. Plant Nutr.* 21:891-902.
- Al-Karaki, G.N. and R.B. Clark. 1999. Varied rates of mycorrhizal inoculum on growth and nutrient acquisition by barley grown with drought stress. *J. Plant Nutr.* 22:775-1784.
- Al-Maskari, S., X. Li and Q. Liu. 2014. An Effective Approach to Handling Noise and Drift in Electronic Noses. In: H. Wang and M.A. Sharaf (eds.), *Databases Theory and Applications. Proc. 25th Australasian Database Conference, Brisbane, Australia*. pp.223-230.
- AOAC. 2002. *Official Methods of Analysis*, 17th Ed. Association of Official Analytical Chemists, Washington, USA.
- Berta, G., A. Copetta, E. Gamalero, E. Bona, P. Cesaro, A. Scarafoni and G. D'Agostino. 2014. Maize development and grain quality are differentially affected by mycorrhizal fungi and a growth-promoting pseudomonad in the field. *Mycorrhiza* 24:61-170.
- Candido, V., G. Campanelli, T. D'Addabbo, D. Castronuovo, M. Rencoand and I. Camele. 2013. Growth and yield promoting effect of artificial mycorrhization combined with different fertiliser rates on field-grown tomato. *Ital. J. Agron.* 8:168-174.
- Celebi, S.Z., S. Demir, R. Celebi, E.D. Durak and I.H. Yilmaz. 2010. The effect of Arbuscular Mycorrhizal Fungi applications on the silage maize (*Zea Mays* L.) yield in different irrigation regimes. *Eur. J. Soil Biol.* 46:302-305.
- Cheeke, T.E., M.B. Cruzan and T.N. Rosenstiel. 2013. Field evaluation of arbuscular mycorrhizal fungal colonization in *Bacillus thuringiensis* toxin-expressing (Bt) and non Bt maize. *Appl. Environm. Microbiol.* 13:4078-4086.
- Covacevich, F., H.E. Echeverria and L.A.N. Aguirrezabal. 2007. Soil available phosphorus status determines indigenous mycorrhizal colonization of field and glasshouse-grown spring wheat from Argentina. *Appl. Soil Ecol.* 35:1-9.
- Davis, E.A., J.L. Young and R.G. Linderman. 1983. Soil lime level (pH) and VA-mycorrhiza effects on growth responses of sweet gum seedlings. *Soil Sci. Soc. Amer. J.* 47:251-257.
- Douds, D.D. Jr., G. Nagahashi, C. Reider and P. Hepperly. 2007. Inoculation with am fungi increases the yield of potatoes in a high P soil. *Biol. Agric. Hortic.* 25:67-78.
- Evans, D.G. and M.H. Miller. 1988. VA-mycorrhiza and the soil disturbance induced reduction of nutrient absorption in maize. I. Causal relations. *New Phitol.* 110:67-75.
- Evans, D.G. and M.H. Miller. 1990. The role of extra-radical mycelial network in the effect of soil disturbance upon vesicular-arbuscular mycorrhizal colonization of maize. *New Phitol.* 114:65-71.
- Fontana, M., S. Reichelt, J. Hempel, S.B. Gershenzon and S.B. Unsicker. 2009. The effects of arbuscular mycorrhizal fungi on direct and indirect defense metabolites of *Plantago lanceolata* L. *J. Chem. Ecol.* 35:833-843.
- George, E. 2000. Nutrient uptake. In: H. Koltai, Y. Kapulnik (eds.), *Arbuscular Mycorrhizas: Physiology and function*. Springer, Berlin. pp.307-344.
- Gilani, R.H., M.A. Mian, S. Muhammad and A. Ghani. 1983. The fertilizer phosphate utilization efficiency in wheat production. *Pak. J. Soil Sci.* 1:7-9.
- Gosling, P., A. Hodge, G. Goodlass and G.D. Bending. 2006. Arbuscular mycorrhizal fungi and organic farming. *Agric. Ecosys. Environm.* 113:17-35.
- Günes, A., M. Turan, M. Güllüce and F. Sahin. 2014. Nutritional content analysis of plant growth-promoting rhizobacteria species. *Europ. J. Soil Biol.* 60:88-97.
- Gyaneshwar, P., G.N. Kumar, L.J. Parekh and P.S. Poole. 2002. Role of soil microorganisms in improving P nutrition of plants. *Pl. Soil* 245:83-93.
- Hazarika, D.K., K.K. Das and L.N. Dubey. 1999. Effect of vesicular arbuscular mycorrhizal fungi inoculation on growth and nutrient uptake of black gram. *J. Mycol. Plant Pathol.* 29:201-204.
- Hao, X., C.M. Cho, G.J. Racz and C. Chang. 2002. Chemical retardation of phosphate diffusion in an acid soil as affected by liming. *Nutr. Cycl. Agroecosyst.* 64:213-224.
- Hines, E.L., E. Llobet and J.W. Gardner. 1999. Electronic noses: A review of signal processing techniques. *IEE Proc. Circuits, Devices and Systems* 146:297-310.

- Isherword, K.F. 1998. Fertilizer use and environment. In: N. Ahmed and A. Hamid (eds.), Proc. Symp. Plant Nutrition Management for Sustainable Agricultural Growth. NFDC, Islamabad, Pakistan; pp.57-76.
- Jasper, D.A., L.K. Abbott and A.D. Robson. 1989. Soil disturbance reduced the infectivity of external hyphae of vesicular-arbuscular mycorrhizal fungi. *New Phytol.* 33:373-381.
- Jensen, A. and I. Jacobsen. 1980. The occurrence of vesicular-arbuscular mycorrhiza in barley and wheat grown in some Danish soils with different fertilizer treatments. *Pl. Soil* 55:403-414.
- Johnson, N.C. 1993. Can fertilization of soil select less mutualistic mycorrhizae? *Ecol. Applic.* 3:749-757.
- Kahiluoto, H., E. Ketoja, M. Vestberg and I. Saarela. 2001. Promotion of AM utilization through reduced P fertilization. 2. Field studies. *Pl. Soil* 231:65-79.
- Krey, T., N. Vassilev, C. Baum and B. Eichler-Löbermann. 2013. Effects of long-term phosphorus application and plant-growth promoting rhizobacteria on maize phosphorus nutrition under field conditions. *Eur. J. Soil Biol.* 55:124-130.
- Leitner, M., R. Kaiser, B. Hause, W. Boland and A. Mithofer. 2010. Does mycorrhization influence herbivore-induced volatile emission in *Medicago truncatula*? *Mycorrhiza* 20:89-101.
- Lekberg, Y. and R.T. Koide. 2005. Is plant performance limited by abundance of arbuscular mycorrhizal fungi? A meta-analysis of studies published between 1988 and 2003. *New Phytol.* 168:189-204.
- Li, T. and Z.W. Zhao. 2005. Arbuscular mycorrhizas in a hot and arid ecosystem in southwest China. *Appl. Soil. Ecol.* 29:135-141.
- Menendez, A.B., J.M. Scervino and A.M. Godeas. 2001. Arbuscular mycorrhizal populations associated with natural and cultivated vegetation on a site of Buenos Aires province, Argentina. *Biol. Fertil. Soils* 33:373-381.
- Nadeem, S.M., M. Ahmadb, Z.A. Zahir, A. Javaid and M. Ashraf. 2014. The role of mycorrhizae and plant growth promoting rhizobacteria (PGPR) in improving crop productivity under stressful environments. *Biotech. Advanc.* 32:429-448.
- Novero, M., A. Faccio, A. Genre, J. Stougaard, K.J. Webb, L. Mulder, M. Parniske and P. Bonfante. 2002. Dual requirement of the LjSym4 gene for mycorrhizal development in epidermal and cortical cells of *Lotus japonicus* roots. *New Phytol.* 154:741-749.
- Oberson, A., D.K. Friesen, I.M. Rao, S. Buhler and E. Frossard. 2001. Phosphorus transformations in an oxisol under contrasting land-use systems: the role of the soil microbial biomass. *Pl. Soil* 237:197-210.
- Passioura, J.B. 1991. Soil structure and plant growth. *Austr. J. Soil Res.* 29:717-728.
- Pellegrino, E., S. Bedini, L. Avio, E. Bonari and M. Giovannetti. 2011. Field inoculation effectiveness of native and exotic arbuscular mycorrhizal fungi in a Mediterranean agricultural soil. *Soil Biol. Biochem.* 43:367-376.
- Raglione, M., P. Lorenzoni, A. Bonifazi, R. Nannelli, E. Bagnarli, P. Bazzoffi and R. Gentile. 2011. Effectiveness of the GAEC cross compliance standard Rational management of set aside (weed control through mowing) in assuring a minimum level of habitats maintenance. *Ital. J. Agron.* 6:92-99.
- Rennenberg, H., M. Dannenmann, A. Gessler, J. Kreuzwieser, J. Simon and H. Papen. 2009. Nitrogen balance in forest soils: nutritional limitation of plants under climate change stresses. *Plant Biol.* 11:4-23.
- Richardson, A.E. and R.J. Simpson. 2011. Soil microorganisms mediating phosphorus availability. *Plant Physiol.* 156:989-996.
- Sabia, E., S. Claps, G. Morone, A. Bruno, L. Sepe and R. Aleandri. 2015. Field inoculation of arbuscular mycorrhiza on maize (*Zea mays* L.) under low inputs: preliminary study on quantitative and qualitative aspects. *Ital. J. Agron.* 10:607-610.
- SAS Institute Inc. 2011. SAS/STAT® 9.3 User's Guide. Cary, USA.
- Shenk, J.S. and M.O. Westerhaus. 1996. Calibration the ISI way. In: A.M.C. Davies and P. Williams (eds.), *Near Infrared Reflectance Spectroscopy: The Future Waves*. NIR Publications, Chichester; pp.198-202.
- Siesler, H.W. 2008. Basic principles of near-infrared spectroscopy. In: D.A. Burns and E.W. Ciurczak (eds.), *Handbook of Near Infrared Analysis*. CRC Press, Taylor & Francis Group, New York. pp.7-20.
- Sheebha, R. and M. Subhadra. 2014. A survey on various signal pre-processing methods for sensor response. *E-Nose. Researcher.* 6:76-80.
- Smith, S.E. and D. Read. 2008. The symbionts forming arbuscular mycorrhizas. In: S.E. Smith and D. Read (eds.), *Mycorrhizal Symbiosis*. Academic Press, New York. pp.13-41.
- Subramanian, K.S., V. Tenshia, K. Jayalakshmi and V. Ramachandran. 2009. Biochemical changes and zinc fractions in arbuscular mycorrhizal fungus (*Glomus intraradices*) inoculated and uninoculated soils under differential zinc fertilization. *Appl. Soil Ecol.* 43:32-39.
- Subramanian, K.S., C. Bharathi, M. Gomathy and N. Balakrishnan. 2014. Role of arbuscular mycorrhizal (*Glomus intraradices*) fungus inoculation on Zn nutrition in grains of field-grown maize. *Aust. J. Crop Sci.* 8:655-665.
- Trouvelot, A., J.L. Kough and V. Gianinazzi-Pearson. 1986. Mesure du taux de Mycorrhization VA d'un système racinaire. Recherche de methods d'estimation ayant une signification fonctionnelle. In: V. Gianinazzi-Pearson

- and S. Gianinazzi (eds.), Physiological and Genetic Aspects of Mycorrhizae. INRA, Paris. pp.217-221.
- Vassilev N., M. Vassileva and I. Nikolaeva. 2006. Simultaneous P-solubilizing and biocontrol activity of microorganisms: potentials and future trends. Appl. Microbiol. Biotechnol. 71:137-144.
- Zhang, G.Y., L.P. Zhang, M.F. Wei, Z. Liu, Q.L. Fan, Q.R. Shen and G.H. Xu. 2011. Effect of arbuscular mycorrhizal fungi, organic fertilizer and soil sterilization on maize growth. Acta Ecol. Sinica 31:192-196.
- Zoppellari, F., E. Malusà, W. Chitarra, C. Lovisolo, F. Spanna and L. Bardi. 2014. Improvement of drought tolerance in maize (*Zea mays* L.) by selected rhizospheric microorganisms. Ital. J. Agromet. 19:5-18.