

POTASSIUM FERTILIZATION MAY IMPROVE STEM STRENGTH AND YIELD OF BASMATI RICE GROWN ON NITROGEN-FERTILIZED SOILS

Umar Zaman^{1,†}, Zeeshan Ahmad^{1,†}, Muhammad Farooq^{2,3}, Salmaan Saeed⁴, Mumtaz Ahmad⁵ and Abdul Wakeel^{1,6,*}

¹Institute of Soil and Environmental Sciences, University of Agriculture, Faisalabad-38040, Pakistan; ²Department of Agronomy, University of Agriculture, Faisalabad-38040, Pakistan; ³College of Food and Agricultural Sciences, King Saud University, Riyadh 11451, Saudi Arabia; ⁴Food and Biotechnology Research Center, PCSIR Labs. Complex, 54600 Lahore; ⁵Engro Eximp (Pvt.) Ltd. DeraSarawan, Sakham, Sheikhpura, Pakistan; ⁶International Potash Institute, Baumgärtlistrasse 17, CH-8810 Horgen, Switzerland

[†]Both authors contributed equally

*Corresponding Author: abdulwakeel77@gmail.com

Imbalance use of fertilizers is one of the important factors for lodging of rice leading to paddy-yield reduction. Potassium (K) deficiency may be one of the contributing factors in this regard. Present study was conducted to assess the decrease in lodging losses due to use of K fertilizer in the presence of nitrogen by determining various factors contributing to stem strength. Two levels of K, one with deficient (K_d) while other with recommended (K_r) K and three levels of N *i.e.* deficient (N_d), half of recommended (N_{hr}) and recommended (N_r) N were applied along with recommended dose of phosphorus and zinc to two fine-grain rice cultivars *i.e.* Super basmati and Basmati 515. The experiment was laid out in randomized complete block design in factorial arrangement with four replicates. Application of K fertilizer significantly improved K concentration in rice stem, and paddy yield of Basmati rice. Furthermore, K fertilization improved plant parameters such as basal internode space, cellulose content and stem-fiber contents. Increase in silicon content in stem due to application of K was also observed, which may also contribute to stem strength against environmental stresses. The present study suggests further work to understand the physiological and biochemical mechanisms contributing to stem strength in response to K fertilization.

Keywords: potassium, balanced fertilization, stem strength, nitrogen, lodging

INTRODUCTION

Rice is the second most important staple crop after wheat in Pakistan and contributes greatly to food security and foreign exchange. Area under rice cultivation in Pakistan is about 2.78 million hectares, giving 6.79 million tons annual production with an average of 2.44 tons ha⁻¹ (Anonymous, 2014). Pakistan is at number eleven in total rice production (FAO, 2012) however fine grained Basmati rice of Pakistan is famous for its quality, taste and aroma throughout the world. Therefore, rice is an important source of income for majority of the farmers in the Punjab and Sindh provinces of Pakistan. Different factors causing low rice production in Pakistan include water scarcity, high input prices, unavailability of skilled labour, less plant population, weeds and pest infestation, lodging and marketing issues (Baloch *et al.*, 2004). Lodging causes severe damage to yield and quality of grain and forage in cereals especially in rice. Broken rice plant stem disturbs the vascular tissues, lowers the photosynthetic rate, increases respiration, and reduces nutrients translocation for grain filling and other plant needs (Tripathi *et al.*, 2003). Lodging also increases disease attack

and reduces pest resistance. Mechanical harvesting problems are also associated with lodging (Kono, 1995) and have great significance in modern agriculture systems.

Generally crops are more susceptible to lodging near to its maturity and is often caused by rain, wind and hailing. Its distribution may often not be uniform throughout the field, may be in spots or scattered over different field sections (Berry *et al.* 2004). Lodging reduces cereals yield especially under mechanized harvesting (Rajkumara, 2008) and restricts the opportunity of increase in rice yield even by additional fertilizer application. Imbalance and limited use of potassium (K) fertilizers may weaken the stem strength making it more susceptible to lodging (Chang, 1964).

Potassium stands on third in macronutrients after nitrogen (N) and phosphorus (P) and is the most abundant macronutrient in soil. Its availability to plants depends upon the concentration of K in soil solution and on exchange sites. Potassium is essential for enzyme activation, charge balance and osmotic regulation in plants (Wakeel *et al.*, 2011; Zörb *et al.*, 2014). It is present in cation form in plant cell to maintain ionic balance and up to 10% of plant dry matter is made up of K (Marschner, 1995). Its ease of access to plants

depends upon its status in soil solution, exchangeable K and rate of exchange of K from exchangeable form to soil solution form (Wakeel, 2013). Specific functions that occur in cytoplasm require small amount of potassium while a major portion (90%) of K is present in vacuole where it acts as an osmoticum contributing to extension growth of plants (Wakeel *et al.*, 2010, 2011).

High K contents in plant tissue increase its resistant to sudden environmental variations like temperature stress, rain in late season, cold and frost (Chamak, 2005). Potassium accumulation in the plant tissue also decreases damage to plant in response to osmotic stress and physiological burdens (Rajkumara, 2008). There is a close interaction between K content of the lower part of culm and lodging resistance in stem. Appropriate K nutrition is closely related to lignification of sclerenchyma cells, vascular bundles and stem strength to enhance lodging resistance (Kong *et al.*, 2014).

Increased use of N improves the vegetative growth and height increasing its susceptibility to lodging and decreased K fertilization weakened the plant stem. In Pakistan, use of K fertilizers is neglected; however, N application is increasing and N: K use ratio is going to increase continuously (NFDC, 2013-14). It was hypothesized that application of K fertilizer under N fertilized conditions may improve stem strength and yield of Basmati rice. This study was conducted to determine the potential role of K fertilization in stem strength, lodging resistance and paddy yield of Basmati rice in major rice growing area of Pakistan.

MATERIALS AND METHODS

Study was conducted during July-November, 2012 at Research Station of the Engro Eximp. (Private) Limited Dera Sarawan near Sakham Muridkay, Sheikhpura. The experimental soil was clay loam and is located in rice belt of Punjab, Pakistan. Physicochemical properties (Estefane *et al.*, 2013; Table 1) show that soil was deficient in K and N nutrition.

Table 1. Physicochemical properties of experimental soil

Soil Characteristics	Values
Soil texture	Clay loam
pH	8.0
EC	0.30 dS m ⁻¹
Exchangeable sodium	1.7 %
Organic matter	0.81 %
Nitrogen	0.051 %
Phosphorus (Olsen)	6 mg kg ⁻¹
Exchangeable potassium	86 mg kg ⁻¹
Zinc	1.08 mg kg ⁻¹
Boron	0.65 mg kg ⁻¹
Iron	8.75 mg kg ⁻¹

Crop husbandry: The study was carried out in field using randomized complete block design in factorial arrangement with four replicates. Seeds of two rice cultivars Super basmati and Basmati 515 were soaked in water for two days before nursery sowing. Seed rate was 12.5 kg for one hectare nursery sown on 15 June. Field was irrigated followed by puddling and planking and was repeated with an interval of one week to prepare field for nursery transplanting. Seedbed preparation was started keeping nursery age in view. Thirty days old nursery seedlings were transplanted manually on 15th of July. Potassium and nitrogen fertilizers were applied as deficient K (K_d) and deficient N (N_d) as (control), deficient K (K_d) and recommended N (N_r) *i.e.* 120 kg ha⁻¹ N, recommended K (K_r) *i.e.* 60 kg ha⁻¹ K and half recommended N (N_{hr}) *i.e.* 60 kg ha⁻¹ N, and K_r and N_r; whereas phosphorus was applied at the rate of 60 kg ha⁻¹ at seed bed preparation in the form of single super phosphate. Zinc was applied three weeks after the nursery transplantation at 9 kg ha⁻¹. Cartap (Cartap Hydrochloride) was applied at 22.5 kg ha⁻¹ on 20th August and repeated on first week of September to control insect-pests, especially the rice stem borer and rice leaf folder. In second week of September Nativo (trifloxystrobin and tebuconazol) was sprayed to cure crop from fungal attack. During 1st week of October, Actara and Score were sprayed to prevent the plant hopper.

Last irrigation was applied on 15th October 2012. Crop was harvested on 11th of November 2012 when the grain moisture content was ~19%. After manual harvesting, plant height, basal internode space, number of grains per panicle, paddy yield, acid detergent fiber concentration, cellulose concentration, and concentration of K, N and Si were determined.

Biochemical analysis: Plant samples were digested following McGill and Figueiredo (1993). Half gram of dried and ground rice shoot sample was taken in each of 100 mL Pyrex digestion tube. Five milliliter of concentrated H₂SO₄ was added and allowed to predigest overnight in a fume hood. Next day, the contents of digestion tube were swirled; 2 mL of 35 % H₂O₂ was added in each tube and allowed to cool. Cooled tubes were heated on block digester at 150 C°. Cooled the tubes again and 2 mL of 35 % H₂O₂ was added. Temperature was raised to 380 C° and samples were digested for 10 min. The above step was repeated until digest remained clear/transparent. The digested samples were diluted to 50 mL volume and stored in stoppered plastic bottles for the estimation of N, P and K concentration.

Nitrogen determination: Nitrogen contents were measured in this digest by distillation method as described by Bremner and Mulvaney (1982). Ten mL of aliquot were pipetted out into a 100 mL distillation flask and distilled for 90 minutes with 10 mL of 12 N NaOH. The distillate was collected in receiver flask containing 5 mL of mixed indicator 2 % boric acid solution. The distillation time of 90 sec. allowed 35 mL

distillate collection. The distillate was titrated against 0.01 N H₂SO₄ and nitrogen was calculated as follow;

$$N (\%) = \frac{R - B}{V} \times 14.01 \times W_t$$

Where, R = Ratio between total digest volume and distillation volume, B = Digested blank V= Titration volume (mL), W_t = Mass of dry plant (g), 14.01=Constant

Potassium determination: Potassium was determined by flame photometer according to the method described by Chapman and Pratt (1961). The instrument was calibrated using a series of K standard solutions and emission of unknown sample was taken. The concentration of K in test samples was determined using regression equation drawn between standard solution concentrations and their emission. The corrected K concentration was calculated as follow;

$$K (mg\ kg^{-1}) = (K\ in\ extract - K\ in\ blank) \times A/W_t$$

Where, A = Total volume of the digest (mL),

W_t = Mass of plant material taken for digestion (g)

Acid detergent fiber (ADF) and cellulose determination:

Acid detergent fiber, cellulose, fiber and silica determination is done by sequence wise procedure which is called acid detergent fiber method and is proposed by Goering and Soest (1970). By this method in first step we find acid detergent fiber which is constituent of cellulose, lignin and finally silica and then from this material cellulose, lignin and finally silica is determined. One gram rice plant sample was taken in 250 mL conical flask. One hundred mL acid detergent was added in flask containing sample. The flask was heated on hot plate for one hour and filtered it giving two washings with hot boiling water followed by one washing of acetone. Residue was collected in previously tarred dry crucible. Then the residue was dried for 1 hour at 100- 105 °C for constant weight.

$$ADF (\%) = \frac{\text{Loss in weight after acid detergent treatment}}{\text{Weight of sample}} \times 100$$

Five milliliter 72 % H₂SO₄ was added in crucible containing ADF. Paste was made of this material and then crucible was covered and placed for one hour. After one hour again 5 mL 72 % H₂SO₄ was added and stirred thoroughly and let the

crucible again for one hour. 72 % H₂SO₄ was added again, stirred and let it for one hour. Material was filtered giving two washings distilled water to make material acid free. Checked the material by phenolphthaline and NaOH solution either it is acid free or not. Final washing was done by acetone to make it water free. Residue was taken in china dish, placed china dish in oven at 100 °C for one hour. Cooled and weighed the china dish, the loss in mass was cellulose.

Cellulose (%) = W_t loss by digestion/Mass of sample × 100
Statistical analysis: Means of four replications were taken for each treatment and statistical significance was calculated by ANOVA using Statistix 8.1 (Analytical Software, 2005) and linear correlation was plotted using MSTAT.

RESULTS

Plant height and number of grains per panicle: Basmati 515 genetically is high stature than the Super basmati, and N fertilization further increased plant height of both genotypes but more in Basmati 515 (Table 2). Maximum (130.2 cm) was in N_rK_r in Basmati515, whereas maximum plant height in Super basmati was 110 cm. Number of grains per panicle has significant contribution towards paddy yield. More the number of grains panicle⁻¹, more will be the paddy yield if the grain weight is similar. Fertilizer treatments and cultivars showed significant difference in number of grains per panicle. More number of grains (175 per panicle) was produced with N_rK_r than the other treatments, while Basmati 515 produced more number of grains per panicles than Super basmati (Table 2). Number of grains per panicle ranged between 110 to 134 and 127 to 175 in Super basmati and Basmati 515, respectively.

Basal internode space and paddy yield: There was no significant difference between Super basmati and Basmati 515 regarding basal internode spaces and minimum basal internode space (10.77, 10.68 cm) was recorded in N_r K_r in both genotypes while maximum basal internode length (8.47 cm) was recorded with N_dK_d in Super Basmati (Fig 1A). The ultimate objective of rice crop is to get paddy yield. Super basmati produced paddy better than basmati 515

Table 2. Effect of N and K fertilization on plant height, number of grains panicle⁻¹, shoot-ADF concentration, and shoot-cellulose concentration of two rice genotypes

Treatment	Plant height (cm)		Grains per panicle		ADF (%)		Cellulose (%)	
	Super Basmati	Basmati 515	Super Basmati	Basmati 515	Super Basmati	Basmati 515	Super Basmati	Basmati 515
N _d K _d	96.52±1.2f	105.62±1.2e	110.50±4.8f	127±1.9cde	42.26±1.16cd	37.16±2.9d	34.25±1.54bc	32.85±1.2c
N _{hr} K _d	104.33±1.9e	115.08±1.9bed	117± 2.4ef	132.7±2.9bc	49.11±1.8bc	46.16±3.3c	39.36±1.5abc	38.14±2.2abc
N _r K _d	110.33±1.3cde	121.08±1.3b	125.75±2.6def	140±3.4b	50.04±2.2bc	44.56±2.1cd	41.19±2.7ab	37.10±1.2abc
N _{hr} K _r	107.08±2.33e	117.25±2.33bc	131±1.7def	153±5.7b	44.58±1.70cd	46.71±2.81c	34.92±1.5bc	39.87±1.8abc
N _r K _r	108.75±3.14de	130.92±3.14a	134.5±4.09bcd	175±5.4a	57.65±2.8a	55.96±3.8ab	43.84±2.3a	44.45±4.1a

Five treatments (N_dK_d= no N and no K, N_{hr}K_d = 60 kg ha⁻¹ N and no K, N_rK_d = 120 kg ha⁻¹ N and no K, N_{hr}K_r = 60 kg ha⁻¹ N and 60 kg ha⁻¹ K, N_rK_r = 120 kg ha⁻¹ N and 60 kg ha⁻¹ K)

because of more number of tillers in super basmati. Maximum 5.45 tons ha⁻¹ and minimum 4.31 tons ha⁻¹ paddy yield were recorded in N_rK_r and N_dK_d respectively (Fig. 1B).




Figure 1. Effect of N and K fertilization on (A.) Internode space of two rice genotypes (B.) paddy yield per hectares. Five treatments N_dK_d= no N and no K, N_{hr}K_d = 60 kg ha⁻¹ N and no K, N_rK_d = 120 kg ha⁻¹ N and no K, N_{hr}K_r = 60 kg ha⁻¹ N and 60 kg ha⁻¹ K, N_rK_r = 120 kg ha⁻¹ N and 60 kg ha⁻¹ K. Column shows mean values of four replicates and bar shows the standard error of means. Column labeled with different alphabets show significant difference according to LSD test at 5% probability.

Acid detergent fiber and cellulose concentrations:

Cellulose is high molecular weight compound in plant stem and it plays important role in stem strengthening. It has largest proportion in acid detergent fiber than that of silicon and lignin. Maximum cellulose concentration (44.1%) was determined in N_rK_r while minimum (33.5%) was with N_dK_d (Table 2). Cellulose content of rice plant was increased with increasing the dose of K and N fertilizers. However, when N was applied without K the cellulose concentration was less. Maximum acid detergent fiber content (56.5%) was produced with N_rK_r while minimum (40%) was at N_dK_d (Table 2). Lignin concentration in shoot was not significantly different in response to K or/and N fertilization

(data not shown).

Shoot N, Si and K concentrations: Maximum N concentration (0.98 mg g⁻¹ dry weight) was recorded in N_rK_r (Table 3). Super Basmati showed more N concentration than the Basmati 515. Potassium fertilization has also synergistic effect on N uptake by rice but major effect of increased N concentration was due to N fertilization.

Generally, silicon (Si) is non-essential element for plants but it is beneficial for plants and play role in many important functions including stem strengthening. Maximum Si concentration (60.1 mg g⁻¹) was recorded in N_rK_r or Basmati 515 and minimum (21.9mg g⁻¹) was in N_dK_d (Table 3) while in Super basmati maximum Si was observed in N_{hr}K_r treatment *i.e.* 57.6 mg g⁻¹. Maximum K ion concentration was determined in shoot at treatment N_rK_r which may show synergistic effect of K on Si uptake by plant in case of Basmati 515.

DISCUSSION

Application of K fertilizer plays role in osmoregulation, maintaining turgor pressure in cell, cell elongation, and growth consequently (Marschner, 1995). In our results, the application of potassium and nitrogen increased plant height. The increase in plant height may be due to the increased amount of N fertilizer because it increases the plant vegetative growth by enhancing cell division (Haghighi *et al.*, 2011). Potassium fertilization also improve the plant height by improving its metabolism, however K reduced the growth rate and strengthen the stem. Increase in height by application of N as well as K fertilizers (Table 2) is similar to Shakouri *et al.* (2012) who reported that higher fertilizer rate increases plant height.

Nitrogen and K fertilizer increased grain yield significantly by improving photosynthetic rate and grain health (Bahmaniar and Ranjbar, 2007). Application of K with N plays a supportive role for N functioning and causes increase in yield production (Usherwood and Segars, 2001). The increase in yield of rice by application N and then further increase by application of K fertilizer may be due to the positive interaction of nitrogen and potassium as described

Table 3. Effect of K and N fertilization on potassium, silicon and nitrogen concentration (mg g⁻¹) in shoot of rice cultivars

Treatment	Silicon(mg g ⁻¹)		Potassium (mg g ⁻¹)		Nitrogen (mg g ⁻¹)	
	Super Basmati	Basmati 515	Super Basmati	Basmati 515	Super Basmati	Basmati 515
N _d K _d	11.6±0.1d	11.8±0.2d	19.4±3.1e	21.9 ± 0.6e	0.76±0.02e	0.69 ±0.03f
N _{hr} K _d	12.7± 0.1c	12.1±0.1d	42.6±2.4bc	27.7 ± 2.4de	0.84±0.01cd	0.79±0.01de
N _r K _d	13.0±0.2bc	12.4±0.2bc	41.2±1.4b	44.6 ± 1.2cd	0.98±0.01a	0.87±0.02bc
N _{hr} K _r	14.9±0.2a	14.6± 0.2a	57.6±2.8a	47.3 ± 0.6a	0.8±0.02bc	0.81±0.02e
N _r K _r	15.0 ±0.1a	14.7±0.3a	54.8 ± 3.7a	60.1 ± 3.7a	1.03±0.03a	0.89±0.02b

Values show the mean of four replication plus standard error for five treatments (N_dK_d= no N and no K, N_{hr}K_d = 60 kg ha⁻¹ N and no K, N_rK_d = 120 kg ha⁻¹ N and no K, N_{hr}K_r = 60 kg ha⁻¹ N and 60 kg ha⁻¹ K, N_rK_r = 120 kg ha⁻¹ N and 60 kg ha⁻¹ K)

by Usherwood and Segars (2001). Application of K fertilizer increased the number of grains per panicle due to increased fertilization (Bahmaniar and Ranjbar, 2007) and also due to healthy grains i.e. 1000 grains mass (data not shown) may be due to efficient translocation of sugars to grains. Application of N and K in combination produced maximum number of grains per panicle (Bahmaniar and Ranjbar, 2007) and overall performance of crop was better than N application only (Uddin *et al.*, 2013). Application of N fertilizer with deficient K conditions may increase the vegetative growth but may not produce more and healthy grains due to inefficient fertilization as well as sugar translocation to grains. Qiangsheng *et al.*, (2004) reported increase in yield with recommended K fertilizer application and positive effect of combined application of N and K on paddy yield (Bahmaniar *et al.*, 2007; Fig. 1B).

Lodging is among the major factor responsible for growth and yield reduction in rice, especially. Improvement in stem strength will definitely reduce the lodging losses. A number of factors are responsible for stem resistance against lodging which includes plant height, basal internode space, fiber content *i.e.* cellulose, lignin (Islam *et al.*, 2007). Potassium plays role in hormonal balance and may increase the number of internode so as the decrease in first internode space (Beringer, 1980). Present study has clearly shown that plant height was not decreased by application of K even was increased in basmati 515; however significant decrease in basal internode space may contribute stem strength to resist against lodging (Fig. 1A). Similar results has been shown earlier by Wakhloo, (1975). Furthermore, Rahman *et al.*, (2007) noticed that high N dose produced more vegetative growth and increased plant height which may increase length of basal internode. It was further explained that heavy N application decreases upper internode length, stem thickness and diameter, and increased lower internode length. High N dose causes reduced stem base strength, stem wall thickness and stem diameter (Hobbs *et al.*, 1998). Application of N at transplanting causes lodging while fertilization at early booting stage is ideal causing lower lodging percentage. Generally N has less effect on root: shoot ratio making the plant more vulnerable to lodging (Pinthus, 1973). Basal internode space in comparison with stem K contents has significant results that show a positive relation between them (Fig. 2A).

Concentrations of stem strengthening components increased with application of K nutrition. Plants with sufficient K synthesize high molecular weight compounds like cellulose (Marschner, 2012). Increase in cellulose concentration by K application along N fertilization was also observed by Fengzhan *et al.*, (2010) and Zhou *et al.*, (2012). Shang Lian *et al.*, (2009) also observed that cellulose concentration increased with combined application of N and K. Silicon concentration of plant parts differ from each and generally Si

accumulation decreased with increased N application which may reduce stem strength. Application of K along with N fertilizer increased Si concentration in rice shoot (Table 3) which is clear indication of stem strengthening by K application.




Figure 2. Correlation graphs between K concentration and (A) basal internode space (B) cellulose % (C) silicon %. Five treatments N_dK_d= no N and no K, N_{hr}K_d = 60 kg ha⁻¹ N and no K, N_rK_d = 120 kg ha⁻¹ N and no K, N_{hr}K_r = 60 kg ha⁻¹ N and 60 kg ha⁻¹ K, N_rK_r = 120 kg ha⁻¹ N and 60 kg ha⁻¹ K.

Stem K contents have positive correlation with both stem strengthening components silicon (R= 0.75) and cellulose (R= 0.35) and may reduce rice lodging (Fig. 2B, 2C). Silicon accumulation in shoot is beneficial for plant due to the formation of silica gel that is accumulated on leaf surface,

stems and plants other organs (Ma *et al.*, 2001). A number of researchers have already reported that application of K increase the Si concentration in plant shoot as well as K concentration (Qiangsheng *et al.*, 2004; Bahmaniar *et al.*, 2007). Potassium fertilization affect the plants K concentration, K fertilized plants contained more K than the unfertilized plants (Sulok-kevin *et al.*, 2007). Early-season plant K fertilization increases plant K uptake than the late season application (Slaton *et al.*, 2004). More K content of plant of fertilized plots was due to the application of K fertilizer than the un-fertilized plots and more K in super basmati may be due to its specific genotypic character. Increase in N concentration by application of higher dose of K may show the synergistic effect of K fertilization for N uptake by rice plants (Table 3).

Conclusions: Application of K fertilizer in combination with N fertilization increased paddy yield owing to increase in grains per panicle and improved grain quality. Increased concentration of acid detergent fiber and cellulose in shoot by application of K along with N may also contribute to stem strength. Positive correlation of shoot-K concentration with shoot-cellulose and Si and negative correlation with basal internode space indicate the contribution of K to stem strength of rice plants to decrease lodging losses.

REFERENCES

- Analytical Software. 2005. Statistix 8.1 User's Manual. Analytical Software, Tallahassee, Florida.
- Anonymous. 2014. Economic Survey of Pakistan Ministry of Food, Agriculture and Livestock, Federal Bureau of Statistics, Islamabad.
- Bahmaniar, M.A. and G. A. Ranjbar. 2007. Response of rice cultivars to rates of nitrogen and potassium application in field and pot conditions. *Pak. J. Biol. Sci.* 10:1430-1437.
- Bahmaniar, M.A., G. A. Ranjbar and S. H. Ahmadian. 2007. Effects of N and K applications on agronomic characteristics of two Iranian and landrace rice (*Oryza sativa*L.) cultivars. *Pak. J. Biol. Sci.* 10:880-886.
- Baloch, M. S., I. U. Awan, G. Hassan, M. A. Khan, K. Ahmad and M. Z. Sulemani. 2004. Quantitative assessment of social and soma input variables relating to rice production in Dera Ismail Khan, Pakistan. *Pak. J. Agron.* 3: 52-58.
- Beringer, H. 1980. Nutritional and environmental effects on yield formation. p155-173, In: *Physiological Aspects of Crop Production*, proceeding of 15th Colloquium of the International Potash Institute. Horgen, Switzerland.
- Berry, P. M., M. Sterling, J. H. Spink, C. J. Baker, R. Sylvester-Bradley, S. J. Mooney, A. R. Tams and A. R. Ennos. 2004. Understanding and reducing lodging in cereals. *Adv. Agron.* 84: 217 - 271.
- Bremner, J. M. and C. S. Mulvaney. 1982. Nitrogen total. P. 595-624. In: A.L. Page (ed) *Methods of soil analysis*. Agron. No. 9, Part 2: Chemical and microbiological properties, 2nd ed., American Society of Agronomy, Madison, WI, USA.
- Cakmak, I. 2005. The role of potassium in alleviating detrimental effects of abiotic stress in plants. *J. Plant Nutr. Soil Sci.* 168: 521-523.
- Chang, T. T. 1964. Virtual differences in lodging resistance. *Int. Rice Comm. Newsl.* 13: 1-11.
- Chapman, H. D. and P. F. Pratt. 1961. *Methods of analysis for soils, plants and waters*. Division Agriculture Sciences, University California Riverside, USA.
- Estefan, G., R. Sommer, J. Ryan. 2013. *Methods of Soil, Plant, and Water Analysis: A manual for the West Asia and North Africa region*. ICARDA, Beirut, Lebanon
- FAO. 2012. *World Food and Agriculture*. Food and Agriculture Organization, Rome, Italy
- Feng-zhuan, Z., J. Zheng-xun, M. Guo-hui, S. Wen-nan, L. Hai-ying, X. Mei-lan, and L. Yan. 2010. Relationship between lodging resistance and chemical contents in culms and sheaths of japonica rice during grain filling. *Rice Sci.* 17:311 - 318.
- Goering, H. K. and P. J. V. Soest. 1970. *Forage Fiber Analysis (Apparatus, Reagents, Procedures and Some Application)*. Agricultural Handbook No. 379, Agricultural Research Service, USDA.
- Haghighi, H., M. S. Daliri, H. R. Mobaser, and A. A. Moosavi. 2011. Effect of different nitrogen and potassium fertilizer levels on quality and quantity yield of flue-cured tobacco. *World Appl. Sci. J.* 15: 941-946.
- Hobbs, P. R., K. D. Sayre, and I. Ortiz-Monasterio. 1998. Increasing wheat yields through agronomic means. Mexico, D.F. Mexico.
- Islam, M. S. S. Peng, R. M. Visperas, N. Ereful, M. S. U. Bhuiya and A. W. Julfiquar. 2007. Lodging-related morphological traits of hybrid rice in a tropical irrigated ecosystem. *Field Crops Res.* 101:240 – 248.
- Kong, L., M. Sun, F. Wang, J. Liu, B. Feng, J. Si, B. Zhang, S. Li, and H. Li. 2014. Effects of high NH₄⁺ on K⁺ uptake, culm mechanical strength and grain filling in wheat. *Front. Plant Sci.* doi: 10.3389/fpls.2014.00703.
- Kono, M. 1995. Physiological aspects of lodging. *Sci. Rice Plant* 2: 971-991.
- Ma, J. F., Y. Miyake, and F. Takahashi. 2001. Silicon as a beneficial element for crop plants. In: *Datonoff L, Korndorfer G, Snyder G, eds. Silicon in Agriculture*. New York: Elsevier Science Publishing. 17– 39.
- Marschner, H. 1995. *Mineral Nutrition of Higher Plants*. 2nd ed. p. 889 Academic Press, San Diego, USA.
- Marschner, P. 2012. *Marschner's Mineral Nutrition of Higher Plants*. 3rd ed. p-178 Academic Press; London, UK.

- McGill, D. and C. T. Figueiredo. 1993. Total Nitrogen, p. 201 - 211. In: M.R. Carter (ed.), Soil sampling and methods of analysis. Lewis Publication, Boca Raton, FL, USA.
- NFDC. 2014. Annual Review of Fertilizers, National Fertilizer Development Center, Islamabad Pakistan.
- Pinthus, M. J. 1973. Lodging in wheat, barley and oats: The phenomenon, its causes and preventive measures. *Adv. Agron.* 25: 209-263.
- Qiangsheng, W., Z. Ruohong, D. Yanfeng, C. Weixing, and H. Pisheng. 2004. Effects of potassium fertilizer application rates on plant potassium accumulation and grain quality of japonica rice. *Sci. Agric. Sinica.* 37: 1444-1450.
- Rahman, M. H., M. H. Ali, M. M. Ali and M. M. Khatun. 2007. Effect of different level of nitrogen on growth and yield of transplantaman rice cv. BRRI dhan32. *Int. J. Sus. Crop Prod.* 2: 28-34.
- Rajkumara, S. 2008. Lodging in cereals – a review. *Agric. Rev.* 29: 55 - 60.
- Shakouri, M. J., A. V. Vajargah, M. G. Gavabar, S. Mafakheri, M. Zargar. 2012. Rice vegetative response to different biological and chemical fertilizer. *Adv. Environ. Biol. J.* 6:859-863.
- Shang Lian, H., C. Ying, L. Xue Qin, H. Ying, J. Ju Qing, S. Xia, Z. Meng. 2009. Effects of nitrogen and potassium on dynamic accumulation of lignin and cellulose in *Neosinocalamus affinis*. *Bull. Bot. Res.* 29: 728 - 733.
- Slaton, N. A., D. Dunn, B. Pugh. 2004. Potassium nutrition of flood irrigated rice. *Better Crops.* 88: 20 - 23.
- Sulok-Kevin M. T., O. H. Ahmad, W. Asrin, A. Rajan and M. Ahzam. 2007. Towards growing Bario rice on low land soils: A preliminary nitrogen and potassium fertilization trial. *Am. J. Agric. Biol. Sci.* 2: 99 - 105.
- Tripathi, S. C., K. D. Sayre, J. N. Kaul, and R. S. Narang. 2003. Growth and morphology of spring wheat (*Triticum aestivum* L.) culms and their association with lodging: effects of genotypes, N levels and ethephon. *Field Crops Res.* 84: 271-290.
- Uddin, S., M. A. R. Sarkar and M. M. Rahman. 2013. Effect of nitrogen and potassium on yield of dry direct seeded rice cv. NERICA 1 in ausseason. *Int. J. Agron. Plant Prod.* 4: 69-75.
- Usherwood, N. R. and W. I. Segars. 2001. Nitrogen interactions with phosphorus and potassium for optimum crop yield, nitrogen use effectiveness, and environmental stewardship. *Sci. World J.* 2: 57- 60.
- Wakeel, A. 2013. Potassium-sodium interactions in soil and plant under saline-sodic conditions. *J. Plant Nutr. Soil Sci.* 176: 344-354.
- Wakeel, A., M. Farooq, M. Qadir and S. Schubert. 2011. Potassium substitution by sodium in plants. *Crit. Rev. Plant Sci.* 30: 401- 413.
- Wakeel, A., S. Hanstein, B. Pitann, S. Schubert. 2010. Hydrolytic and pumping activity of H⁺-ATPase from leaves of sugar beet (*Beta vulgaris* L.) as affected by salt stress. *J. Plant Physiol.* 167: 725-731.
- Wakhloo, J. L. I. 1975. Interaction between foliar potassium and applied giberellic acid and 6-furfurylaminopurine. *J. Exp. Bot.* 26: 433-440.
- Zhou, D., J. Chen, J. She, J. Tong and Y. Chen. 2012. Temporal dynamics of shearing force of rice stem. *Biomass Bioenergy* 47: 109-114.
- Zörb, C., M. Senbayram, E. Peiter. 2014. Potassium in agriculture – Status and perspectives. *J. Plant Physiol.* 171: 656-669.