



Scientific Inquiry and Review (SIR)

Volume 4, Issue 1, January to March 2020

ISSN (P): 2521-2427, ISSN (E): 2521-2435

Journal DOI: <https://doi.org/10.32350/sir>

Issue DOI: <https://doi.org/10.32350/sir.41>

Homepage: <https://journals.umt.edu.pk/index.php/SIR/Home>

Journal QR Code:



Article

Design, Synthesis and Characterization of 2, 4-Dimethylphenyl Hydrazine based Chemosensors

Author(s) Saema Salim
Qamar Ali

Online Published January to March 2020

Article DOI <https://doi.org/10.32350/sir.41.04>

QR Code
of Article



Saema Salim

To cite this
Article

Salim S, Ali Q. Design, synthesis and characterization of 2, 4-dimethylphenyl hydrazine based chemosensors. *Sci Inquiry Rev.* 2020;4(1): 46–58.

[Crossref](#)

Copyright
Information

This article is open access and is distributed under the terms of Creative Commons Attribution – Share Alike 4.0 International License.



A publication of the
School of Science, University of Management and Technology
Lahore, Pakistan.

Indexing Agency



Design, Synthesis and Characterization of 2,4-Dimethylphenyl Hydrazine Based Chemosensors

Saema Salim*, Qamar Ali

Department of Chemistry, Division of Science & Technology,
University of Education, Lahore, Pakistan

*saema786salim@gmail.com

Abstract

Hydrazones are a unique division of compounds found in various syntheses. They have an important role in synthetic chemistry due to their different biological properties, such as antifungal, anticonvulsant, antibacterial, antimalarial, anti-inflammatory and anti-TB properties. This paper reports the syntheses of a series of 3 hydrazones based on the condensation of 2,4-dimethylphenyl hydrazine HCl with different aromatic carbonyl compounds. The structures of the synthesized compounds were confirmed by EIMS. These compounds can act as ideal candidates in chemosensor chemistry.

Keywords: chemosensors, hydrazone, sensors, 2,4-dimethylphenyl hydrazine HCL

Introduction

Sensors quantify a physical quantity converting it into a signal which can be examined through electronic equipment. For example, a glass thermometer filled with mercury measures the temperature range based on the contraction and expansion of liquid mercury, which can be calibrated on the glass tube. A thermocouple is an instrument that converts the measuring temperature into an outgoing current through a voltmeter. For the sake of precision, well-known standards are used to calibrate sensors. They are based on the principles of reflectance, absorbance, light scattering, fluorescence, luminescence, refractive index, and optothermal effect [1].

Depending on their principles, the sensors are classified as magnetic sensors, electrical sensors, electrochemical sensors, thermometric sensors, mass sensitive sensors and chemosensors. Chemical sensors are small devices that display actual time and give online information about the existence of particular ions and/or compounds in complex samples. Depending on their type, they are classified as direct and distance chemosensors. Olfaction receiving

neurons exemplify distance chemosensors which can detect chemicals in the gas state [2]. The design and synthesis of chemosensors for heavy and/or transition metal (HTM) ions with a high accuracy continues to be an active part of supramolecular chemistry. They are designed to incorporate three components: firstly, a chemical receptor that receives the guest of interest frequently with considerable selectivity; secondly, a transducer or signaling unit that converts the binding component into a measurable physical change; and lastly, a design of measuring that change and converting it into useful information [3]. The following is a schematic diagram of this type of chemosensors.



Figure 1. Quantum dot sensor for the specific detection of the explosive 2, 4, 6-trinitrotoluene (TNT) in aqueous environments [4]

The identification and sensing of ionic species such as cations and anions have been an area of interest for many scientific research groups. Transition metal complexes of Schiff bases are the most extensively studied coordination compounds because they have a number of uses. The property that displays electronic relation and structural relation with Schiff bases also plays a vital role in catalysis [5].

Hydrazine (N_2H_4) is a very rigorous base and a reducing agent that plays an important role in pharmaceutical, chemical and agricultural industries [6, 7]. It is used in projectile propulsion methods such as maximum energy fuel and it is the primary material used in a variety of products including foaming agents for antioxidants, plastics and polymer. Still, it has been categorized as a possible human carcinogen with a threshold limit value (TLV) of 10 ppb by the U.S. Environmental Protection Agency (EPA). It also causes mutations and has a neurotoxic impact on the kidneys, liver and lungs [8]. Thus, numerous recognition methods for hydrazine have been devised lately. Analyzing techniques such as MS-Gas chromatography, MS-Liquid chromatography and

capillary electrophoresis are used for the detection of hydrazine [9, 10, 11].

Hydrazones are a form of organic compounds with the structural formula $R^1R^2C=NNH_2$ [12] and they are identified in many useful synthetic products. Due to their extensive biological properties including anti-inflammatory, antifungal, antimalarial, antituberculosis, antibacterial, and anticonvulsant properties, hydrazine-hydrazone have gain considerable importance [13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25]. This paper reports the synthesis, design and characterization of a series of 3 hydrazones based on the condensation of 2,4-dimethylphenyl hydrazine HCl with different aromatic carbonyl compounds. The structures of the synthesized compounds were confirmed by EIMS. These compounds can act as ideal candidates in chemosensor chemistry.




Figure 2. Structure of hydrazone functional group

Taking into consideration the significant roles that transition metal ions played in many biomedical and environmental researches, the growth of hydrazone derivative ligands as artificial receptors was used for selective sensing in the current research. The development of aromatic hydrazone derived is known for measuring the concentration of low molecular weight ketones and aldehydes, for instance in gas flowing. The process of elution and analyzing is further proceeded via high performance liquid chromatography using a UV detector. Hydrazone based coupling techniques are utilized in medical biotechnologies to couple drugs with the targeted antibodies (see ADC), such as the antibodies against a specific form of cancer cell [26].

2. Material and Method

Chemosensors are based on molecular recognition that is emerging as a very important research field in the area of supramolecular chemistry,

which permits the determination of guest by binding-induced modification in spectroscopic or electrochemical properties [27]. A typical chemosensor comprises two parts: a receptor site that selectively attaches an analyte and a signaling unit connected through a spacer. Receptor components vary from simple metal-chelating compounds to a synthetic host compound or to proteins or protein fragments in case of biosensors. By coupling these molecules to a sensitive fluorescent readout system, molecular sensors are devised for analytes ranging from metal ions to simple organic compounds [28].





Figure 3. General scheme for the synthesis of chemosensors

2.1. Syntheses of Hydrazone Derivatives Acting as Chemosensors


All compounds were synthesized by mixing hydrazine hydrochloride acid and various acetophenones or aldehydes in 1:1 molar ratios which is a straightforward approach to synthesize the series of hydrazones derivatives.

2.1.1. Synthesis of 2, 4, 5-trimethoxybenzaldehyde-2, 4 dimethylphenyl hydrazine. 113.63 mg (0.579 mmole) of 2, 4, 5-trimethoxybenzaldehyde was mixed in 20 ml of ethanol in a round bottomed flask. After heating the mixture for 10 minutes at 78°C, 5 droplets of acetic acid were added to it and the reaction mixture was heated to another 5 minutes. Then, 100 mg (0.579 mmole) of 2,4-dimethylphenyl hydrazine HCl was added and the mixture was refluxed for one hour. After refluxing, the reaction mixture was cooled down and 10 ml of sodium bicarbonate (1M) was added to it. Then, 10 ml of dichloromethane was added to separate the layers. The evaporation of the solvent resulted in fine crystals of compound 1 giving 90% yield.




Scheme 1: Synthesis of 2, 4, 5-trimethoxybenzaldehyde-2,4-dimethylphenyl hydrazone (I)

2.2.2. Synthesis of 3, 4, 5-trimethoxybenzaldehyde-2,4-dimethylphenyl hydrazone. 113.63 mg (0.579 mmole) of a 3, 4, 5-trimethoxybenzaldehyde was mixed in 20 ml of ethanol in a round bottomed flask. After heating the mixture for 10 minutes at 78°C , 5 droplets of acetic acid were added to it and the reaction mixture was heated to another 5 minutes. Then, 100 mg (0.579 mmole) of 2, 4-dimethylphenyl hydrazine HCl was added and the mixture was refluxed for one hour. After refluxing, the reaction mixture was cooled down and 10 ml of sodium bicarbonate (1M) was added to it. Then, 10 ml of dichloromethane was added to separate the layers. The evaporation of the solvent resulted in fine crystals of compound 2 giving 88% yield.



Scheme 2: Synthesis of 3, 4, 5-Trimethoxybenzaldehyde-2,4-Dimethylphenyl Hydrazone (2)

2.2.3. Synthesis of acetophenone-2,4-dimethylphenyl hydrazine. 0.1 ml of acetophenone was mixed in 20 ml of ethanol in a round bottomed flask. After heating the mixture for 10 minutes at 78°C, 5 droplets of acetic acid were added to it and the reaction mixture was heated to another 5 minutes. Then, 100 mg (0.579 mmole) of 2,4-dimethylphenyl hydrazine HCl was added and the mixture was refluxed for one hour. After refluxing, the reaction mixture was cooled down and 10 ml of sodium bicarbonate (1M) was added to it. Then, 10 ml of dichloromethane was added to separate the layers. The evaporation of the solvent resulted in fine crystals of compound 3 giving 92% yield.



Scheme 3: Synthesis of acetophenone-2, 4-dimethylphenyl hydrazone (3)

3. Results and Discussion

Different types of hydrazones were synthesized by reacting a variety of aromatic aldehydes and ketones with 2,4-dimethylphenylhydrazine hydrochloride. The synthesized compounds were confirmed by mass spectrometry.

3.1. Compound 1

3.1.1. 2, 4, 5-trimethoxybenzaldehyde-2,4-dimethylphenyl hydrazine. Molecular mass of Compound 1 having the molecular formula $C_{18}H_{22}N_2O_3$ was 314, which was found to be comparable with the calculated molecular mass. Molecular mass was determined using JEOL JMS 600-H. Compound 1 showed different fragment peaks such as 212, 157 and 312.




Figure 4. EIMS of compound (1)

3.2. Compound 2

3.2.1. 3, 4, 5-trimethoxybenzaldehyde-2,4-dimethylphenyl hydrazine. Molecular mass of Compound 2 having the molecular formula $C_{18}H_{22}N_2O_3$ was 314, which was found to be comparable with the calculated molecular mass. Molecular mass was determined using JEOL JMS 600-H. Compound 2 showed different fragment peaks such as 212, 157 and 312.




Figure 5. EIMS of compound (2)

3.3. Compound 3

3.3.1. Acetophenone-2,4-dimethylphenyl hydrazine. Molecular mass of Compound 3 having the molecular formula $C_{16}H_{18}N_2$ was 238, which was found to be comparable with the calculated molecular mass. Molecular mass was determined using JEOL JMS 600-H. Compound 3 showed different fragment peaks such as 212, 147 and 128.




Figure 6. EIMS of compound (3)

4. Conclusion

Chemosensors are sensory receptors that transform a chemical signal into an active potential. Generally, they determine the chemical stimuli present in an environment. Chemosensors have a variety of applications in biological, medicinal and environmental sciences. Hydrazones can act as synthetic chemosensors. Hydrazones are mostly produced through the reaction of hydrazines with ketones or aldehydes. Hydrazones are a significant class of compounds present in many synthetic products. They also show variable biological properties including anticonvulsant, antibacterial, antifungal, antimalarial, anti-inflammatory, and anti-tuberculosis properties. In order to synthesize different hydrazones, a variety of aromatic aldehydes and ketones were refluxed with 2,4-dimethylphenyl hydrazine in the presence of acetic

acid acting as a catalyst and ethanol acting as a solvent. The reaction was quenched by adding 1M aqueous solution of NaHCO_3 . The required compound was extracted by adding 10 ml dichloromethane. The evaporation of the solvent resulted in fine crystals of target molecules which were characterized by EIMS. The synthesized compounds can act as ideal candidates in chemosensor chemistry. Target molecules were synthesized by the condensation of 2,4-dimethylphenyl hydrazine with the following aromatic aldehydes and ketones (3, 4, 5-trimethoxybenzaldehyde, 2, 4, 5-trimethoxybenzaldehyde, benzophenone). The synthesized compounds were obtained as fine crystals with good yield and were characterized by EIMS. These compounds are expected to act as chemosensors.

5. Acknowledgement

Thanks to Allah Almighty, Who pulled us through the time when every stone was turned against us. All respects for the Holy Prophet Muhammad (SAW). My thanks to my parents and lastly, all respect to my research supervisor and the university which fully funded our research and provided the required research infrastructure.

References

- [1] Malele CN. *Design, synthesis and photophysics of dipyrrolequinoxaline receptor loaded fluorescent conjugated polymer chemosensors.* 2019. <https://www.antoineonline.com/Product.aspx?productCode=0009781244072510>
- [2] Shi P, Zhang J. Extraordinary diversity of chemosensory receptor gene repertoires among vertebrates. *Results Probl Cell Differ.* 2009;47: 57–58.
- [3] Jing W, Chang H-S, Xu ZC, et al. A colorimetric and fluorescent turn-on chemosensor for Zn^{+2} based on an azobenzene-containing compound. *J Tetrahedron.* 2009;65: 2307–6964.
- [4] Ellen R, Goldman IL, Medintz JL, et al. A hybrid quantum dot–antibody fragment fluorescence resonance energy transfer-based TNT sensor. *J Am Chem Soc.* 2005;127(18): 6744–51.
- [5] Marwan S, Abdussalam S, Tan T, Saleh MI, Bahrudin S. Spectroscopic studies of 1,4-bis(4-dimethylaminobenzyl)-2,3-

- diaza-1,3-butadieneas Colorimetric Reagent for Cu^{+2} . *World Appl Sci J*. 2011;15(4): 598–605.
- [6] Ragnarsson U. Synthetic methodology for alkyl substituted hydrazines. *Chem Soc Rev*. 2011;30: 205–213.
- [7] Narayanan SS, Scholz F. A comparative study of the electrocatalytic activities of some metal hexacyanoferrates for the oxidation of hydrazine. *Electroanalysis*. 1999; 11(7): 465–469.
- [8] Khomsan T, Rattha N, Sinchai C, et al. A chromogenic and fluorogenic rhodol-based chemosensor for hydrazine detection and its application in live cell bioimaging. *Spectrochim Acta A Mol Biomol Spectrosc*. 2018; 195: 136–141. <https://www.researchgate.net/publication/322459741>
- [9] Bhutani H, Singh S, Vir S, et al. LC and LC-MS study of stress decomposition behaviour of isoniazid and establishment of validated stability-indicating assay method. *J Pharm Biomed Anal*. 2007; 143: 1213–1220.
- [10] Liu J, Zhou W, You T, Li F, Wang E, Dong S. Detection of hydrazine, methylhydrazine, and isoniazid by capillary electrophoresis with a palladium-modified microdisk array electrode. *Anal Chem*. 1996;68: 3350–3353.
- [11] Sun T, Wetzel SJ, Johnson ME, Surlow BA, Patton J-V. Development and validation of a hydrophilic interaction liquid chromatography-tandem mass spectrometry method for the quantification of lipid-related extracellular metabolites in *Saccharomyces cerevisiae*. *J Chromatogr B*. 2012;897: 1–9. doi:[10.1016/j.jchromb.2012.03.034](https://doi.org/10.1016/j.jchromb.2012.03.034)
- [12] March J. *Advanced organic chemistry: reactions, mechanisms, and structure* (3rd ed.). New York: Wiley; 1985.
- [13] Rahman VM, Mukhtar S, Ansari WH, Lemiere G. Synthesis, stereochemistry and biological activity of some novel long alkyl chain substituted thiazolidin-4-ones and thiazan-4-one from 10-undecenoic acid hydrazide. *Eur J Med Chem*. 2005;40: 173–184.
- [14] Dimmock JR, Vashishtha SC, Stables JP. Anticonvulsant properties of various acetylhydrazones, oxamoylhydrazones and

- semicarbazones derived from aromatic and unsaturated carbonyl compounds. *Eur J Med Chem.* 2000; 35: 241–248.
- [15] Yapia R, La Mara MP, Massieu GH. Modifications of brain glutamate decarboxylase activity by pyridoxal phosphate- β -glutamyl hydrazone. *Biochem. Pharmacol.* 1967;16: 1211–1218.
- [16] Sava G, Perissin L, Lassiani L, Zabucchi G. Antiinflammatory action of hydrosoluble dimethyl-triazenes on the carrageen induced edema in guinea pigs. *Chem Biol Interact.* 1985;53: 37–43.
- [17] Xia YL, Chuan-Dong F, Zhao BX, Zhao J, Shin DS, Miaom JY. Synthesis and structure-activity relationships of novel 1-arylmethyl-3-aryl-1H-pyrazole-5-carbohydrazide hydrazone derivatives as potential agents A549 lung cancer cells. *Eur J Med Chem.* 2008;43: 2347–2353.
- [18] Melnyk P, Leroux V, Serghergert C, Grellier P. Design, synthesis and in vitro antimalarial activity of an acylhydrazone library. *Bioorg Med Chem Lett.* 2006;16: 31–35.
- [19] Ajani OO, Obafemi CA, Nwinyi OC, Akinpelu DA. Microwave assisted synthesis and antimicrobial activity of 2- quinoxalinone-3-hydrazone derivatives. *Bioorg Med Chem.* 2010;18: 214–221.
- [20] Zheng LW, Wu LL, Zhao BX, Dong WL, Miao YJ. Synthesis of novel substituted pyrazole-5-carbohydrazide hydrazone derivatives and discovery of a potent apoptosis inducer in A549 lung cancer cells. *Bioorg Med Chem.* 2009;17: 1957–1962.
- [21] Bhagavan NV. *Medical biochemistry.* 4th ed. Amsterdam, The Netherlands: Elsevier; 2002.
- [22] Saulnier MG, Velaparthi U, Zimmermann K. Progress in heterocyclic synthesis. In: Gordon W. Gribble, J. Joule, ed. *Five-membered ring system: with N and S (SE) atoms.* Vol. 16. Amsterdam, The Netherlands: Elsevier; 2005.
- [23] Short EI. Studies on the inactivation of isonicotinyl acid hydrazide in normal subjects and tuberculous patients. *Tubercle.* 1962;43: 33–42.
- [24] Holdiness MR. A review of blood dyscrasias induced by the antituberculosis drugs. *Tubercle.* 1987;68: 301–309.

- [25] Faroumadi A, Kiano Z, Soltani F. Antituberculosis agents VIII: synthesis and in vitro antimycobacterial activity of alkyl a-[5-(5-nitro-2-thienyl)-1,3,4- thiadiazole-2-ylthio] acetates. *Farmaco*. 2003;58: 1073–1076.
- [26] Wu AM, Senter PD. Arming antibodies: prospects and challenges for immunoconjugates. *Nat Biotechnol*. 2005;23(9): 1137–46.
- [27] Yuee Y, Yan H, Guopo Z, Xingming K. A novel azathia-crown ether dye chromogenic chemosensor for the selective detection of mercury (II) ion. *Dyes Pigm*, 2008;79(2): 210–215.
- [28] Chen C. Fluorescent, sequence-selective peptide detection by synthetic small molecules. *Science*. 1998;279(5352): 851–853.