

## Research Article



## Correlation of Obesity with Hypertension in Medical Students

Rana Khurram Aftab<sup>1\*</sup>, Farhat Ijaz<sup>2</sup>, Samia Jawed<sup>3</sup>

<sup>1</sup>Assistant Professor of Physiology, King Edward Medical University, Lahore; <sup>2</sup>Assistant Professor of Physiology, CMH Lahore Medical College, Lahore; <sup>3</sup>Professor of Physiology, King Edward Medical University, Lahore, Pakistan.

**Abstract** | The relationship between obesity and hypertension has long been recognized. Overweight and obesity are disorders of balance of energy affecting people of various ethnic groups, age and socio-economic statuses. It has been found that obesity is closely associated and correlated with hypertension. Preventive strategies for obesity may offer a cost-effective approach towards lowering blood pressure.

**Objectives:** To determine the frequency of obesity according to body mass index and waist hip ratio and its relationship with hypertension among medical students.

**Methodology:** It was a descriptive cross sectional study where a total of 213 students of 1<sup>st</sup> year MBBS and BDS were involved. Their waist circumference, hip circumference, height and weight were recorded. Moreover, on these subjects body mass index (BMI), waist circumference (WC), waist hip ratio (WHR) were calculated and blood pressure was recorded.

**Results:** Amongst 213 students, a total of 97(45.5%) were male and 116 (54.5%) were females. Among studied subjects, 2 (2.1%) males and 8 (6.9%) females were underweight, whereas 55(56.7%) male and 52(44.8%) were of normal weight. The prevalence of overweight in male and females was 17(17.5%) and 22(19.0%), prevalence in obese I in male and female was 20(20.6%) and 23(19.8%), and obese II was 3(3.1%) and 11(9.5%) respectively. Frequency of obesity was found to be more in females than in males according to BMI and WHR. Hypertension was present in 12(30.8%) normal weight students, 6(15.4%) overweight, 14(35.9%) obese I and 7(17.9%) obese II subjects. Correlation between BMI, WC, and WHR was positively significant with SBP and DBP.

**Conclusion:** High prevalence of overweight and obesity was observed in students. Strong positive correlation of BMI, WC and WHR with systolic and diastolic blood pressures was shown in our study. Results of this study indicate that increase in BMI leads to increase in hypertension.

Received | August 08, 2017; Accepted | December 10, 2017; Published | March 18, 2018

\*Correspondence | Dr. Rana Khurram, King Edward Medical University, Lahore; Email: drnanakurram81@outlook.com

Citation | Khurram, R. 2018. Correlation of obesity with hypertension in medical students. *Annals of King Edward Medical University*, 24(1): 637-642

DOI | <http://dx.doi.org/10.21649/journal.akemu/2018/24.1.637-642>

Keywords | Body mass index, Diabetes mellitus, Hypertension, Obesity, Waist hip ratio, Waist circumference

### Introduction

Overweight and obesity is a medical condition in which abnormal or excessive fat accumulates in the body<sup>(1)</sup>. It is the most accustomed form of malnutrition in not only developed but developing countries as well. More than 1.5 billion people in the world are overweight or obese<sup>(2)</sup>. Obesity is considered to be

known public health problem internationally with rapidly rising prevalence in several developed countries because nearly two third adult population are overweight<sup>(3)</sup>. Prevalence of obesity in urban Pakistani population is 22-37%<sup>(4)</sup>. Increased prevalence of diabetes mellitus (DM), hypertension, and cardiovascular diseases (CVS) has been reported among overweight and obese persons as a result of unhealthy

lifestyle<sup>(5)</sup>. Obesity is found to be major risk factor for the development of hypertension<sup>(6)</sup>. It is more likely that obese patients will develop arterial hypertension and may require antihypertensive medication<sup>(7)</sup>. Early detection of obesity and its intervention with treatment could prevent the long-term consequences like the development of cardiovascular diseases.

Estimated prevalence for HTN in the world is almost one billion and it is increasing in developing countries. Hypertension has almost two times higher prevalence in the obese than in non-obese subjects<sup>(8)</sup>. Abdominal or central obesity is an emergent clinical and public health problem. It is closely associated with increased risk of coronary heart diseases (CHD)<sup>(9)</sup>. Anthropometric measurements of central obesity such as waist circumference (WC), waist hip ratio (WHR) are found to be more strongly associated with cardiovascular risk factors as compared to BMI<sup>(10)</sup>. Obesity associated hypertension is linked with many serious conditions e.g. congestive cardiac failure, atherosclerosis, cerebrovascular and coronary artery disease, atrial fibrillation and renal insufficiency. Along with medical treatment, dietary and life style modifications are necessary for the management of obesity related HTN. Weight loss plays an important role to reverse several mechanisms that are responsible for hypertension in obese<sup>(11)</sup>.

In last few decades, there is growing trend of taking junk food in the form of fast food, lack of taking fresh fruits, fibre containing foods and vegetables. Junk food like snacks, soft drinks are preferred. Younger generation spend more time on television, mobile phones and computers thereby increasing screen time. Lack of physical activity and sedentary behaviour are responsible for rising prevalence of non-communicable diseases like diabetes mellitus (DM), hypertension (HTN), ischemic heart disease (IHD) and obesity. These conditions are leading cause of mortality and disability all parts of the world<sup>(11)</sup>.

Overweight and obesity in college students is almost 40% that is very alarming. Most of them are unaware of the increased health risks linked with being overweight<sup>(12)</sup>. A small percentage of about 8.5% of college students eat the proposed amount of fruits and vegetables. It was noticed that many college students are lacking variety in their diets. Obesity is the important challenge for Pakistani population because we are genetically prone to it<sup>(13)</sup>.

In present study, we found out the prevalence of obesity by three indexes (BMI, WHR and WC) and their association with hypertension prevalence in medical students, age group of 18-21. This young age group is important mentally, emotionally and physically as this age group is the transition period when the individuals are entering adulthood, so this age group is very important. Students joining Medical Colleges represent this group. Due to less physical activity as they have to give maximum time to their studies, they are easily predisposed to obesity and its related complications. Thus it is very important to assess their weight, this is specifically crucial in case of medical students who are future health guard of population because they have to undergo hard medical training to excel in their profession and they get a very little time for physical activities and outdoor games.

## Methodology

Our study was a descriptive cross sectional study conducted in Physiology department, CMH Lahore Medical College from December 2016 to March 2017. Total 213 students of 1<sup>st</sup> year MBBS and 1<sup>st</sup> year BDS were included in the study after approval from institutional review board & ethical committee. Informed consent was taken. Demographic profile (name, age, address, class roll number, marital status, CNIC # & Cell number) was noted on a especially designed questionnaire.

For general obesity, Student's height in centimetres and weight in kilograms was taken by stadiometer. They were advised to remove shoes for their height measurement and extra clothing before taking their weight. We calculated BMI by weight in kilograms divided by square of height in meters. On the basis of their BMI, they were divided into five groups: under weight, normal weight, over weight, obese 1 and obese 2<sup>(14)</sup>.

For abdominal obesity, (WC and hip (HC) were taken. Waist circumference was taken at the midpoint between iliac crest and lower border of rib cage and average of 3 readings was taken. Hip circumference was measured at the level of greater trochanter and 2cm above and below the trochanter was calculated. Average of 3 readings was taken. We calculated waist hip ratio (WHR) as waist circumference divided by hip circumference. The recommendations of international diabetic federation (IDF) for waist circumference in

South Asians are: Men >90cm; Women >80cm. The cut-off value for WHR for male and female in South Asians  $\geq 0.90$ cm (M);  $\geq 0.85$ cm (W)<sup>(15)</sup>.

Blood Pressure measurements were taken using stethoscope and mercury sphygmomanometer. Students were asked to sit and relax for 5 minutes and blood pressure was taken in sitting position first in right arm then in left arm. We heard First and fifth Korotkoff sounds for systolic and diastolic readings respectively. We took two readings separated by two minutes and it was averaged. Subjects were divided into four groups on the basis of blood pressure, normal, pre-hypertension, stage 1 and stage 11 HTN<sup>(16)</sup>.

Analysis was done by SPSS 21. Quantitative variables like height, weight, BMI, WC, WHR, systolic and diastolic blood pressures were presented as mean $\pm$ S.D. While qualitative variables like gender, hypertension and obesity were presented as frequency and percentages. P- Value  $\leq 0.05$  was taken as significant. For correlation between Obesity parameters & blood pressure, Pearson correlation test was used.

## Results

Among 213 students, 97(45.5%) were male and 116(54.5%) were females. A total of 2(2.1%) males and 8(6.9%) females were underweight (Table 1). So there were 10(4.7%) students who were underweight.

55(56.7%) males and 52(44.8%) females were of normal weight. Prevalence of overweight in male and females were 17(17.5%), 22(19.0%), respectively. In obese category I, males and females were 20(20.6%), 23(19.8%), while in obese category II, there were 3(3.1%) males, 11(9.5%) females. Prevalence of underweight, overweight and obesity was more in females than males according to BMI and WHR (Table 2). Hypertension was present in 12(30.8%) normal weight students, 6(15.4%) overweight, 14(35.9%) obese I and 7(17.9%) obese II subjects (Table 3). Correlation between BMI, WC and WHR was positively significant with SBP and DBP. Mean $\pm$ SD of height among medical students was 165.5 $\pm$ 9.357cm, weight was 64 $\pm$ 12.99kg, WC was 118 $\pm$ 84cm, HC was 129 $\pm$ 97cm, BMI 35 $\pm$ 23.31, WHR 0.96 $\pm$ 0.8696, SBP 138.54 $\pm$ 13.90, DBP 78.83 $\pm$ 10.30.

For comparing mean of study variables between male and female medical students independent sample t-test was used. The mean  $\pm$  standard deviation of body mass Index, Waist Hip Ratio and Waist Circumference in men and women were, respectively, 22.91 $\pm$ 3.3 and 23.6 $\pm$ 4.37Kg/m<sup>2</sup> (P=0.043), 89 $\pm$ .02 and 0.84 $\pm$ 0.04 (P=0.0000), 86.24 $\pm$ 7.06 and 83.62 $\pm$ 9.186 cm (P=0.057). The mean of SBP and DBP in men and women was 125.90 $\pm$ 13.540 and 113.88 $\pm$ 12.906 (P= .648), 83.45 $\pm$ 10.467 and 74.96 $\pm$ 8.431 (P=0.000).

**Table 1:** Comparison of study variables between male and female students

|                                      | Gender | N   | Mean    | Std. Deviation | P-value |
|--------------------------------------|--------|-----|---------|----------------|---------|
| Height (cm)                          | Male   | 97  | 173.747 | 5.7803         | .357    |
|                                      | Female | 116 | 158.720 | 5.4591         |         |
| Weight (Kg)                          | Male   | 97  | 69.412  | 12.5098        | .470    |
|                                      | Female | 116 | 59.603  | 11.6722        |         |
| Waist circumference (cm)             | Male   | 97  | 86.24   | 7.062          | .057    |
|                                      | Female | 116 | 83.62   | 9.186          |         |
| Hip circumference (cm)               | Male   | 97  | 96.43   | 7.241          | .152    |
|                                      | Female | 116 | 98.45   | 8.963          |         |
| Body Mass Index (Kg/m <sup>2</sup> ) | Male   | 97  | 22.9199 | 3.39985        | .043*   |
|                                      | Female | 116 | 23.6431 | 4.37380        |         |
| Waist to Hip Ratio                   | Male   | 97  | .8942   | .02664         | .000*   |
|                                      | Female | 116 | .8489   | .04277         |         |
| Systolic Blood Pressure (mmHg)       | Male   | 97  | 125.90  | 13.540         | .648    |
|                                      | Female | 116 | 113.88  | 12.906         |         |
| Diastolic Blood Pressure (mmHg)      | Male   | 97  | 83.45   | 10.467         | .000*   |
|                                      | Female | 116 | 74.96   | 8.431          |         |

**Table 2:** Prevalence of general and central obesity in male and female students

|                              |               | Male    | Female    | Total   |      |
|------------------------------|---------------|---------|-----------|---------|------|
| BMI group                    | Underweight   | 2       | 8         | 10      | .113 |
|                              |               | 2.1%    | 6.9%      | 4.7%    |      |
|                              | Normal weight | 55      | 52        | 107     |      |
|                              |               | 56.7%   | 44.8%     | 50.2%   |      |
|                              | Overweight    | 17      | 22        | 39      |      |
|                              |               | 17.5%   | 19.0%     | 18.3%   |      |
|                              | Obese I       | 20      | 23        | 43      |      |
|                              |               | 20.6%   | 19.8%     | 20.2%   |      |
|                              | Obese II      | 3       | 11        | 14      |      |
|                              |               | 3.1%    | 9.5%      | 6.6%    |      |
| Grouping on the basis of WHR | Non Obese     | 66      | 64        | 130     | .05  |
|                              |               | 68.0%   | 55.2%     | 61.0%   |      |
|                              | Obese         | 31(32%) | 52(44.8%) | 83(39%) |      |

There were 15.4% overweight students who were hypertensive, in obese I 35.9% and obese II 17.9% were hypertensive. Association of obesity on the basis of BMI groups with blood pressure was determined by chi-square test ( $p$  value = 0.000)

**Table 3:** Prevalence of hypertension on the basis of obesity by BMI

|           |               | Hypertensive/<br>non hypertensive |        | Total  | P- value |
|-----------|---------------|-----------------------------------|--------|--------|----------|
|           |               | Present                           | Absent |        |          |
| BMI group | Underweight   | 0                                 | 10     | 10     | .000     |
|           |               | .0%                               | 5.7%   | 4.7%   |          |
|           | Normal weight | 12                                | 95     | 107    |          |
|           |               | 30.8%                             | 54.6%  | 50.2%  |          |
|           | Overweight    | 6                                 | 33     | 39     |          |
|           |               | 15.4%                             | 19.0%  | 18.3%  |          |
|           | Obese I       | 14                                | 29     | 43     |          |
|           |               | 35.9%                             | 16.7%  | 20.2%  |          |
|           | Obese II      | 7                                 | 7      | 14     |          |
|           |               | 17.9%                             | 4.0%   | 6.6%   |          |
|           | Total         | 39                                | 174    | 213    |          |
|           |               | 100.0%                            | 100.0% | 100.0% |          |

BMI, WHR and WC all were positively correlated with blood pressure ( $P$  value= .000). (Table 4)

## Discussion

Since last two decades, studies conducted in Pakistan indicate that under and over nutrition both exist in our population. Transition period of childhood to adulthood has an important position in human's life because it is the phase of rapid growth rate. Due to less physical activity and sedentary life styles prevalence of overweight and obesity is rapidly increasing in young medical students. Thus assessing frequency

of underweight, overweight and obesity among adolescents is very important to prevent morbidity and mortality related to them.

Prevalence of obesity and hypertension were determined in medical students of 1<sup>st</sup> year MBBS and 1<sup>st</sup> year BDS. BMI, WC and WHR was used for grouping students into obese and non obese on the basis of South-Asians obesity guidelines<sup>(17)</sup>. BMI, WC and WHR are important indicators for assessing CVD risk factors particularly hypertension.

**Table 4:** Correlation of BMI, WC and WHR with systolic diastolic blood pressure

|                                  |                          | Systolic<br>Blood Pres-<br>sure (mmHg) | Diastolic<br>Blood Pres-<br>sure (mmHg) |
|----------------------------------|--------------------------|----------------------------------------|-----------------------------------------|
| BMI group                        | Pearson Cor-<br>relation | .375 <sup>+</sup>                      | .351 <sup>+</sup>                       |
|                                  | Sig. (2-tailed)          | .000                                   | .000                                    |
|                                  | N                        | 213                                    | 213                                     |
| Waist to<br>Hip Ratio            | Pearson Cor-<br>relation | .250 <sup>+</sup>                      | .265 <sup>+</sup>                       |
|                                  | Sig. (2-tailed)          | .000                                   | .000                                    |
|                                  | N                        | 213                                    | 213                                     |
| Waist cir-<br>cumference<br>(cm) | Pearson Cor-<br>relation | .372 <sup>+</sup>                      | .374 <sup>+</sup>                       |
|                                  | Sig. (2-tailed)          | .000                                   | .000                                    |
|                                  | N                        | 213                                    | 213                                     |

In National Health Survey of Pakistan (NHSP), over-

weight prevalence was 13.5% for males and 19.6% for females that is comparable with our study, while underweight population was 25% in NHSP that is higher than our study in which prevalence of underweight was 4.7%. Mean BMI was 20.9kg/m<sup>2</sup> for males and 21.7kg/m<sup>2</sup> for females according to NHSP while in our study mean BMI in males was 22.9kg/m<sup>2</sup> and in females was 23.6kg/m<sup>2</sup> that is slightly higher than NHSP <sup>(18)</sup>. In another comprehensive study conducted on Pakistani population, the prevalence of obesity, overweight and underweight was 2.2%, 14% and 11%, respectively. Another study conducted by Shams N et al on female medical students <sup>(19)</sup>, there were 47(15%) underweight, 62(20%) overweight and 78(25.4%) obese female students. Change in lifestyle and eating behavior in youngsters in the past decades are the key factors for difference of obesity values in studies. Study on large sample size at national level is needed.

Our study noted another important finding that a very high prevalence 39% of central obesity in medical students existed. Specifically, there were 31(32%) obese males and 52(44.8%) obese females on the basis of WHR. Study by Bertias G, showed prevalence of central obesity as 13% <sup>(20)</sup>. Another study conducted by Afzal M indicated a prevalence of 12.3% that are lower than our study. Strong positive correlation of BMI, WC and WHR with systolic and diastolic blood pressures was shown in our study. It indicates that increase in BMI leads to increase in hypertension in male and female students. The link between these two CVD risk factors has well shown in various studies <sup>(21-22)</sup>. Importance of prevention as well as control of weight and obesity is constant with our study results. These findings also emphasize that measurement of blood pressure and body weight and timely diagnosis and control are essential for overweight and obese people.

## Conclusions

We concluded that high prevalence of overweight and obesity was seen in medical students and it was positively correlated with hypertension. HTN was significantly higher in males as compared to females while obesity was more pronounced in females. It is concluded that medical students should be screened for obesity and hypertension along with dietary advice, implementation of life style modification, and pharmacological intervention if needed. Our study also

suggests that not only being overweight and obesity is a problem of importance among medical students but being under-weight is also a significant problem among students of medical college. Overweight and obese students were advised to do regular exercise, modify their life style and dietary habits to prevent obesity related complications. Underweight students were advised to improve BMI by taking balanced diet.

## Author's Contribution

**Rana Khurram Aftab:** Designed the study, analysed the results and provided references for writing the article.

**Farhat Ijaz:** Approved the study and helped in writing the article.

**and Samia Jawed:** Helped in article-writing and proof reading.

## References

1. Flegal KM, Carroll MD, Kit BK.. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999-2010. *JAMA*. 2012;307(5):491-7. <https://doi.org/10.1001/jama.2012.39>
2. Nguyen T, Lau DCW. Obesity epidemic and its impact on hypertension (Review). *Can J Cardiol*. 2012; 28(3): 326-33. <https://doi.org/10.1016/j.cjca.2012.01.001>
3. Tuck ML, Corry DB. Prevalence of obesity, hypertension, diabetes, and metabolic syndrome and its cardiovascular complications. *Curr Hypertens Rev*. 2010; 6(2): 73-82. <https://doi.org/10.2174/157340210791171010>
4. NananDJ. The Obesity Pandemic- Implications for Pakistan. *J Pak Med Assoc*. 2002;52(8):342-6.
5. Sullivan PW, Ghushchyan VH, Ben-Joseph R. The impact of obesity on diabetes, hyperlipidemia and hypertension in the United States. *Qual Life Res*. 2008; 17(8): 1063-71. <https://doi.org/10.1007/s11136-008-9385-7>
6. Wakabayashi I. Stronger associations of obesity with prehypertension and hypertension in young women than in young men [Abstract]. *J Hypertens*. 2012; 30(7): 1423-9. <https://doi.org/10.1097/HJH.0b013e3283544881>
7. Jordan J, Yumuk V, Schlaich M, Nilsson PM, Zahorska-Markiewicz B, Grassi G, et al. Joint statement of the European Association for the Study of Obesity and the European Society of Hyperten-

- sion: Obesity and difficult to treat arterial hypertension. *J Hypertens*.2012;30(6):1047-55. <https://doi.org/10.1097/HJH.0b013e3283537347>
8. National Heart Lung and Blood institute. The Seventh Report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure (JNC 7) Complete Report; 2004.
  9. Balkau B, Deanfield JE, Després JP, Bassan JP, Fox KAA, Smith SC, et al. International Day for the Evaluation of Abdominal Obesity (IDEA): a study of waist circumference, cardiovascular disease, and diabetes mellitus in 168,000 primary care patients in 63 countries. *Circulation* 2007; 116(17): 1942–51. <https://doi.org/10.1161/CIRCULATIONAHA.106.676379>
  10. Koning L, Merchant AT, Pogue J, Anand SS. Waist circumference and waist-to-hip ratio as predictors of cardiovascular events: meta-regression analysis of prospective studies. *Eur Heart J*. 2007; 28(7): 850–6. <https://doi.org/10.1093/eurheartj/ehm026>
  11. Wenzel UO, Krebs C. Management of arterial hypertension in obese patients. *Curr Hypertens Rep*.2007;9(6):491–97. <https://doi.org/10.1007/s11906-007-0090-7>
  12. Boyle JR, LaRose NR. Personal beliefs, the environment and college students' exercise and eating behaviors. *Am J Health Stud*. 2008;23(4):195–200.
  13. Huang TT, Harris KJ, Lee RE, Nazir N, Born W, Kaur H. Assessing overweight, obesity, diet, and physical activity in college students. *J Am Coll Health*.2003;52(2):83– 6. <https://doi.org/10.1080/07448480309595728>
  14. World Health Organization, Regional Office for the Western Pacific, International Association for the Study of Obesity. International Obesity Task Force. The Asia-Pacific perspective: redefining obesity and its treatment. Melbourne, Health Communications Australia, 2000
  15. World health organization. Waist circumference and waist hip ratio: report of WHO expert consultation. Geneva; 2011:27–34
  16. National Heart Lung and Blood institute. The Seventh Report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure (JNC 7) Complete Report; 2004.
  17. Asia-Pacific Steering Committee. The Asia-Pacific Perspective: Redefining Obesity and Its Treatment; International Diabetes Institute. Health Communications Australia; St Leonards, Australia: 2000;19.
  18. National Health Survey of Pakistan 1990-1994: Pakistan Medical Research Council, 1998.
  19. Shams N, Niaz F, Motwani R, Shaikh Z, Saleem F. Obesity and Hypertension in Female Medical Students; Frequency and Risk Factors. *J Liaquat Uni Med Health Sci*. 2015;14(01):26–32.
  20. Bertsiadis G, Mammas I, Linardakis M and Kafatos A. Overweight and obesity in relation to cardiovascular disease risk factors among medical students in Crete, Greece. *BMC Public Health* 2003; 3(3): 34–39. <https://doi.org/10.1186/1471-2458-3-3>
  21. Hwang LC, Bai CH, Sun CA, Chen CJ. Prevalence of metabolically healthy obesity and its impacts on incidences of hypertension, diabetes and the metabolic syndrome in Taiwan. *Asia Pac J Clin Nutr*. 2012; 21(2): 227–33.
  22. Warsy AS, El-Hazmi MAF, Al-Hazmi AM. Prevalence of co-existing hypertension and obesity in Saudis. *Biomedical and Pharmacology J*. 2011; 4(1): 269–74. <https://doi.org/10.13005/bpj/293>