

GENUS *SORBUS* L.: A PHYTOCHEMICAL REVIEW

Sadia Khan^{1*}, Mehdi Hassan Kazmi¹, Farah Inamullah¹, Shagufta Afaq¹, M. Zahid Farhad¹, Sadia Ferheen² and Sarwat Ismail³

¹Department of Applied Chemistry, University of Karachi, Karachi-75270, Pakistan

²Pharmaceutical Research Centre, PCSIR Laboratories Complex Karachi, Pakistan

³PCSIR, Head office, 1-Constitution Avenue, G-5/2, Islamabad, Pakistan

*corresponding author: sakhan@uok.edu.pk

ABSTRACT

The current review is focused on the phytochemical and pharmacological aspects of the genus *Sorbus*. Its numerous species have been utilized as a traditional medicine in the world. Despite, the therapeutic importance of genus *Sorbus*, out of 100 species only 13 species have been screened phytochemically. Up till now, around 150 secondary metabolites have been reported from this genus, which are summarized in this review.

Keywords: Rosaceae, *Sorbus*, phytochemical, pharmacological potential.

INTRODUCTION

The *Sorbus* L. is a genus from the Rose family (Rosaceae). *Sorbus* comprises of 100 species among which many are medicinally important and are being used in the folklore medications all over the world (Bhattacharjee, 2003). Several taxa from *Sorbus* are found in customary and local medicines that are used as anti-diarrhea, diuretic, anti-inflammatory, anti-diabetic, vaso- protective, broncho- and vaso-relaxant, along with potent antioxidative qualities (Hukkanen *et al.*, 2006, Olszewska and Michel, 2009, Olszewska *et al.*, 2010, Olszewska, 2012; Perry and Metzger, 1980; Krishna, 1972; Jayaweera, 1982; Krachmal, 1980).

The fruit of *S. aucuparia* L. is bitter in nature and having characteristics of, astringent, laxative, diuretic and cholagogue. Traditionally these are used as drugs in the form of tea, syrup, jelly or liqueur in Austria for the treatment of ailments related to respiratory tract, fever, infections, colds, flu, rheumatism and gout. The bark of *S. aucuparia* L. is used as astringent for the treatment of diarrhea and leucorrhoea as a vaginal injection. Its infusion is also used for the cure of constipation, kidney disorders and painful menstruation (Gillham, 1980; McAllister, 1996; Sperens, 1997; Vogl *et al.*, 2013).

S. commixta Hedl is used as medication for asthma, cough, and other bronchial disorders. The plant exhibits promising antioxidative, anti-inflammatory, anti-atherosclerotic, anti-athero genic and vascular relaxant effects. It decreases hepatic lipid peroxidation by lowering the bioavailability of alcohol and its oxidative metabolites by the protection of hepatic catalase (Gabsik and Hyo-Jin, 2014). Ethnomedicinal uses of *S. amurens* varies with the plant parts such as its bark is Koehne and is used to cure nausea, cough, cleansing of blood and heart diseases whereas the fruits are used to treat severe wounds, stroke, and heart disease, gall bladder irritation and digestive disturbance (Bhattacharjee, 2003). The infusion of the leaves to some extent is laxative and used as pectoral in cough and bronchitis (Krachmal, 1980). Uses of *S. decora* C.K. Schneid is associated with the diabetes mellitus (Guerro-Analco *et al.*, 2010). The bark of *S. cashmiriana* Hedlung, (Monog tree) is considered to be helpful in controlling nausea and its fruit is used to cure scurvy and a rich source of vitamin C (Kazmi *et al.*, 2009). Traditionally, the fruits of *S. domestica* L. have antioxidant properties, and used as a remedy against long term diabetic complications.

Most of the *Sorbus* species depicts antioxidant behavior due to the presence of phenolic constituents, which are mainly phenolic acid, esters with quinic acid, flavonols (quercetin, kaempferol and sexangularetin), anthocyanins (cyaniding and pelargonidin derivatives) and tannin-type proanthocyanidins (Hukkanen *et al.*, 2006, Termentzi *et al.*, 2006, Olszewska and Michel, 2009, Olszewska, 2012). Presence of phenolic compounds in the fruits of *S. domestica* causes aldose reductase inhibitory activity which reduces diabetic complications (Termentzi *et al.*, 2008). In Xinjiang, *S. tianschanica* Rupr has been used as a pharmacologically important plant for the treatment of tuberculosis, asthma, cough and gastritis for a long time (Yu *et al.*, 2012). Various parts of the *S. pohuashanensis* Hance such as fruits, stems and bark have been extensively used in Chinese ethno- medicine for the treatment of edema, tuberculosis and chronic tracheitis (Huiyong *et al.*, 2012). Genus *Sorbus* L. contains Caffeoylquinic acid possess cardio protective, neuro protective, antidiabetic and hepatoprotective effects (Zymone *et al.*, 2018).

Secondary Metabolites

From 13 species of the genus *Sorbus*, 180 secondary metabolites have been report up till now. These compounds are presented in Table1.

Table 1. Summarized account of various secondary metabolites reported from *Sorbus* species up to Dec 2018.

Species	Compounds	References
<i>S. aucuparia</i>	Sorbikortal I	Lawrie <i>et al.</i> (1960)
	Sorbikortal II	Lawrie <i>et al.</i> (1960)
	β -Xyloside	Arya <i>et al.</i> (1962)
	S-Parasorbic acid	Wilhelm (1963)
	Isoquercitrin	Borisov and Zhuravl'ov (1965)
	Spireoside	Borisov and Zhuravl'ov (1965)
	Ursolic acid	Shavva <i>et al.</i> , (1969)
	Sexangularetin 3-glucoside	Jerzmanowska and Kamecki (1973)
	Sorbic acid	Pyysalo and Kuusi (1974)
	Dihydrosinapic aldehyde	Malterud and Opheim (1989)
	Aucuparin	Kokubun <i>et al.</i> (1995)
	2'-Methoxyaucuparin	Kokubun <i>et al.</i> (1995)
	4'-Methoxyaucuparin	Kokubun <i>et al.</i> (1995)
	2'-Hydroxyaucuparin	Kokubun <i>et al.</i> (1995)
	Isoaucuparin	Kokubun <i>et al.</i> (1995)
	L-Iditol	Izumori (2006)
	Caproic acid	Krivoruchko <i>et al.</i> (2013)
	3-Hexenoic acid	Krivoruchko <i>et al.</i> (2013)
	Caprylic acid	Krivoruchko <i>et al.</i> (2013)
	Oxalic acid	Krivoruchko <i>et al.</i> (2013)
	Pelargonic acid	Lawrie <i>et al.</i> (1960)
	Malonic acid	Lawrie <i>et al.</i> (1960)
	Fumaric acid	Arya <i>et al.</i> (1962)
	Succinic acid	Wilhelm (1963)
	Capric acid	Borisov and Zhuravl'ov (1965)
	Benzoic acid	Borisov and Zhuravl'ov (1965)
	Methoxysuccinic acid	Shavva <i>et al.</i> (1969)
	Phenylacetic acid	Jerzmanowska and Kamecki (1973)
	Salicylic acid	Pyysalo and Kuusi (1974)
	Lauric acid	Malterud and Opheim (1989)
	Myristic acid	Kokubun <i>et al.</i> (1995)
	Pentadecanoic acid	Kokubun <i>et al.</i> (1995)
	Azelaic acid	Kokubun <i>et al.</i> (1995)
	Palmitic acid	Kokubun <i>et al.</i> (1995)
	Palmitoleic acid	Kokubun <i>et al.</i> (1995)
	Margaric acid	Izumori (2006)
	Citric acid	Krivoruchko <i>et al.</i> (2013)
	Stearic acid	Krivoruchko <i>et al.</i> (2013)
	Oleic acid	Krivoruchko <i>et al.</i> (2013)
	Vaccenic acid	Krivoruchko <i>et al.</i> (2013)
	Linoleic acid	Krivoruchko <i>et al.</i> (2013)
	Linolenic acid	Krivoruchko <i>et al.</i> (2013)
	Arachic acid	Krivoruchko <i>et al.</i> (2013)
	11-Eicosenic acid	Krivoruchko <i>et al.</i> (2013)
	11,13-Eicosadienoic acid	Krivoruchko <i>et al.</i> (2013)
	Heneicosanoic acid	Krivoruchko <i>et al.</i> (2013)
	Behenic acid	Krivoruchko <i>et al.</i> (2013)
	Erucic acid	Krivoruchko <i>et al.</i> (2013)
	Lignocerinic acid	Krivoruchko <i>et al.</i> (2013)
	Ferulic acid	Krivoruchko <i>et al.</i> (2013)
	Kerotinic acid	Krivoruchko <i>et al.</i> (2013)

(Continued)

Table 1.(Continued).

Species	Compounds	References
<i>S. aria</i>	Ursolic acid	Rolland and Raynaud (1979)
	Oleanolic acid	Rolland and Raynaud (1979)
	Quercetin acid	Rolland and Raynaud (1979)
	Cyanidin acid	Rolland and Raynaud (1979)
	Leucocyanidin	Rolland and Raynaud (1979)
	Isorhamnetin 3-O- β -D-glucopyranoside	Olszewska and Michel (2012)
	Astragalin	Olszewska and Michel (2012)
	Isoquercitrin	Olszewska and Michel (2012)
	Hyperoside	Olszewska and Michel (2012)
	Kaempferol 3-O- β -D-glucopyranoside-7-O- α -rhamnopyranoside	Olszewska and Michel (2012)
	Quercetin 3-O- β -D-glucopyranoside-7-O- α -rhamnopyranoside	Olszewska and Michel (2012)
	Rutin	Olszewska and Michel (2012)
	Chlorogenic acid	Olszewska and Michel (2012)
	Neochlorogenic acid	Olszewska and Michel (2012)
	Caproic acid	Krivoruchko <i>et al.</i> (2013)
	3-Hexenoic acid	Krivoruchko <i>et al.</i> (2013)
	2-Hexenoic acid	Krivoruchko <i>et al.</i> (2013)
	Oxalic acid	Krivoruchko <i>et al.</i> (2013)
	Malonic acid	Krivoruchko <i>et al.</i> (2013)
	Fumaric acid	Krivoruchko <i>et al.</i> (2013)
	Succinic acid	Krivoruchko <i>et al.</i> (2013)
	Benzoic acid	Krivoruchko <i>et al.</i> (2013)
	Lauric acid	Krivoruchko <i>et al.</i> (2013)
	Myristic acid	Krivoruchko <i>et al.</i> (2013)
	Pentadecanoic acid	Krivoruchko <i>et al.</i> (2013)
	Azelaic acid	Krivoruchko <i>et al.</i> (2013)
	Palmitic acid	Krivoruchko <i>et al.</i> (2013)
	Palmitoleic acid	Krivoruchko <i>et al.</i> (2013)
	Margaric acid	Krivoruchko <i>et al.</i> (2013)
	Citric acid	Krivoruchko <i>et al.</i> (2013)
	Stearic acid	Krivoruchko <i>et al.</i> (2013)
	Oleic acid	Krivoruchko <i>et al.</i> (2013)
	Linoleic acid	Krivoruchko <i>et al.</i> (2013)
	Linolenic acid	Krivoruchko <i>et al.</i> (2013)
	Arachic acid	Krivoruchko <i>et al.</i> (2013)
	Behenic acid	Krivoruchko <i>et al.</i> (2013)
	Lignoceric acid	Krivoruchko <i>et al.</i> (2013)
	Ferulic acid	Krivoruchko <i>et al.</i> (2013)
<i>S. cashmiriana</i>	Sorbinol A	Kazmi <i>et al.</i> (2007)
	Sorbinol B	Kazmi <i>et al.</i> (2007)
	Cashmirol A	Kazmi <i>et al.</i> (2007)
	Cashmirol B	Kazmi <i>et al.</i> (2007)
	Sorbicin A	Kazmi <i>et al.</i> (2007)
	Sorbicin B	Kazmi <i>et al.</i> (2007)
	Sorbusin	Rizvi <i>et al.</i> (2013)
	Lyonoside	Rizvi <i>et al.</i> (2013)
	β -Sitosterol	Rizvi <i>et al.</i> (2013)
	Benzyl- β -D-glucoside	Rizvi <i>et al.</i> (2013)
	Methyl- β -D-glucoside	Rizvi <i>et al.</i> (2013)
	2(R)-Prunasin	Rizvi <i>et al.</i> (2013)
	(2R/2S)-Prunasin	Rizvi <i>et al.</i> (2013)

(Continued)

Table 1.(Continued).

Species	Compounds	References
	Betulin	Rizvi <i>et al.</i> (2013)
	Anemosapogenin	Rizvi <i>et al.</i> (2013)
	Ursolic acid	Rizvi <i>et al.</i> (2013)
	Rutin	Rizvi <i>et al.</i> (2013)
	Isoquercitrin	Rizvi <i>et al.</i> (2013)
	Ursolic acid	Khan <i>et al.</i> (2013)
	Stigmasterol	Khan <i>et al.</i> (2013)
	Myricadiol	Khan <i>et al.</i> (2013)
	Taraxerol	Khan <i>et al.</i> (2013)
	Butanolic acid	Khan <i>et al.</i> (2013)
	Betulinic acid	Khan <i>et al.</i> (2013)
	5 α , 8 α -Epidioxyergosta-6, 22-diene-3 β -ol	Khan <i>et al.</i> (2013)
	3 β , 5 α -Dihydroxy-6- β -methoxy- ergosta-7, 22-diene	Khan <i>et al.</i> (2013)
	β -Sitosteryl acetate	Khan <i>et al.</i> (2013)
	5 α , 8 α -Epidioxyergosta-6, 9 (11),22-trien-3 β -ol	Rizvi <i>et al.</i> (2013)
	Stigmasterol 3-O- β -D-glucopyrano- side	Rizvi <i>et al.</i> (2013)
	Cashmin A	Khan <i>et al.</i> (2015)
	Cashmin B	Khan <i>et al.</i> (2015)
<i>S. commixta</i>	Sorbic acid	Asahina and Shinoda (1930)
	Sorbitol (II)	Asahina and Shinoda (1930)
	Prunasin	Takaishi and Kuwajima (1976)
	Amygdalin	Takaishi and Kuwajima (1976)
	Luteolin 7-glucoside	Bylka <i>et al.</i> (1977)
	Quercetin 3-glucoside	Bylka <i>et al.</i> (1977)
	Quercetin 3-glucosyloside	Bylka <i>et al.</i> (1977)
	Quercetin	Bylka <i>et al.</i> (1977)
	Catechin 7-apioside	Na <i>et al.</i> (2002)
	Catechin 7-xyloside	Na <i>et al.</i> (2002)
	Lupenone	Na <i>et al.</i> (2009)
	Lupeol	Na <i>et al.</i> (2009)
	Chalcone glycoside	Bhatt <i>et al.</i> (2009)
	β -sitosteryl-3-O- β -D-glucopyrano- side	Gabsik and Hyo-Jin (2014)
	1,2,4-trimethoxydibenzofuran-3,9-diol	Choi <i>et al.</i> (2018)
	Sorcomisides A	Kim <i>et al.</i> (2018)
	Sorcomisides B	Kim <i>et al.</i> (2018)
<i>S. cortex</i>	Lupenone	Lee and Lee (1999)
	Lupeol	Lee and Lee (1999)
<i>S. decora</i>	Aucuparin	Kokubun <i>et al.</i> (1995)
	2'-Methoxyaucuparin	Kokubun <i>et al.</i> (1995)
	3 β , 23, 28-Trihydroxy-12-ursene	Guerro-Analco <i>et al.</i> (2010)
	23, 28-Dihydroxyursan-12-ene-3 β -caffeate	Guerro-Analco <i>et al.</i> (2010)
	23, 28-Dihydroxylupan-20(29)-ene-3 β -caffeate	Guerro-Analco <i>et al.</i> (2010)
	23-Hydroxybetulin	Guerro-Analco <i>et al.</i> (2010)
	Uvaol	Guerro-Analco <i>et al.</i> (2010)
	Betulin	Guerro-Analco <i>et al.</i> (2010)
	α -Amyrin	Guerro-Analco <i>et al.</i> (2010)
	Betulinic acid	Guerro-Analco <i>et al.</i> (2010)
	(+)-Catechin	Guerro-Analco <i>et al.</i> (2010)
	(-)-Epicatechin	Guerro-Analco <i>et al.</i> (2010)
<i>S. domestica</i>	Vannilic acid 4-O- α -L-rhamnopyra- noside	Termentzi <i>et al.</i> (2009)
	Protocateuic acid anhydride	Termentzi <i>et al.</i> (2009)

(Continued)

Table 1.(Continued).

Species	Compounds	References
	Trivanilloyl-(1,3,4-trihydroxybenzol) ester	Termentzi <i>et al.</i> (2009)
	Quercetin 3-O-β-D-glucopyranosyl (1"→2")-α-L-rhamnosyl	Termentzi <i>et al.</i> (2009)
	3-{4-(Bis[4-hydroxy-3-(5-hydroxy-pentanoyloxy)phenyl]-methoxy]-3,5-dihydroxyphenyl}propanoic acid	Termentzi <i>et al.</i> (2009)
	(1"→3")-α-L-rhamnosyl(1"→3")-α-L-arabinofuranoside	Termentzi <i>et al.</i> (2009)
	Quercetin 3-O-α-L-rhamnosyl- (1"→3")-β-D-glucopyranoside	Termentzi <i>et al.</i> (2009)
	5,7,3',6'-tetrahydroxyflavanol 7-O-β-D-glucopyranoside	Termentzi <i>et al.</i> (2009)
	(7-O-4",4'-O-7") Quercetin dimer	Termentzi <i>et al.</i> (2009)
	[2,2'-dihydroxy, 4-(propionic acid hexyl ester), 4'-(propionic acid heptyl ester)] biphenyl	Termentzi <i>et al.</i> (2009)
	[2,6,2',6'-tetrahydroxy,4,4'-bis-(propionic acid hexyl ester)] biphenyl	Termentzi <i>et al.</i> (2009)
<i>S. lanata</i>	β-Pyrufuran	Yousuf <i>et al.</i> (2010)
	1,2,4-Trimethoxydibenzo[<i>b,d</i>]furan-3-ol	Yousuf <i>et al.</i> (2010)
	Sorlanin	Uddin <i>et al.</i> (2013)
	Sorbanin	Uddin <i>et al.</i> (2013)
	Sorbalanin	Uddin <i>et al.</i> (2013)
	Polystachyol	Uddin <i>et al.</i> (2013)
	Isolariciresinol	Uddin <i>et al.</i> (2013)
	Dihydrodehydrodiconiferyl alcohol	Uddin <i>et al.</i> (2013)
	Tuberculation	Uddin <i>et al.</i> (2013)
	Ovafofinin E	Uddin <i>et al.</i> (2013)
	Aucuparin	Uddin <i>et al.</i> (2013)
	2'-Methoxyaucuparin	Uddin <i>et al.</i> (2013)
	Tetracosyl-3-(3,4-dihydroxyphenyl)- acrylate	Uddin <i>et al.</i> (2013)
	SorbanolicAcid	Latif <i>et al.</i> (2014)
	3β,23-Dihydroxy-lup-20(29)ene-28-oic acid-23-caffeate	Latif <i>et al.</i> (2014)
	3β,23-Dihydroxy-lup-20(29)ene-28-oic acid-3β-caffeate	Latif <i>et al.</i> (2014)
	Lyoniside	Latif <i>et al.</i> (2014)
<i>S. pendula</i>	Quercetin	Pavlai <i>et al.</i> (1966)
	Galactose	Pavlai <i>et al.</i> (1966)
	Hyperoside	Pavlai <i>et al.</i> (1966)
	Sorboside	Makarova (1967)
	Kaempferol 3-gentiobioside	Wagner <i>et al.</i> (1968)
	Quercetin 3-gentiobioside	Wagner <i>et al.</i> (1968)
<i>S. pohuashanensis</i>	Stigma-5-en-3-O-β-glucoside	Huiyong <i>et al.</i> (2012)
	Quercetin	Huiyong <i>et al.</i> (2012)
	Rutin	Huiyong <i>et al.</i> (2012)
	1-Caffeoylquinic acid	Huiyong <i>et al.</i> (2012)
	Phenylmethanol α-L-arabinofuran- osyl-(1/6)-β-D-glucopyranoside	Huiyong <i>et al.</i> (2012)

(Continued)

Table 1.(Continued).

Species	Compounds	References
	Prunasin	Huiyong <i>et al.</i> (2012)
	Amygdalin	Huiyong <i>et al.</i> (2012)
	Butanedioic acid	Huiyong <i>et al.</i> (2012)
	(3S,5S)-3-(β -D-glucopyranosyloxy)-5-hydroxyhexonic acid ethyl ester	Huiyong <i>et al.</i> (2012)
	Parasorboside	Huiyong <i>et al.</i> (2012)
	1-Glycerol linolate	Huiyong <i>et al.</i> (2012)
	3S,5S)-3-(β -D-glucopyranosyloxy)-5-hydroxyhexonic acid ethyl ester	Huiyong <i>et al.</i> (2012)
	Parasorboside	Huiyong <i>et al.</i> (2012)
	1-Glycerol linolate	Huiyong <i>et al.</i> (2012)
	Pomolic acid	Li <i>et al.</i> (2014)
	Pomolic acid-3 β -acetate	Li <i>et al.</i> (2014)
	3 β -Acetoxy-urs-12-ene-28-oic acid	Li <i>et al.</i> (2014)
	Ursolaldehyde	Li <i>et al.</i> (2014)
	Euscaphic acid	Li <i>et al.</i> (2014)
	3 β -Acetoxy-urs-11-en-28,13-olide	Li <i>et al.</i> (2014)
	Betulinic acid	Li <i>et al.</i> 92014)
<i>S. quercifolia</i>	Quercetin 3-gentiobioside	Wagner <i>et al.</i> (1968)
	Quercetin 3- β -gentiobioside	Pavlii and Makarova (1970)
	Quercetin 3- β -D-galactopyranoside	Pavlii and Makarova (1970)
	Quin 3-rutinoside	Pavlii and Makarova (1970)
	Apigenin 7-glucoside	Pavlii and Makarova (1970)
	Caffeic acid	Pavlii and Makarova (1970)
	Chlorogenic acid	Pavlii and Makarova (1970)
	Rutin	Pavlii and Makarova (1970)
	Hyperin	Pavlii and Makarova (1970)
	Chlorogenic acid	Pavlii and Makarova (1970)
	Caffeic acid	Pavlii and Makarova (1970)
	Neochlorogenic acid	Pavlii and Makarova (1970)
<i>S. tianschanica</i>	Quirzutrin	Zapesochnaya <i>et al.</i> (1973)
	Chlorogenic acid	Zapesochnaya <i>et al.</i> (1973)
	Benzoic acid	Chang <i>et al.</i> (2009)
	Benzyl-O- β -D-glucopyranoside	Chang <i>et al.</i> (2009)
	Ursolic acid	Chang <i>et al.</i> (2009)
	2 α -Hydroxyursolic acid	Chang <i>et al.</i> (2009)
	Hyperoside	Chang <i>et al.</i> (2009)
	Quercetin-3-O-glucoside	Chang <i>et al.</i> (2009)
	Chlorogenic acid	Yu <i>et al.</i> (2012)
	Quercetin-3-O-(6"-O-malonyl)- β -D-glucoside	Yu <i>et al.</i> (2012)
	Kaempferol-3-O-(6"-O-malonyl)- β -D-glucopyranoside	Yu <i>et al.</i> (2012)
	Kaempferol-3-O- β -D-glucopyranoside	Yu <i>et al.</i> (2012)
<i>S. torminalis</i>	Ursolic acid	Ivanovskaya <i>et al.</i> (1963)
	Cholesterol	Tsitsa-Tzardi <i>et al.</i> (1991)
	Campesterol	Tsitsa-Tzardi <i>et al.</i> (1991)
	Stigmasterol	Tsitsa-Tzardi <i>et al.</i> (1991)
	Sitosterol	Tsitsa-Tzardi <i>et al.</i> (1991)
	Myristic acid	Tsitsa-Tzardi <i>et al.</i> (1991)
	Palmitic acid	Tsitsa-Tzardi <i>et al.</i> (1991)
	Palmitoleic acid	Tsitsa-Tzardi <i>et al.</i> (1991)
	Stearic acid	Tsitsa-Tzardi <i>et al.</i> (1991)

(Continued)

Table 1.(Continued).

Species	Compounds	References
	Oleic acid	Tsitsa-Tzardi <i>et al.</i> (1991)
	Linoleic acid	Tsitsa-Tzardi <i>et al.</i> (1991)
	Linolenic acid	Tsitsa-Tzardi <i>et al.</i> (1991)
	p-Coumaric acid	Tsitsa-Tzardi <i>et al.</i> (1992)
	Caffeic acid	Tsitsa-Tzardi <i>et al.</i> (1992)
	Quercetin	Tsitsa-Tzardi <i>et al.</i> (1992)
	Chlorogenic acid	Tsitsa-Tzardi <i>et al.</i> (1992)
	Luteolin-7-O-glycoside	Tsitsa-Tzardi <i>et al.</i> (1992)
	Apigenin-7-O-glycoside	Tsitsa-Tzardi <i>et al.</i> (1992)
	Tormaloside	Olszewska and Roj (2011)
	Hyperoside	Olszewska and Roj (2011)
	Isoquercitrin	Olszewska and Roj (2011)
	Chlorogenic acid	Olszewska and Roj (2011)
	Neochlorogenic acid	Olszewska and Roj (2011)
	5,7,4'-Trihydroxy-3'-methoxyflavone-7-O- β -D-glucopyranoside	Olszewska and Roj (2011)
	3,5,7,4'-Tetrahydroxy-8,3'-dimethoxy- flavone-3-O- β -D-glucopyranoside	Olszewska and Roj (2011)
	3,5,7,4'-Tetrahydroxy-8-methoxy- flavone-3-O- β -D-glucopyranoside	Olszewska and Roj (2011)
	3,5,7,4'-Tetrahydroxy-3'-methoxy- flavone-3-O- β -D-galactopyranoside	Olszewska and Roj (2011)
	3,5,7,4'-Tetrahydroxy-3'-methoxy- flavone-3-O- β -D-glucopyranoside	Olszewska and Roj (2011)

Pharmacological studies

Studies on the sapwood tissue of *S. aucuparia* depicted that aucuparin and its derivatives are basically not present in the healthy tissues, and are only formed as phytoalexins after the fungal infection (Kokubun *et al.*, 1995). A significant antidiabetic activity has been elucidated by 23, 28-Dihydroxylupan-20(29)-ene-3 β -caffeate which was isolated from *S. decora* (Guerro-Analco *et al.*, 2010). The lupene type triterpenes sorbicins A and B extracted from *S. cashmiriana* illustrated strong inhibitory activity against the enzymes urease and α -chymotrypsin (Kazmi *et al.*, 2011). The triterpenes Cashmirols A and B are isolated from *S. cashmiriana*, are lanostane type which displayed characteristics of inhibitory potential enzyme, lipoxygenase (Kazmi *et al.*, 2009). The phenolic secondary metabolites namely tuberculatin, dihydrodehydrodiconiferyl alcohol, 20-methoxyaucuparin and tetracosyl-3-(3,4-dihydroxyphenyl) acrylate, isolated from *S. lanata* (D. Don.) Schauer have expressed radical scavenging activity through DPPH assay (Uddin *et al.*, 2013). Significant antioxidant properties were also observed in Sorbanolic acid along with three known compounds 3 β , 23-dihydroxy-lup-20(29)ene-28-oic acid-23-caffeate, 3 β , 23-dihydroxy-lup-20(29)ene-28-oic acid-3 β -caffeate and lyoniside attained from *S. lanata* (Latif *et al.*, 2014). Two more triterpenes, lupenone and lupeol extracted from *S. commixta* exhibited inhibition of protein tyrosine phosphatase 1B (PTP1B) (Na *et al.*, 2009). Likewise, β -sitosteryl-3-O- β -glucopyranoside also obtained from *S. commixta* possess anti-inflammatory activity (Gabsik and Hyo-Jin, 2014). Moreover, following compounds isolated from *S. aria* (L.) Crantz have also shown strong free radical-scavenging activity; isoquercitrin, quercetin 3-O- β -glucopyranoside-7-O- α -rhamnopyranoside, rutin, chlorogenic acid and neochlorogenic acid (Olszewska and Michel, 2012).

CONCLUSIONS

Hence it is concluded from the foregoing brief that genus *Sorbus* has tremendous phytochemical and pharmacological value. Nevertheless, still numerous species have not been analyzed in respect to systematic phytochemical screening.

REFERENCES

- Arya, V.P., H. Erdtman, M. Krolikowska and T. Norin (1962). A Lignanxyloside from the sapwood of *Sorbus aucuparia*. *Acta Chem. Scand.*, 16: 518-520
- Asahina, Y. and H. Shinoda (1930). Constituents of fruits of *Sorbus commixta*, Hedlund. *Yakugaku Zasshi.*, 50: 1-7.

- Bhatt, L.R., M. S. Bae, B.M. Kim, G.S. Oh and K.Y. Chai (2009) A chalcone glycoside from the fruits of *Sorbus commixta* Hedl. *Molecules*, 14: 5323-5327.
- Bhattacharjee, S.K. (2003). *Hand Book of Medicinal Plants*, Pointer Publisher, Jaipur India, p. 30
- Borisov, M.I. and M.S. Zhurav'ov (1965). Flavonoids of *Sorbusaucuparia*. *Farm.Zh. (Kiev)*, 20: 50-52.
- Bylka, W., Z. Kowalewski and K.Oleksiak (1977). Flavonoid compounds in fruits of *Sorbus commixta*. *Herba Polonica*, 23: 105-110.
- Chang, J., L. Yang and K. Rena (2009). Studies on chemical constituents from EtOAc fraction of *Sorbustianschanica*. *Zhongguo Zhong Yao ZaZhi*, 34: 175-176.
- Choi, seong yeon, B. Jeong, H. S. Jang, J.L.K. Ko, H. Kim, J. Hang, Y. S. Bae and H. Yang (2018). A new Dibenzofuran from the bark of *Sorbus commixta*, *Rec.Nat.Prod.*, 12 (2): 179-183.
- Gabsik, Y. and A. Hyo-Jin (2014). β -sitosteryl-3-O- β -glucopyranoside isolated from the bark of *Sorbuscommixta* ameliorates pro-inflammatory mediators in RAW 264.7 macrophages. *Immunopharmacol.Immunotoxicol.*, 36: 70-77.
- Gillham, C.M. (1980). The Biology of Rowan (*Sorbus aucuparia* L.). The ecology of *Sorbus aucuparia*, taxonomy of *Sorbus* section aucuparia, and the use of those species as amenity trees. MSc Thesis, University of Liverpool, Liverpool, UK.
- Guerro-Analco, J.A., L. Martineau, A. Saleem, P. Madiraju, A. Muhammad, T. Durst, P. Haddard and J.T. Arnason (2010). Bioassay-guided isolation of the antidiabetic principle from *Sorbus decora* (Rosaceae) used traditionally by the EeyouIstchee Cree First Nations. *J. Nat. Prod.*, 73: 1519-1523.
- Huiyong, L., M. Mariko, L. Wei, L. Qin, Z. Qingbo and K. Kazuo (2012). Chemical constituents from the fruits of *Sorbus pohuashanensis*. *Biochem.Sys. Ecol.*, 43: 166-168.
- Hukkanen, A.T., S.S. Pölönen, S.O. Kärenlampi and H.I.Kokko (2006).Antioxidant capacity and phenolic content of sweet rowanberries. *J. Agric. Food Chem.*,54: 112–119.
- Ivanovskaya, L.Y., A.A. Ryabinin and L.P. Barabanova (1963). Ursolic acid in plants. *ZhurnalObshcheiKhimii*, 33: 3446-3447.
- Izumori, K. (2006). A strategy for bioproduction of all hexoses. *J. Biotechnol.*, 124: 717-722.
- Jayaweera, D.M.A. (1982). *Medicinal Plants*, N. S. C of Sri Lanka, 4715 Maitland Place Colombo, Vol. 4, p. 257
- Jerzmanowska, Z. and J. Kamecki (1973). Phytochemical analysis of inflorescence of mountain ash (rowan tree) – *Sorbusaucuparia* L. Part II.*Rocz. Chem.*, 47: 1629-1638.
- Kazmi, M.H., E. Ahmed, S. Hameed, A. Malik and M. Ashraf (2007). Isolation and structural determination of sorbinols A and B, new triterpenes from *Sorbus cashmariana*, by 1D and 2D NMR spectroscopy. *Magn.Reson. Chem.*, 45: 416-419.
- Kazmi, M.H., E. Ahmed, S. Hameed, A. Malik, I. Fatima and M. Ashraf (2009). Cashmirols A and B, new lipoxygenase inhibiting triterpenes from *Sorbus cashmiriana*. *Chem. Biodiversity*, 6: 1471-1476.
- Kazmi, M.H., I. Fatima, A. Malik, L. Iqbal, M. Latif and N. Afza (2011). Sorbicins A and B, new urease and serine protease inhibitory triterpenes from *Sorbus cashmiriana*. *J. AsianNat. Prod. Res.*, 13: 1081-1086.
- Khan, S., M.H. Kazmi, E. Ahmed, A. Malik and A. Sharif (2013). Phytochemical studies on *Sorbus cashmiriana*. *J. Chem. Soc. Pak.*, 35: 130-134.
- Khan, sadia; F. Itrat, K. Mehdi, M. Abdul, A. Afza, I. Lubna and L. Mehreen (2015). Cashmin A and B, potent antioxidant coumarins from *Sorbus cashmiriana*. *Chem. Nat Comps.*, 4: 626-629.
- Kim, Chung sub, J. Oh, L. Subedi, S. Y. Kim, S. U. Choi and K. R. Lee (2018). Two new phenolic glycosides from *Sorbus commixta*. *Chem.Pharm.Bull.*, 66: 839-842.
- Kokubun, T., J.B. Harborne, J. Eagles and P.G. Waterman (1995). Antifungal biphenyl compounds are the phytoalexins of the sapwood of *Sorbus aucuparia*. *Phytochemistry*, 40: 57-59.
- Krachmal, C. (1980). *A Guide to the Medicinal Plants of United States*, Quadrangle New York Times Book Co., pp. 9, 207
- Krishna, A. (1972). *The Wealth of India*, CSIR, New Delhi, Vol. 9, p. 435.
- Krivoruchko, E.V., O.A. Andrushchenko and A.V. Kononenko (2013). Carboxylic Acids from *Sorbusaucuparia* and *S. aria*.*Chem. Nat. Comp.*, 49: 742-743.
- Latif, A., S.H. Hussain, M. Ali, M. Arfan, M. Ahmed, R.J. Cox, T.J. Simpson and G. Uddin (2014). A new antioxidant triterpenoid from the stem wood of *Sorbus lanata*.*Rec. Nat. Prod.*,8: 19-24.
- Lawrie, W., J. McLean and G.R. Taylor (1960). Triterpenoids in the bark of mountain ash (*sorbus aucuparia* L.). *J. Chem. Soc.*, 4303-4308.
- Lee, S.M. and C.G. Lee (1999). Isolation and gas chromatographic analysis of lupenone and lupeol from *Sorbus cortex*. *Anal. Sci., Tech.*, 12: 136-140.

- Li, D., W. Li, K. Higai and K. Koike (2014). Protein tyrosine phosphatase 1B inhibitory activities of ursane- and lupane-type triterpenes from *Sorbus pohuashanensis*. *J. Nat. Med.*, 68: 427-431.
- Makarova, H.V. (1967). A study of *Sorbus pendula* flavonoids. *Farm.Zh. (Kiev)*, 22: 50-53
- Malterud, K.E. and K. Opheim (1989). 3-(4-Hydroxy-3,5-dimethoxyphenyl)-propanal from *Sorbus aucuparia* sapwood. *Phytochemistry*, 28: 1548-1549.
- McAllister, H.A. (1996). *Sorbus*: mountain ash and its relatives. *The Garden, J. R. Hortic. Soc.*, 121: 561-567.
- Na, M., B.Y. Kim, H. Osada and J.S. Ahn (2009). Inhibition of protein tyrosine phosphatase 1B by lupeol and lupenone isolated from *Sorbus commixta*. *J. Enzyme Inhib. Med. Chem.*, 24: 1056-1059.
- Na, M.K., R.B. An, S.M. Lee, B.S. Min, Y.H. Kim, K.H. Bae and S.S. Kang (2002). Antioxidant compounds from the stem bark of *Sorbus commixta*. *Nat. Prod. Sci.*, 8: 26-29.
- Olszewska, M.A. (2012). *In vitro* antioxidant activity and total phenolic content of the in florescences, leaves and fruits of *Sorbus torminalis* (L.) Crantz. *Acta. Pol. Pharm.*, 68: 945-953.
- Olszewska, M.A. and P. Michel (2009). Antioxidant activity of in florescences, leaves and fruits of three *Sorbus* species in relation to their polyphenolic composition. *Nat. Prod. Res.*, 23: 1507-1521.
- Olszewska, M.A. and P. Michel (2012). Activity-guided isolation and identification of free radical scavenging components from various leaf extracts of *Sorbus aria* (L.) Crantz. *Nat. Prod. Res.*, 26: 243-254.
- Olszewska, M.A., S. Nowak, P. Michel, P. Banaszczyk and A. Kicel (2010). Assessment of the content of phenolics and antioxidant action of inflorescences and leaves of selected species from the genus *Sorbus sensu stricto*. *Molecules*, 15: 8769-8783.
- Olszewska, M.A. and J.M. Roj (2011). Phenolic constituents of the inflorescences of *Sorbus torminalis* (L.) Crantz. *Phytochem.Lett.*, 4: 151-157.
- Pavlii, A.I. and G.V. Makarova (1970). Polyphenols of *Sorbus quercifolia*. *Chem. Nat. Comp.*, 6: 773.
- Pavlii, O.I., G.V. Makarova and Y.G. Borisyuk (1966). Flavonoids of *Sorbus pendula*. I. Flavonoids of the flowers of *Sorbus pendula*. *Farm. Zh. (Kiev)*, 21: 55-59.
- Perry, L.M. and J. Metzger (1980). *Medicinal Plants of East and South Asia*, The MIT Press, Cambridge England, pp. 341, 347
- Pyysalo, H. and T. Kuusi (1974). Phenolic compounds from the berries of mountain ash, *Sorbus aucuparia*. *J. Food Sci.*, 39: 636-638.
- Rizvi, T.S., L. Ali and F. Shaheen (2013). *Phytochemical investigations on Sorbus cashmiriana: isolation, structure elucidation, and biological evaluation of the chemical constituents*. LAP Lambert Academic Publishing, pp.84.
- Rolland, O. and J. Raynaud (1979). Phytochemical study of the buds on *Sorbus aria* (Rosaceae). *Bulletin des Travaux de la Societe de Pharmacie de Lyon*, 23: 3-6.
- Shavva, A.G., L.G. Matyukhina and I.A. Saltykova (1969). Ursolic acid in *Sorbus aucuparia*. *Chem. Nat. Comp.*, 5: 372.
- Sperens, U. (1997b). Fruit production in *Sorbus aucuparia* L. (Rosaceae) and pre-dispersal seed predation by the apple fruit moth (*Argyresthia conjugella* Zell.). *Oecologia*, 110: 368-373.
- Takaishi, K. and H. Kuwajima (1976). Prunasin and amygdalin from *Sorbus commixta*. *Phytochemistry*, 15: 1984.
- Termentzi, A., P. Alexiou, V.J. Demopoulos and E. Kokkalou (2008). The aldose reductase inhibitory capacity of *Sorbus domestica* fruit extracts depends on their phenolic content and may be useful for the control of diabetic complications. *Pharmazie*, 63: 693.
- Termentzi, A., P. Kefalas and E. Kokkalou (2006). Antioxidant activities of various extracts and fractions of *Sorbus domestica* fruit at different maturity stages. *Food Chem.*, 98: 599-608.
- Termentzi, A., M. Zervou and E. Kokkalou (2009). Isolation and structure elucidation of novel phenolic constituents from *Sorbus domestica* fruits. *Food Chem.*, 116: 371-381.
- Tsitsa-Tzardi, E., A. Loukis and S. Philianos (1991). Constituents of *Sorbus torminalis* fruits. *Fitoterapia*, 62: 282-283.
- Tsitsa-Tzardi, E., A. Loukis and S. Philianos (1992). Constituents of *Sorbus torminalis* leaves. *Fitoterapia*, 63: 189-90.
- Uddin, G., A. Latif, M. Arfan, M. Ali, S.H. Hussain, T.J. Simpson, R.J. Cox and M.I. Choudhary (2013). Phytochemicals from the stem wood of *Sorbus lanata* (D. Don.) Schauer. *Phytochem.Letts.*, 6: 84-89.
- Van, M.C.C., C.B. Almeida, R. Kort, M.N. Groot and T. Abee (2012). Germination inhibition of *Bacillus cereus* spores: impact of the lipophilic character of inhibiting compounds. *Int. J. Food Microbiol.*, 160: 124-130.
- Vogl, S., P. Picker, J. Mihaly-Bison, N. Fakhrudin, A.G. Atanasov, E.H. Heiss, C. Wawrosch, G. Reznicek, V.M. Dirsch, J. Saukel and B. Kopp (2013). Ethnopharmacological *in vitro* studies on Austria's folk medicine--an unexplored lore in vitro anti-inflammatory activities of 71 Austrian traditional herbal drugs. *J. Ethnopharmacol.*, 149: 750-751.

- Wagner, H., L. Hoerhammer, R. Dirscherl, G. Hitzler, L. Farkas and M. Nogradi (1968). Über die Synthese von Quercetin-3-glykosiden, IV. Synthese des Quercetin- und Kämpferol-3- β -gentiobiosids. *Chem. Ber.*, 101: 3419-3422.
- Wilhelm, H. (1963). On the occurrence of parasorbic acid in the fruits of the genus *Sorbus*. *Flora Oder Allg. Bot. Zeit.*, 153: 514-520.
- Yousuf, S., A. Latif, M. Arfan and M.I. Choudhary (2010). 1,2,4-Trimethoxy dibenzofuran-3-ol. *Acta Cryst. E*, 66: o3066.
- Yu, C., Y. Zheng, L. Li, G. Wang, Y. Li, T. Wu and H. Tang (2012). Isolation of chemical compounds from *Sorbus tianschanica* Rupr by high-speed counter-current chromatography. *J. Med. Plants Res.*, 6: 5142-5149.
- Zapesochaya, G.G., R.K. Aitbaeva and A.I. Ban'kovskii (1973). Flavonoids and chlorogenic acid of *Sorbus tianschanica* II. *Khim. Prir. Soedin.*, 9: 118.
- Zymone, Kristina, L. Raudone, R. Raudonis, M. Marksa, L. Ivanauskas and V. Janulis (2018). Phytochemical profiling of Fruit powders of twenty *Sorbus* L. cultivars. *Molecules*, 23: 2593.

(Accepted for publication September 2015)