

## FUNGICIDAL ACITIVITY OF *THUJA OCCIDENTALIS* L. LEAVES AGAINST ROOT ROT FUNGI ON CROP PLANTS

Asma Hanif and Shahnaz Dawar

Department of Botany, University of Karachi, Karachi-75270, Pakistan

Corresponding author: shahnaz\_dawar@yahoo.com

---

### ABSTRACT

In the present study, we investigated the effect of *T. occidentalis* L. leaves due to its fungicidal activity, therefore leaves extract as well as its whole powder form was taken to control *Fusarium* spp., *Macrophomina phaseolina* and *Rhizoctonia solani*. We found a significant increase in plant growth parameters when 1.0% w/v concentration of leaves extract was used as soil drenching, and seed treatment. Our data also indicated an increase in plant growth parameters when leaves powder was used as soil amendment at 0.5% w/w. We also found a measurable reduction in the colonization of root rot fungi. Although soil amended with 2.5% w/w leaves powder showed complete suppression in the colonization of pathogenic fungi, but failed to improve plant growth seems to be toxic to test plants as it ceased the germination and caused stunted and wilting symptoms were observed.

**Key-words:** *Thuja occidentalis*, root rot fungi, plant growth

---

### INTRODUCTION

Plant pathogenic fungi, *Rhizoctonia solani*, *Macrophomina phaseolina* and *Fusarium* spp. have a broad host range and worldwide distribution (Agrios, 2005). For many years, synthetic fungicides have been used as a quick and effective management strategy for crop protection from pathogenic fungi infestation (Osman and Al-Rehiayam, 2003). However, due to the extensive use of fungicides, there have been negative effects on ecosystems and human health including increased risk of human cancer (Stranger and Scott, 2005). In Western countries, a large number of agrochemicals are banned because of their undesirable attributes such as; extensive degradation period, accumulation in the food chain, high toxicity (Gatto *et al.*, 2011). In addition, most of the available agrochemicals can also destroy beneficial organisms (Gatto *et al.*, 2011). Thus there is a need to search of alternative control strategies to minimize the reliance on the synthetic fungicides. This can be achieved by exploring crop plants that may contain a rich source of antimicrobial agents (Mahesh and Satish, 2008).

Extracts from a number of plants have been reported to have fungicidal activity against many plant pathogenic fungi. The natural plant products possess anti-microbial activity, biodegradability as well as non- phytotoxicity (Gómez *et al.*, 1990; Talibi *et al.*, 2012), produce a great deal of secondary metabolites which exhibit antifungal activity (Bennett and Wallsgrove, 1994; Osbourne, 1996) and has been adequately explored against phytopathogenic fungi (Gangadevi *et al.*, 2008). Currently, plant extracts has received attention because of their phytochemicals (Jasuja *et al.*, 2012) which exhibit great potential as microbial inhibitors are naturally toxic to fungi and bacteria (Basile *et al.*, 1999). Use of medicinal plants having antifungal properties could be used against plant pathogens in an organic farming system because of their environment friendly mode of disease management (Srivastava *et al.*, 1997; Shanmugavalli *et al.*, 2009). For instance, *Thuja occidentalis* L., belonging to family Cupressaceae is characteristically coniferous and monoecious tree measuring from 12-21 meters in height (Naser *et al.*, 2005). The tree has given numerous names, including white cedar of the east, arbor vitae, white cedar of the north, white cedar, tree of life, swamp cedar and yellow cedar (Tsiri *et al.*, 2009). *T. occidentalis* leaves have oil and resin which gives the characteristic of intense odor, a sharp taste, including camphoric and balsamic (Farmacopeia, 2011).

Fresh plant contains reducing sugar (2.07%), water soluble minerals (2.11%), essential oil (0.6%), free acid (1.67%), tannic agents (1.31%) and water-soluble polysaccharides (4.9%) (Harnischfeger and Stolze, 1983). The essential oil of fresh leaves contains isothujone (8%), thujone (65%), fenchone (8%), sabinene (5%) and  $\alpha$ -pinene (2%) as the main monoterpenes (Witte *et al.*, 1983). Thujones compounds are known as insect repellent which was confirmed by Hwang *et al.*, (1985) which can cause high mortality in root worm larvae in western corn. Thujones also exhibit antimicrobial activity against gram positive bacteria, gram negative bacteria and many fungal strains (Tsiri *et al.*, 2009). In addition, flavonoids compound is also present which are known to be synthesized by plants in response to microbial infection (Cowan, 2002). Flavonoid contains amentoflavone that exhibits strong antifungal activity against numerous pathogenic fungi (Jung *et al.*, 2006) and antiviral activity against respiratory syncytial

virus (Ma *et al.*, 2001). *T. occidentalis* leaves exhibits anticancer, antidiabetic, antitumor, anti-ulcerative, antioxidant, hepatoprotective and hypolipidemic activities (Dubey and Batra, 2009; Madhuri and Pandey, 2009).

In the present study, we investigated the fungicidal potential of *T. occidentalis* leaves in the management of root rot fungi and its effect on the growth of crop plants.

## MATERIALS AND METHODS

**Collection of plants:** Fresh *Thuja occidentalis* L. leaves were collected from trees growing in Karachi University (KU) campus. After collection, the collected leaves were dried under shade at room temperature. Then leaves were grinded into powder form using an electric grinder machine. The seeds powder thus obtained were stored in glass jars for further studies.

**Preparation of aqueous extract:** For the aqueous extraction, *T. occidentalis* leaves @ 0.5, 1.0, 1.5, 2.0 and 2.5 g were separately soaked in 100 mL sterilized distilled water, respectively, for 48 hours at room temperature (28-34°C). The extract was separately filtered through two layers of Whatman No. 1 filter paper in 250mL Pyrex flask. Filtrate stored at 4°C for further study.

**Soil properties:** Soil used for this study was collected from an experimental plot in the Botany Department (Karachi University). The sandy loam soil contained; sand 73%, silt 12%, clay, 15%, pH 7.3 with moisture holding capacity (MHC) of 42% (Keen and Raczowski, 1922). Natural soil infestation contains 6-8 sclerotia/g soil of *M. phaseolina* found by wet sieving technique (Sheikh and Ghaffar, 1975), 15-20% of *R. solani* on sorghum seeds used as baits (Wilhelm, 1955) and 2300 CFU/g soil of *Fusarium* spp., was assessed by soil dilution technique (Nash and Synder, 1962).

### Pot experiment:

**Soil amendment:** *T. occidentalis* leaves powder was amended in the 300g soil at 0.5, 1.0, 1.5, 2.0 and 2.5% w/w respectively. Soil without leaves powder (control) was used for comparison. Pots containing soil were watered daily to allow the decomposition of organic substrate. After one week, four seeds of mung bean (*Vigna radiata* (L.) R. Wilczek. cv. NM-2006), mash bean (*Vigna mungo* (L.) Hepper cv. NM-97), okra (*Abelmoschus esculentus* (L.) Moench cv. Arka anamika) and sunflower (*Helianthus annuus* L. cv. Hysun-38) were planted in each pot containing various amount of leaves powder.

**Seed treatment:** Test seeds after surface sterilization at 1% Ca (OCl)<sub>2</sub>, dried aseptically under laminar flow hood. Surface disinfested seeds were separately treated with different concentrations of *T. occidentalis* aqueous leaves extracts for 5-10 minutes, whereas seeds without any treatments served as control. Four treated seeds of test crops with different concentrations along with the control were separately planted in each pot.

**Soil drenching:** Twenty milliliter of aqueous *T. occidentalis*, leaves extracts at 0.5, 1.0, 1.5, 2.0, 2.5% w/concentrations, respectively, were separately drenched in each plastic pot containing 300g soil. Then 4 seeds were planted in each pot. Soil drenched with 20 mL sterilized distilled water served as control.

Pots were placed in completely randomized design (CRD) on a greenhouse bench in three replicates per treatment under the natural sunlight. All plants were watered regularly in order to maintain sufficient moisture content required for the growth of test plants. Growth parameters including shoot weight and height, root weight and length were recorded after one month of seed germination.

**Determinations colonization of root rot fungi:** Plant roots were washed in running tap water, after washing roots were surface sterilized with 1% Ca (OCl)<sub>2</sub> for 2-3 minutes, cut into one cm and dried on blotter paper under laminar flow hood. Transferred root pieces (5 root pieces/plate) on Petri plates containing Potato dextrose agar medium supplemented with antibiotics to inhibit the growth of bacteria. Petri plates were incubated at room temperature (27-33°C) for a week and the colonization of roots by root rot fungi was recorded.

**Statistical analysis:** Data were subjected to analysis of variance (ANOVA) followed by the least significant difference (LSD) test as P = 0.05 and Duncan's multiple range test to compare treatment means, using Statistica software according to Sokal and Rohlf (1995).

## RESULTS

We found that seeds of mash bean and mung bean treated with *T. occidentalis* leaves extracts @ 2.0 and 2.5% w/v showed a higher growth as compared to other concentrations. Soil amendment and soil drenched with *T. occidentalis* leaves @ 0.5 and 1.0% concentrations effectively increased both the plant height and the weight over 2.5% in both methods. Colonization of root rot pathogens were significantly ( $P < 0.01$ ) reduced at all concentrations (Fig. 1)




Fig. 1. Effect of seed treatment, soil amendment and soil drenching with *T. occidentalis* leaves on growth parameters of crop plants.

When seed treatment method used, we found a complete inhibition of root rot colonization in both mash bean and mung bean crops. When the soil was drenched at 2.5% w/v ( $P < 0.001$ ) with *T. occidentalis* aqueous leaves extracts, we also found a complete suppression of *R. solani* colonization in mash bean plants that were planted in soil amended with 2.5% w/w leaves powder. Both seed treatments and soil drenched with *T. occidentalis* leaves extracts used @ 0.5, 1.0 and 1.5% w/v concentrations ( $P < 0.001$ ) enhanced the growth parameters in both sunflower and okra plants (Fig. 2).




Fig 2. Effect of seed treatment, soil amendment and soil drenching with *T. occidentalis* leaves on growth parameters of crop plants.






Fig 3. Effect of seed treatment, soil amendment and soil drenching with *T. occidentalis* leaves on root rot fungi of crop plants.

The highest growth was noticed, when the soil was amended with 0.5% w/w leaves powder as compared to 2.0 and 2.5% w/w. In all three methods, 2.5% leaves extracts ( $P < 0.05$ ), did not improve the plant growth, but rather completely suppressed root rot colonization (Fig. 3). Greater inhibition of *R. solani*, *M. phaseolina* and *Fusarium* spp. colonization ( $P < 0.01$ ) was recorded in seed treatment, soil drenching and soil amendment methods, when *T. occidentalis* leaves used @ 1.0%, followed by 0.5% concentration which not only increased the plant weight and height but also recorded maximum control of pathogenic fungi. When *T. occidentalis* leaves powder amended @ 2.5% w/w in soil, it completely inhibited the *R. solani* colonization observed in sunflower plants, but in case of okra it suppressed both *R. solani* and *M. phaseolina* colonization (Fig. 4).




Fig 4. Effect of seed treatment, soil amendment and soil drenching with *T. occidentalis* leaves on root rot fungi of crop plants.

## DISCUSSION

Application of *Thuja occidentalis* leaves used @ 1.0% in seed treatment, soil drenching and soil amendment not only enhanced the plant weight and height but also showed maximum inhibition in the colonization of root rot fungi in both leguminous and non-leguminous crops. Ikram and Dawar (2012) also found a similar result. They used

*Aerva javanica* leaves powder at 1% w/w mixed with soil and noticed significant inhibition of *R.solani*, *M. phaseolina* and *Fusarium* spp. colonization.

Organic amendments not only improve crop productivity, but also suppress soil borne diseases caused by plant pathogenic fungi and nematodes (Stone *et al.*, 2003; Alam, 1990; Mital and Gowswami, 2001). Addition of organic amendments such as; animal manure, green manure, sawdust, compost, wood chips, grass clippings, sphagnum peat, wood ash and many others are commonly used in farmlands conditions suitable for improving plant growth (Hadar *et al.*, 1992; Muchovej and Pacovsky, 1997). Tariq *et al.* (2006) and (2008) reported that when *Avicennia marina* and *Rhizophora mucronata* applied to the soil as organic amendment at different ratios not only increased the plant weight and height but also controlled plant parasitic nematode and root rot fungi infestation on different crops. Application of soil amendment with plant powder inhibits plant pathogens by releasing toxic substances like phenols (Ali *et al.*, 2001; Shaukat *et al.*, 2001). Seed treatment inhibit the fungi residing either on the surface of the seed or maybe inside the seed but also protect from the pathogen that are inhabited in the soil causes seed rot, damping off and root rot diseases (Martha *et al.*, 2003).

In previous studies, okra and mash bean seeds treated with aqueous extracts of *Sida pakistanica* and *Senna holosericea* @ 25, 50 and 100% w/v concentrations have shown reduced colonization of root rot fungi (Emmanuel *et al.*, 2010). In addition, *In vitro* study, the alcoholic extracts of *T. occidentalis* twigs have not only shown antibacterial activity against *Citrobacter*, *Escherichia coli*, *Staphylococcus aureus*, *Shigella flexneri*, *Yersinia aldovae* and *Pseudomonas aeruginosa* but have also shown antifungal activity against *Trichophyton rubrum*, *Macrophomina* spp., *Fusarium solani*, *Saccharomyces cerevisiae*, *Candida albicans* and *Aspergillus parasiticus* (Jahan *et al.*, 2010). Both aqueous and alcoholic extracts of *T. occidentalis* twigs exhibited antioxidants and anti-inflammatory activity (Dubey and Batra, 2009). *T. occidentalis* leaves contain carotenoid compounds as a secondary metabolite (Krinsky 1979). Carotenoids are an effective inhibitor of various diseases such as heart and cancer disease (Kotikova *et al.*, 2009). Chaudhary *et al.* (2015) reported that carotenoid present in *T. occidentalis* leaves have potent antibacterial activity against *Bacillus subtilis*, *Bacillus megaterium*, *Bacillus amyloliqui faciens*, *Proteus vulgaris* and *Salmonella typhi*. Aqueous extract of leaves, bark, fruits and stem of *Eucalyptus* sp., used @ 5% w/v were found to inhibit *M. phaseolina*, *R. solani* and *Fusarium* sp. Infestation on mung bean and chick pea (Dawar *et al.*, 2007). Soil drenching has been shown as an effective method of controlling root rot diseases reported by Abdel-Kader *et al.*, (2012) studied with different bio-agents such as; *Trichoderma harzianum*, *Trichoderma viride*, *Bacillus subtilis*, *Pseudomonas flourescens* and *Sacchchromyces serivisae* and drenched into the soil, showed their efficacy in reducing root rot incidence of post-emergence growth stage of Cucumber, Cantaloupe, Tomato and Pepper.

We suggest that low concentration of soil drenching and soil amended with *T. occidentalis* leaves powder may decrease the colonization of fungal pathogens inhabiting in the soil. However, seed treatment is the best method either used at low or high concentration protects seeds from external attack of the pathogens considered as biological control and eco-friendly. We found *T. occidentalis* leaves can be best used as a fungicide against root rot fungi and may also improve plant growth.

## ACKNOWLEDGEMENT

We are grateful to Dean Faculty of Science for providing financial support in order to carry out this research work.

## REFERENCES

- Abdel-Kader, M.M., N.S. EL-Mougy and M.D.A. Lashin (2012). Different approaches of bio-control agents for controlling root rot incidence of some vegetables under greenhouse conditions. *International Journal of Agriculture and Forestry*, 2(1): 115-127.
- Agrios, G.N. (2005). *Plant pathology*, 6<sup>th</sup> edition. Academic Press, New York.
- Alam, M. (1990). Neem in nematode control. In: *Nematode biocontrol (Aspects and prospects)*: M.S. Jairajpuri, M.M. Alam and I. Ahmed. BS Publishers and Distributors (Pvt) Ltd. Dehli-110032, India.pp.51-55.
- Ali, N.I., I.A. Siddiqui, M.J. Zaki and S.S. Shaukat (2001). Nematicidal potential of *Lantana camara* against *Meloidogyne javanica* in mung bean. *Nematologica Mediterranea*, 29: 99-102.
- Bennett, R.N. and R.M. Wallsgrove (1994). Secondary metabolites in plant defense mechanisms. *New Phytology*, 127: 617-633.
- Basile, A., S. Giordano, J.A. López-Sáez and R.C. Cobiánchi (1999). Antibacterial activity of pure flavonoids isolated from mosses. *Phytochemistry*, 52: 1479-1482.
- Chaudhary, P., G. Bhagwati and M. Kavita (2015). Carotenoid and antibacterial analysis of *Thuja occidentalis*. *Indian journal of applied research*, 5(7): 112-114.

- Cowan, M.M. (2002). Plant products as antimicrobial agents. *Clin. Microbiol. Rev.*, 12: 564-582.
- Dawar, S., S. Younus., M. Tariq and M.J. Zaki (2007). Use of Eucalyptus sp., in the control of root infecting fungi on mung bean and chick pea. *Pak. J. Bot.*, 39 (3): 975-979.
- Dubey, S.K. and A. Batra (2009). Antioxidant activity of *Thuja occidentalis* linn. *Asian Journal of Pharmaceutical and Clinical Research*, 2: 73-76.
- Emmanuel, A., S. Dawar and M. J. Zaki (2010). Effect of *Sida Pakistanica* S. Abedin and *Senna holosericea* Fresen on growth and root rot diseases of okra and mash bean. *Pak. J. Bot.*, 42(1): 391-400.
- Farmacopeia (2011). Homeopática Brasileira: *Thuya occidentalis*. Fundação Oswaldo Cruz, third edition.
- Gangadevi, V., S. Yogeswari, S. Kamalraj, G. Rani and J. Muthumary (2008). The antibacterial activity of *Acalypha indica* L. *Indian J. Sci. Technol.*, 1(6): 1-5.
- Gatto, M.A., A. Ippolito, V. Linsalata, N.A. Cascarano, F. Nigro, S. Vanadia and D. Di-Venere (2011). Activity of extracts from wild edible herbs against postharvest fungal diseases of fruit and vegetables. *Postharvest Biol. Technol.*, 261: 72-82.
- Gómez, G.R., R. Chilpa, L. Quijano, J.S. Calderón Pardo and T. Ríos Castillo (1990). Methoxifurans auranols with fungostatic activity from *Lonchocarpus castilloi*. *Phytochemistry*, 29: 459-463.
- Hadar, Y., R. Mandelbaum and B. Gorodecki (1992). Biological control of soilborne plant pathogens by suppressive compost. In: *Biological control of plant diseases*. Tjamos, E.C., G.C. Papavizas and R.J. Cook (Eds.). Plenum Press, New York. pp. 79-83.
- Harnischfeger, G. and H. Stolze (1983). Bewahrte Pflanzendrogen in Western Medicine. Notamed Verlag, Bad Homburg/Melsungen. pp. 250-915.
- Hwang, Y.S., K.H. Wu, J. Kumamoto, H. Axelrod and M.S. Mulla (1985). Isolation and identification of mosquito repellents in *Artemisia vulgaris*. *Journal of Chemical Ecology*, 11: 1297-1306.
- Ikram, N. and S. Dawar (2012). Soil amendment with *Aerva javanica* (Burm. f.) Juss. ex Schult. in the control of root rot fungi of cow pea (*Vigna unguiculata* (L.) Walp.) and mung bean [*Vigna radiata* (L.)]. *Acta Agrobotanica*, 65(3): 69-74.
- Jahan, N., M. Ahmad, M. Mehjabeen, S. Zia-ul-haq, M. Alam and M. Qureshi (2010). Antimicrobial screening of some medicinal plants of Pakistan. *Pak. J. Bot.*, 42(6): 4281-4284.
- Jasuja, N.D., R. Saxena, S. Chandra and R. Sharma (2012). Pharmacological characterization and beneficial uses of *Punica granatum*. *Asian J. Plant Sci.*, 11: 251-267.
- Jung, H.J., W.S. Sung, S. Yeo, H.S. Kim, I. Lee, E. Woo and D.G. Lee (2006). Antifungal effect of amentoflavone derived from *Selaginella tamariscina*. *Archives of Pharmacol. Research*, 29 (9): 746-751.
- Keen, B.A. and H. Raczkowski (1922). Clay contents and certain physical properties of soil. *J. Agric. Sci.*, 11: 441-449.
- Kotikova, Z., A. Hejtmankova and J. Lachman (2009). Determination of the influence of variety and level of maturity on the content and development of carotenoids in tomatoes. *Czech J. Food Sci.*, 27: 200 - 204.
- Krinsky, N.I. (1979). Carotenoid protection against oxidation. *Pure Appl. Chem.*, 51: 649-660.
- Ma, S., P.P. But, V.E. Ooi, Y. He, S.H. Lee, S. Lee and R. Linn (2001). Antiviral amentoflavone from *Selaginella sinensis*. *Biological & Pharmaceutical Bulletin*, 24 (3): 311-312.
- Mahesh, B. and S. Satish (2008). Antimicrobial activity of some important medicinal plant against plant and human pathogens. *World J. Agri. Sci.*, 4: 839-843.
- Madhuri, S. and G. Pandey (2009). Some anticancer medicinal plants of foreign origin. *Current Science*, 96 (6): 779-783.
- Martha, M., J. Riesselman, D. Mathre, B. Jhonston and S. Blodgett (2003). *Manual of small seed grain treatment guide*, pp. 55.
- Mital, A and B. K. Gowswami (2001). Role of undi oil seed cake for the management of disease complex caused by *Fusarium solani* and *Meloidogyne incognita* on brinjal. *Pak. J. Nematol.*, 19(1&2): 87-90.
- Muchovej, R.M.C. and R.S. Pacovsky (1997). Future directions of by-products and wastes. In: *Agricultural uses of by-products and wastes*. Rechcigl, J.E. and H.C. MacKinnon (Eds.). ACS Symposium Series, American Chemical Society Washington, D.C. pp. 1-19.
- Nash, S.M. and W.C. Synder (1962). Quantitative estimations by plate counts of propagules of the bean root rot, *Fusarium* in field soils. *Phytopathology*, 52: 567-572.
- Naser, B., C. Bodinet, M. Tegtmeier and U. Lindequist (2005). *Thuja occidentalis* (Arbor vitae): A Review of its Pharmaceutical, Pharmacological and Clinical Properties. *Evid Based Complement Alternat Med.*, 2(1): 69-78.
- Osbourne, A.E. (1996). Preformed antimicrobial compounds and plant defense against fungal attack. *The Plant Cell*, 8: 1821-1831.



- Osman, K.A. and S. Al-Rehiayam (2003). Risk assessment of pesticide to human and the environment. *Saudi J. Biol. Sci.*, 10:81-106.
- Shaukat, S.S., I.A. Siddiqui, N.I. Ali and M.J. Zaki (2001). Biological and chemical control of root infecting fungi and their effect on growth of mung bean. *Pak. J. Biol. Sci.*, 4: 1240-1242.
- Shanmugavalli, N., V. Umashankar and Raheem (2009). Antimicrobial activity of *Vanilla planifolia*. *Indian J. Sci. Technol.*, 2(3): 37-40.
- Sheikh, A.H and A. Ghaffar (1975). Population study of Sclerotia of *Macrophomina phaseolina* in cotton field. *Pak. J. Bot.*, 7: 13-17.
- Sokal, R. R. and F.J. Rohlf (1995). *Biometry: The principles and practices of statistics in biological research*. Freeman, New York. Pp.887.
- Srivastava, A.K., B. Bihari and B. Lal (1997). Studies on biofungicidal properties of leaf extract of some plants. *Indian Phytopathol.*, 50(3): 408-411.
- Stranger, R.N. and P.R. Scott (2005). Plant Diseases: a threat to global food security. *Annual Review of phytopathology*, 43: 83-116.
- Stone, A.G., G.E. Vallad, L.R. Cooperband, D.R. Rotenburg, H.M. Darby, W.R. Stevenson and R.M. Goodman (2003). Impact of annual organic amendment on disease incidence in a three year vegetable rotation. *Plant Disease*, 87: 1037-1042.
- Talibi, I., L. Askarne, H. Boubaker, E.H. Boudyach, F. Msanda, B. Saadi and A. A. B. Aoumar (2012). Antifungal activity of Moroccan medicinal plants against citrus sour rot agent *Geotrichum candidum*. *Lett. Appl. Microbiol.*, 55:155–161.
- Tariq, M., S. Dawar, F.S. Mehdi and M.J. Zaki (2006). Use of *Avicennia marina* in the control of root infecting fungi on okra and mash bean. *Pak. J. Bot.*, 38(3): 811-815.
- Tariq, M., S. Dawar, F.S. Mehdi and M.J. Zaki (2008). The effect of mangroves amendments to soil on root rot and root knot of potato (*Solanum tuberosum* L.). *Acta Agrobotanica*, 61(1): 115-121.
- Tsiri, D., K. Graikou, L. Pobłocka-Olech, M. Krauze-Baranowska, C. Spyropoulos and I. Chinou (2009). Chemosystematic value of the essential oil composition of *Thuja* species cultivated in Poland-antimicrobial activity. *Molecules*, 14 (11): 4707-4715.
- Wilhelm, S. (1955). Longevity of the *Verticillium* with fungus in the laboratory and field. *Phytopathology*, 45: 180-181.
- Witte, I., J. Berlin, V. Wray, W. Schubert, W. Kohl, G. Hofle and J. Hammer (1983). Mono- und diterpenes from cell cultures of *Thuja occidentalis*. *Planta Med.*, 49: 216–221.

(Accepted for publication March 2017)






Fig 1. Effect of seed treatment, soil amendment and soil drenching with *T. occidentalis* leaves on growth parameters of crop plants.




Fig 2. Effect of seed treatment, soil amendment and soil drenching with *T. occidentalis* leaves on growth parameters of crop plants.




Fig 3. Effect of seed treatment, soil amendment and soil drenching with *T. occidentalis* leaves on root rot fungi of crop plants.