

ASSESSMENT OF GENETIC DIVERSITY IN EXOTIC GERMPLASM OF WHEAT (*TRITICUM AESTIVUM* L.) BY USING PCA AND CLUSTER ANALYSIS

Shazia Parveen¹, Tasveer Zahra Bokhari¹, Muhammad Faheem Siddiqui³, Manzoor Hussain² and Faheem Iqbal²

¹Institute of Pure and Applied Biology, Bahauddin Zakariya University, Multan, Pakistan.

²Regional Agricultural Research Institute, Bahawalpur, Pakistan.

³Department of Botany, University of Karachi, Karachi 75270, Pakistan.

ABSTRACT

The objective of study was to assess genetic diversity in exotic germplasm (8th EBWYT) of wheat (*Triticum aestivum* L.) based on cluster analysis and principle component analysis (PCA). There was a significant relationship among the studied traits i.e. days to heading, plant height, spike length, canopy temperature and grain yield per plant. A positive significant correlation was observed between spike length and number of spikelet per spike while a negative significant correlation was assessed between canopy temperature and thousand kernel weight. Out of seven, three PCA's axes inhibited more than one Eigen value but the level of dissimilarity was high which indicated that the germplasm was with broad genetic base. Cluster analysis expressed high level of diversity because two main groups were observed, one with 14 and the other with 16 genotypes.

Keywords: Exotic germplasm, Genetic diversity, PCA, Cluster analysis, *Triticum aestivum*.

INTRODUCTION

Wheat (*Triticum aestivum* L.) is the most important food crop of the world which is being cultivated on 217.2 Million hectare area with 651 million tons production and average yield of 2906 kg / ha (Anonymous, 2010). With regard to its production and utilization it is one of the most important sources of sustenance for mankind since its domestication from 15000 to 10000 BC. Wheat occupies a primary position in daily food consumption because it is the cheapest source of calories and protein (Maqbool *et al.*, 2010). The contribution of wheat among total cereal intake is 84% in Pakistan. Genetic diversity of plants enhances their potential efficiency for breeding which may result in enhanced food production (Mostafa *et al.*, 2011). The necessity of performing breeding experiment for the production of resistant plant varieties under different condition showed prolonged food production (Martin *et al.*, 2008). The development of new varieties with desirable genetic makeup takes great efforts by breeders to overcome the consumption pressure of ever increasing population by improving the yield potential of wheat (Memon *et al.*, 2007). To reconstruct the ideotypes of plant, available genetic resources are utilizing by wheat breeders to meet the ever increasing requirements of the population (Memon *et al.*, 2007). A measure of the genetic relationship between parents and progeny is determined by heritability value (Memon *et al.*, 2007). A lot of research work has been done to insert the desirable genes in present wheat varieties to enhance the crop productivity. The insertion of semi dwarfing genes Rht1 Rht2 are associated with development of high yielding varieties with grain yield associated traits (Sial *et al.*, 2010). Ajmal *et al.*, (2009) estimated genetic parameters and characterization associated in wheat while genetic heritability for grain yield and its related characters in spring wheat (*Triticum aestivum* L.) was studied by Shabana *et al.*, (2007). Eivazi *et al.*, (2007) suggested that identification of genetic diversity analyzed with the help of some appropriate methods i.e. PCA, factor analysis and cluster analysis. Shah *et al.*, (1999) performed genetic analysis of agronomic traits controlled by wheat chromosome 3A.

The present research was designed to sort out better performing genotypes of wheat against WG-99 (a race of stem rust fungus), under local conditions (Bahawalpur-Pakistan) and to observe diversity in valuable CIMMYT germplasm.

MATERIALS AND METHODS

A CIMMYT nursery of "8th Elite bread wheat yield trail" was selected for present study. This valuable germplasm is resistant to stem rust race i.e. WG-99. This trail was consisted of 30 wheat genotypes, sown at Regional Agricultural Research Institute (RARI), Bahawalpur- Pakistan, for performance under local environment and genetic diversity in the germplasm. Plot size was kept as 5m in length and 1.8 m in width with 6 rows. Distance between rows was 30cm. Randomized Complete Block Design (RCBD) was used with three replicates. Sowing was

done on mid November 2013-2014 cropping season. The experiment was conducted by using local recorded management practices.

During the growing season, following traits were measured by random selection and tagging 10 plants from each experimental unit. Plant height of the main tiller of each selected plant was measured in centimeters from the ground level to tip of spike excluding awn. At maturity the spike length of main tillers of nominated plant measured in centimeter, from the base to the tip of the spike excluding awn. Number of spikelet per spike, days to heading, 1000 grain weight, CTD (C), grain yield per plot (kg/ha) were measured.

Analysis of variance was done by using statistical formula given by Steel and Torrie (1980). Other statistical techniques like Principal Component Analysis (PCA) and Cluster Analysis were also used in the present study.

RESULTS AND DISCUSSION

Elite bread wheat yield trial (8th EBWYT) was planted along with 3 reps and 30 entries in each. Mean grain yield was recorded about 4080 kg/hectare with a maximum of 4720.7 kg/hectare. Mean Thousand kernel weight was 36.7 grams with a maximum of 40.33 grams (Table 1).

Table 1. Descriptive statistics of studied traits in 30 genotypes of the 8th EBWYT.

Sr. No.	Description	CT	DH	GY	NS	PLH	SL	TKW
1	Mean	25.575	93.701	4080	9.4893	99.823	11.622	36.7
2	SD	1.2551	2.4525	316.55	0.8566	8.5743	0.7086	2.1705
3	Variance	1.5752	6.0149	100202	0.7338	73.519	0.5021	4.7109
4	SE Mean	0.2291	0.4478	57.793	0.1564	1.5654	0.1294	0.3963
5	CV	4.9075	2.6174	7.7584	9.0269	8.5895	6.0971	5.9141
6	Minimum	22.73	90.33	3563	8.33	86.67	10.17	32.67
7	Maximum	27.27	99.67	4720.7	13	131	12.5	40.33

Key to abbreviations: CT: Canopy Temperature °C, DH: Days to heading, GY: Grain Yield (Kg/Hectare), NS: Number of Spikelet per spike, PLH: Plant Height (cm), SL: Spike length (cm), TKW: Thousand kernel weight (g), SD = standard deviation, SE = standard error and CV = cumulative variance.

Analysis of variance of all the studied traits remained non-significant among the replicates (Table 2). By taking genotypes as factor of variability, all the studied traits found with significant variation (Table 3). This indicates possible diversity in genotypes.

Table 2. Analysis of Variance (ANOVA) of agronomic traits showing non significance among replicates.

SOV	Canopy Temperature				Days to Heading		
	DF	MS	F value	P	MS	F value	P
Reps.	2	5.22	1.75	0.1792	3.03	0.43	0.6497
Error	87	2.97			6.99		
Reps.	Grain Yield (Kg/hectare)				Number of Spikelet per spike		
	DF	MS	F value	P	MS	F value	P
Reps.	2	21014	0.14	0.8707	0.544	0.33	0.722
Error	87	151510			1.6712		
Reps.	Plant Height (cm)				Spike length (cm)		
	DF	MS	F value	P	MS	F value	P
Reps.	2	126.544	0.8	0.45	1.17	1.18	0.312
Error	87	157.219			0.99		
Reps.	Thousand kernel weight (g)						
	DF	MS	F value	P			
Reps.	2	1.3	0.22	0.8			
Error	87	5.911					

Key to abbreviations: Reps. = Replicates, dF = degree of freedom, MS = mean square, P = Probability

Correlation coefficient assessed significant between spike length and number of spikelet per spike. Similar result was reported by Ilker *et al.* (2010). Thousand kernel weights were found negatively significant with canopy

temperature indicating that high canopy temperature can shrivel the seed up to ample decrease in weight. Highly significant positive correlation was found between grain yield and thousand kernel weight (Table 4). Akram *et al.*, (2008) reported that grain yield was positively correlated with thousand grain weight. Similar results were also reported by Dokuyueu and Akkaya (1999). Yucel *et al.*, (2009) shows a positive and significant correlation between spikelet number per spike, grain yield per spike, grain number per spike and spike length.

Table 3. Analysis of Variance (ANOVA) of agronomic traits among 30 genotypes.

SOV	Canopy Temperature				Days to Heading		
	DF	MS	F	P	MS	F	P
Genotypes	29	4.72	2.14	0.006648	18	11.8	0
Error	60	2.21			1.5		
	Grain Yield (Kg/hector)				Number of Spikelet per spike		
	DF	MS	F	P	MS	F	P
Genotypes	29	300604.5	4	0.000003	2.201	1.597	0.063375
Error	60	75097.44			1.378		
	Plant Height (cm)				Spike length (cm)		
	DF	MS	F	P	MS	F	P
Genotypes	29	220.6	1.756	0.033201	1.51	2.02	0.010822
Error	60	125.6			0.75		
	Thousand kernel weight (g)						
	DF	MS	F	P			
Genotypes	29	14.1	7.96	0			
Error	60	1.8					

Key to abbreviations: See Table 2

Table 4. Pearson's correlation coefficient (r) among studied traits.

Variables	DH	PLH	SL	NS	TKW	CT
PLH	0.171					
SL	-0.018	-0.002				
NS	-0.072	0.109	0.661			
TKW	0.307	0.034	0.148	0.063		
CT	-0.189	-0.177	0.071	-0.062	-0.441	
GY	0.126	0.180	-0.141	-0.090	0.525	-0.903

Values in bold are different from 0 with a significance level $\alpha=0.05$;
Key to abbreviations: See Table 1

Principal Component Analysis, a technique to show transformation of large number of variables into correlated small number of components was carried out. Factors (F₁ and F₂) explained the variability of about 58.83% with the Eigen value 2.413 and 1.705 respectively. Seven factors explained 100 percent variability of the whole parameters estimated (Table 5).

Table 5. Principal Component Analysis.

	F1	F2	F3	F4	F5	F6	F7
Eigen value	2.413	1.705	1.039	0.957	0.511	0.306	0.069
Variability (%)	34.470	24.363	14.842	13.670	7.301	4.373	0.980
Cumulative %	34.470	58.833	73.675	87.346	94.647	99.020	100.000

Scree plot (Fig. 1) among Eigen values, Factors and cumulative variability indicating effective factors showing maximum variability. In Fig. 2, spike length and number of spikelet per spike has been shown with same pattern of variation explained by F_2 . Plant height, thousand kernel weight, days to heading and yield is briefly explained with respect to their variation with F_1 . Canopy temperature is negative to all of the components.




Fig. 1. Scree plot indicating factors distribution along with Eigen values. Cumulative variability (Red line) indicating F_1 and F_2 as the most effective components of variability.




Fig. 2. Percent variability (58.83) explained by F_1 (X-axis) and F_2 (Y-axis).

Table 6, showed percent contribution of each studied trait toward the appropriate factor. Few genotypes such as 21, 30, 16 and 29 showed high diversity as compared to the others (Fig.3).

Table 6. Contribution of the variables (%).

	F1	F2	F3	F4	F5	F6	F7
DH	6.442	0.008	44.428	30.530	17.608	0.011	0.974
PLH	3.560	0.870	44.350	39.967	10.767	0.279	0.207
SL	0.262	48.297	0.226	1.610	0.309	49.198	0.098
NS	0.021	47.784	0.181	2.256	7.160	40.125	2.473
TKW	20.755	2.426	0.483	18.928	48.719	7.048	1.641
CT	33.762	0.009	4.308	3.640	14.674	0.298	43.309
GY	35.198	0.606	6.024	3.070	0.763	3.040	51.299

Key to abbreviations: See Table 1




Fig. 3. Biplot of genotypes and the studied traits, showing the variation explained by F1 and F2.

In Fig. 4, structural equation of multiple regression model is explained. Grain yield as a dependent Y-variable, along with error variance of 12798.02 depicts relationship with all other studied traits taken as X-variables.

As long as diversity among the genotypes is concerned, a profile shows the pattern of variation (Fig. 5) and the distance among similarities of 30 genotypes is shown in Fig. 6. Cluster analysis also expresses high level of diversity because two main groups have been observed (Fig. 6.) i.e. the one with 14 and the other with 16 genotypes.

Histogram of the data of 30 genotypes shows that thousand kernel weights above 40 is concerned with only three genotypes (Fig. 7). Grain yield is not normally distributed as in Fig. 8, hence above 4600 kg ha⁻¹ is found in two genotypes only.




Fig. 4. Structural equation of multiple regression model showing Grain Yield (GY) as Y-variable being influenced by multiple X variables (DH, PLH, SL, NS, TKW, CT).

Note: $GY = -15.034 \cdot DH + 2.357 \cdot PLH - 0.399 \cdot SL - 62.724 \cdot NS + 29.152 \cdot TKW - 210.964 \cdot CT$ Error var. = 12798.022, $R^2 = 0.872$




Fig. 5. Depending upon the studied traits a diversity profile of 30 genotypes made by using a software PAST






Fig. 6. Cluster Analysis showing Genetic Diversity among 30 Genotypes



The Least Significant Difference (LSD) showing significant groups among genotypes as genotype number 02, 10 and 30 is depicted in (Table 7) and also for thousand kernel weight genotype 14, 29 and 30 expressed significant.

Table 7. Least significant difference (LSD) of studied traits among 30 genotypes.

Gen.	DH Mean	GY Mean	PLH Mean	SL Mean	TKW Mean	CT Mean
1	90.333 K	4048 CDEFGHI	90.667 CDE	11.333 ABCDE	38.333 ABCDE	26.633 ABC
2	92.667 FGHIJ	4720.7 A	103.67 BCDE	11 BCDE	38.333 ABCDE	22.733 G
3	95 CDE	4243.7 BCDEF	131 A	11.5 ABCDE	35.667 FGHI	24.633 BCDEFG
4	97.667 AB	3613.3 IJ	90 CDE	11.833 ABC	36.667 DEFG	27.1 A
5	93 EFGHI	3753.3 HIJ	104 BCDE	10.667 CDE	37 CDEF	27 AB
6	94.333 DEF	4212.7 BCDEFG	101.67 BCDE	11.667 ABCD	35.667 FGHI	25.833 ABCDE
7	93 EFGHI	4157.3 CDEFGH	86.667 E	12.333 AB	38.667 ABCD	25.1 ABCDEFG
8	91.667 HIJK	4417 ABCD	89.667 CDE	12.333 AB	37 CDEF	24.367 CDEFG
9	93.667 DEFGH	4244 BCDEF	102 BCDE	12.5 A	39.333 AB	24.9 ABCDEFG
10	94 DEFG	4494.3 ABC	101.67 BCDE	11 BCDE	39 ABC	23.833 EFG
11	92 GHIJK	4002.7 DEFGHIJ	105.33 BCD	11 BCDE	32.667 J	25.8 ABCDE
12	91 IJK	3785 GHIJ	102.33 BCDE	10.667 CDE	34.667 GHIJ	27.267 A
13	95 CDE	3921 EFGHIJ	103 BCDE	12.333 AB	36.667 DEFG	26.5 ABC
14	92.667 FGHIJ	4217 BCDEFG	103 BCDE	12.333 AB	40.333 A	24.933 ABCDEFG
15	95.667 BCD	3601.7 IJ	103 BCDE	12.333 AB	36.333 EFGH	27.067 A
16	95.667 BCD	3788.3 GHIJ	101.33 BCDE	10.667 CDE	33.333 J	26.767 ABC
17	95 CDE	3563 J	90.667 CDE	12 ABC	34 IJ	27.067 A
18	93 EFGHI	4000 DEFGHIJ	87.333 DE	11.333 ABCDE	34.333 HIJ	25.3 ABCDEF
19	94 DEFG	3602 IJ	93.667 BCDE	12 ABC	37 CDEF	26.767 ABC
20	92.333 FGHIJK	3839 FGHIJ	96.667 BCDE	11 BCDE	37 CDEF	25.967 ABCDE
21	92.333 FGHIJK	3931.7 EFGHIJ	106.67 BC	12.333 AB	36.333 EFGH	25.7 ABCDE
22	99.667 A	4346.7 ABCDE	101.33 BCDE	11.667 ABCD	38.667 ABCD	23.967 DEFG
23	90.333 K	4014.7 DEFGHI	104.67 BCDE	12.5 A	36.667 DEFG	26.267 ABCD
24	92.667 FGHIJ	4337 ABCDE	96.667 BCDE	12.167 AB	37.333 BCDEF	25.333 ABCDEF
25	91 IJK	4068 CDEFGH	93 BCDE	12 ABC	33 J	25.3 ABCDEF
26	93.333 EFGH	4168.7 BCDEFGH	99 BCDE	11.167 ABCDE	33.667 IJ	25.433 ABCDEF
27	92.667 FGHIJ	4414 ABCD	94.333 BCDE	10.167 E	37 CDEF	23.667 EFG
28	90.667 JK	3808.3 FGHIJ	104 BCDE	12.333 AB	35.667 FGHI	27 AB
29	97 BC	4479 ABC	97 BCDE	10.333 DE	40.333 A	25.733 ABCDE
30	99.667 A	4609.3 AB	110.67 B	12.167 AB	40.333 A	23.267 FG
LSD Value	2.0224	447.57	18.302	1.4118	2.1776	2.4283

REFERENCES

- Ajmal S. U., N. Zakir and M.Y. Mujahid (2009). Estimation of Genetic Parameters and Characterization Association in Wheat. *J. Agric. Biol. Sci.*, 1 (1):15-18.
- Akram Z., S. Ajmal, M. Munir (2008). Estimation of correlation coefficient among some yield parameters of wheat under rain-fed conditions. *Pak. J. Bot.*, 40 (4): 1777-1781.
- Anonymous (2010). World Wheat Crop to Be Third Largest Ever." Farmers Weekly. 152.13: 134. Academic Search Premier. Web. 13 Mar. 2013.
- Dokuyeu T. and A. Akkaya (1999). Path coefficient analysis and correlation of grain yield and yield components of wheat genotypes. *RACHIS*. 18: 17-20.
- Eivazi A.R., M.R. Naghavi, M. Hajheidari, *et al.*, (2007). Assessing Wheat (*Triticum aestivum* L.) Genetic diversity using quality traits, amplified fragments length polymorphism, simple sequence repeats and proteome analysis. *Ann Appl Biol.*, 152:81-91.
- Ilker, E., F. A. Tonk and M. Tosun (2010). Heterosis for yield and its components in bread wheat crosses among powdery mildew resistant and susceptible genotypes. *Pak. J. Bot.*, 42(1): 513-522.
- Maqbool, R., M. Sajjad, I. Khaliq, Aziz-ur-Rehman, A.S. Khan, S.H. Khan (2010). Morphological diversity and traits association in bread wheat (*Triticum aestivum* L.). *American-Eurasian J. Agric. Environ. Sci.*, 8: 216-224.
- Martin E., V. Cravero, A. Esposito, F. Lopez, L. Milanesi and E. Cointy (2008). Identification of markers linked to agronomic traits in globe artichoke. *Aust. J Crop Sci.*, 1:43-46.
- Memon, S., M. D. Qureshi, B. A. Ansari and M. A. Sial (2007). Genetic heritability for grain yield and its related characters in spring wheat (*Triticum aestivum* L.). *Pak. J. Bot.*, 39(5): 1503-1509.
- Mostafa K., H.F. Mohammad and M. Mohammad (2011). Genetic diversity of wheat genotypes based on cluster and principal component analyses for breeding strategies. *Aus. J. Crop Sci.*, 5(1): 17-24.
- Shabana M., D.Q. Mueen, A.A. Bashir and A.S. Mahboob (2007). Genetic heritability for grain yield and its related characters in spring wheat (*Triticum aestivum* L.). *Pak. J. Bot.*, 39(5): 1503-1509.
- Shah M. M., P.S. Baenziger, Y. Yen, K.S. Gill, B.M. Silva and K. Haliloglu (1999). Genetic analysis of agronomic traits controlled by wheat chromosome 3A. *Crop Sci.*, 39:96-102.
- Sial M.A., M.U. Dohat, K.A. Laghari, M.A. Arain, S.M. Mangrio and A.J. Pirzada (2010). Agronomic Performance of Semi-dwarf and dwarf Wheat genotypes. Nuclear Institute of Agriculture, Tandojam. *World Appl. Sci. J.*, 8: 30-33.
- Steel, R.G.D. and J.H. Torrie (1980). *Principles and Procedures of Statistics*. McGraw Hill Book Inc., New York.
- Yucel, C., F.S. Baloch, and H. Ozkan (2009). Genetic analysis of some physical of bread wheat grain (*Triticum aestivum* L.). *Turk. J. Agric. For.*, 33: 525-535.

(Accepted for publication April 2016)