

IN VITRO EVALUATION OF BIOCONTROL POTENTIAL OF *PAECILOMYCES* SPECIES AGAINST *SCLEROTIUM ROLFSII* AND *PYTHIUM APHANIDERMATUM*

Zahida Perveen¹, Nadia Ramzan¹, Nayara Noreen¹, Abdul Qayum Rajpuit² and Saleem Shahzad¹

¹Pest & Disease Research Lab., Department of Agriculture and Agribusiness Management, University of Karachi, Karachi-75270, Pakistan

*e-mail: zpbobotany@gmail.com

²Department of Plant Pathology, Sindh Agriculture University, Tandojam

ABSTRACT

Paecilomyces species viz., four strains of *Paecilomyces variotii*, *Paecilomyces lilacinus* and *Paecilomyces fumosoroseus* were evaluated *in vitro* for biocontrol potential against *Sclerotium rolfii* and *Pythium aphanidermatum* by dual culture plate method. Colonies of *Paecilomyces* species and *Sclerotium rolfii* met each other but *Sclerotium rolfii* later overgrew the colonies of tested fungi. whereas, growth of *Pythium aphanidermatum* inhibited by the *Paecilomyces* species.

Key words: *Paecilomyces*, Biocontrol potential, *Sclerotium rolfii*, *Pythium aphanidermatum*.

INTRODUCTION

The presence of *Sclerotium rolfii* (Sacc.), first time reported in Pakistan by Ahmed *et al.* (1984) on maize (*Zea mays* L.), is an economically important pathogen in warm, moist climate worldwide, causing diseases on more than 500 species of plants (Aycok, 1966). The pathogen causes stem and root rots on a wide variety of fruit and vegetable crops (Domsh *et al.*, 1980). This pathogen propagates by sclerotia under favourable conditions. After germination, sclerotia may cause chlorosis and wilting of entire plants (Yaqub and Shahzad, 2005). It causes serious damages to rice, apple, sugar beet, sunflower, and mash and mung beans in Pakistan (Ruqia, 2001; Shahzad and Ghaffar, 1995; Yaqub and Shahzad, 2005). Considering the health risks of living beings farmers prefer to use biocontrol agents against pathogens over fungicides.

Pythium aphanidermatum is a cosmopolitan pathogen with a wide host range (Plaats-Niterink, 1981; Domsh *et al.*, 1980). It is an aggressive pathogen and produces damping-off, root and stem rots, and blights of grasses, fruits, vegetables and crop plants. It is of economic concern on most annuals, cucurbits, and grasses. This fungus was associated with all the agronomic and horticultural crops in Pakistan (Shahzad and Ghaffar, 1995; Abdul-Haq and Shahzad, 1998; Lodhi *et al.*, 2013).

Of the various biocontrol agents, species of *Paecilomyces*, especially *P. lilacinus* and *P. variotii* have shown biocontrol potential against root infecting fungi like *Macrophomina phaseolina* (Shahzad and Ghaffar, 1987, 1989; Abbas *et al.*, 2011; Qureshi *et al.*, 2012). *Rhizoctonia solani* (Shahzad and Ghaffar, 1989; Mansoor *et al.*, 2007), *Fusarium solani* (Shahzad and Ghaffar, 1989; Siddiqui *et al.*, 2000; Mansoor *et al.*, 2007), and *Pythium aphanidermatum* (Hashem-al-Sheikh and Abdelzaher 2010). No literature on the use of *Paecilomyces* species against *S. rolfii*, and *P. aphanidermatum* in Pakistan is available. In our previous studies (Perveen, 2015), species of *Paecilomyces* showed biocontrol potential against root-knot nematodes. In view of the importance of the root rot and root-knot disease complexes caused by the interaction of the root infecting fungi and root knot nematodes (Shahzad and Ghaffar, 1992; Perveen and Ghaffar, 1998; Siddiqui *et al.*, 2000), biocontrol potential of *Paecilomyces* species was evaluated against *S. rolfii* and *P. aphanidermatum* *in vitro*.

MATERIAL AND METHODS

Microorganisms used: *Paecilomyces variotii* and *Paecilomyces fumosoroseus* were isolated from soil using serial dilution technique (Waksman and Fred, 1922). Culture of *Paecilomyces lilacinus* was acquired from ARS collection of entomopathogenic fungal cultures, USDA-ARSRW Centre for Agriculture & Health, USA. *Sclerotium rolfii* used in the present studies was isolated from sugar beet seed (Ruqia, 2003). Isolation of *Pythium aphanidermatum* was by baiting techniques (Harvey, 1925):

In vitro interaction of *Sclerotium rolfii* and *Pythium aphanidermatum* with *Paecilomyces* species:

Dual culture plate method: In dual culture plate essay, a 5mm diameter inoculum disc of four strains of *P. variotii*, *P. lilacinus* and *P. fumosoroseus* placed near the edge of a Petri plate containing PDA medium; a similar inoculums

disc of a root infecting fungus placed at the opposite end of Petri plate. There were three replicates for each treatment and the plates were incubated at $28 \pm 2^\circ\text{C}$. The colony diameters of pathogen and test microorganism recorded daily type of interaction observed using the following key:

A= Colonies of the test microorganism and the pathogen met each other. No further growth of either the microorganism or the pathogen observed.

B= Colonies of test organism and pathogen met each other; the pathogen later overgrew the colony of test microorganism.

C= Test microorganism produced a zone of inhibition.

RESULTS AND DISCUSSION

Colonies of four strains of *Paecilomyces variotii* met with *Sclerotium rolfsii* then pathogen overgrew on the tested fungi (Table 1; Fig. 1B, C, D, E). Colonies of four strains of *Paecilomyces variotii* and *Pythium aphanidermatum* met each other. No further growth of either the microorganism or the pathogen observed (Table 2; Fig. 1b, c, d, e). Colonies of *Paecilomyces lilacinus* produced zone of inhibition against *sclerotium rolfsii* then pathogen later overgrew the colony of test microorganism (Table 3; Fig. 1A). *Paecilomyces fumosoroseus* failed to inhibit the growth of *sclerotium rolfsii* but pathogen overgrew the tested fungi (Table 4; Fig. 1F). *Paecilomyces lilacinus* and *Paecilomyces fumosoroseus* produced a zone of inhibition against *Pythium aphanidermatum* (Table 3 and 4; Fig. 1a, f).

Table 1. Effect of *Paecilomyces variotii* on *in vitro* growth of *Sclerotium rolfsii*.

Biocontrol agents	Days of incubation	Colony Diameter (mm)		Comments
		Pathogen	Test fungi	
<i>Strain 1</i>	5	44	46	B
<i>Strain 2</i>	5	40	50	B
<i>Strain 3</i>	5	41	49	B
<i>Strain 4</i>	5	45	45	B

Table 2. Effect of *Paecilomyces variotii* on *in vitro* growth of *Pythium aphanidermatum*.

Biocontrol agents	Days of incubation	Colony Diameter (mm)		Comments
		Pathogen	Test fungi	
<i>Strain 1</i>	6	30	60	A
<i>Strain 2</i>	6	28	58	A
<i>Strain 3</i>	6	28	62	A
<i>Strain 4</i>	6	25	65	A

Table 3. Effect of *Paecilomyces lilacinus* on *in vitro* growth of *Sclerotium rolfsii* and *Pythium aphanidermatum*.

Biocontrol agents	Days of incubation	Colony Diameter (mm)		Comments
		Pathogen	Test fungi	
<i>Pythium aphanidermatum</i>	6	30	30	C
<i>Sclerotium rolfsii</i>	5	55	22	B

Table 4. Effect of *Paecilomyces fumosoroseus* on *in vitro* growth of *Sclerotium rolfsii* and *Pythium aphanidermatum*.

Biocontrol agents	Days of incubation	Colony Diameter (mm)		Comments
		Pathogen	Test fungi	
<i>Pythium aphanidermatum</i>	6	44	46	C
<i>Sclerotium rolfsii</i>	5	44	47	B




Fig. 1. Interaction of *Sclerotium rolfii* with A) *Paecilomyces lilacinus*, B) *Paecilomyces variotii* strain 1, C) *Paecilomyces variotii* strain 2, D) *Paecilomyces variotii* strain 3, E) *Paecilomyces variotii* strain 4 and F) *Paecilomyces fumosoroseus*. Interaction of *Pythium aphanidermatum* with a) *Paecilomyces lilacinus*, b) *Paecilomyces variotii* strain 1, c) *Paecilomyces variotii* strain 2, d) *Paecilomyces variotii* strain 3, e) *Paecilomyces variotii* strain 4 and f) *Paecilomyces fumosoroseus*.

Colonies of all the four strains of *P. variotii* and *S. rolfii* met each other; the pathogen later overgrew the colony of the test microorganism. Whereas, *P. lilacinus* produced zone of inhibition the pathogen later overgrew the colony of test microorganism. In case of *P. fumosoroseus* first there was inhibition of the growth of pathogen but later on pathogen overgrew the test organism. Shahzad and Ghaffar (1987) has reported that *Paecilomyces lilacinus* produced zone of inhibition against *Sclerotium oryzae* but after 3 days of inhibition, the pathogen over grew the colonies of *P. lilacinus*. Yaqub and Shahzad (2005) have reported that *Trichoderma* species inhibited the growth of

S. rolfsii in dual culture plates. Whereas, four strains of *P. variotii* inhibited the growth of *P. aphanidermatum* Colonies of the test microorganism and the pathogen met each other. No further growth of either the microorganism or the pathogen observed. *Paecilomyces lilacinus* and *P. fumosoroseus* produced zone of inhibition against *P. aphanidermatum*. Hashem-al-Sheikh and Abdelzaher (2010) reported the biological control of *Pythium spinosum* by *P. variotii*, *Aspergillus sulfureus*, and *Penicillium islandicum*. *P. fumosoroseus* produced 2-3mm wide zone of inhibition against *F. solani*, *R. solani* and *P. aphanidermatum*, and inhibited the growth of *M. phaseolina* when colonies met each other. The growth inhibition of pathogens by the biocontrol agents may be attributed to hyperparasitism or antibiosis (We *et al.*, 1986), or production of chitinase and B-1,3-glucanase enzymes that degrade the cell wall of the pathogens (Ahmed and Baker, 1987). Similarly, Muhammad and Amusa (2003) also concluded that the antagonists inhibit the growth of the pathogens either by producing biologically active metabolites, or by colonizing the agar surface much faster as compared to the pathogen.

These results suggest that the chemicals produced by different microorganisms have different effects on pathogens. It could be the reason for inefficacy of *Paecilomyces* species against *Sclerotium rolfsii*

REFERENCE

- Abbas, H., N. Javed, S.A. Khan, Irfan-ul-Haq, M. Asif and A. Safdar (2011). Integration of bioagent and bioproduct for the management of root-knot nematode, *Meloidogyne Incognita* on eggplant. *J. Biology, Agriculture and Healthcare*, 1(4): 31-36.
- Abdul-Haq, M. and S. Shahzad.1998. Oomycetes from soil of Bajour Agency, FATA. Pakistan. *Pak. J. Bot.*, 30: 305-306.
- Ahmed, J.S. and R. Baker (1987). Competitive saprophytic ability and cellulolytic activity of rhizosphere competent mutants of *Trichoderma harzianum*. *Phytopathology*, 77: 358-362.
- Ahmed, Y., M.S. Mirza and M. Aslam (1984). *Sclerotium rolfsii* on maize. *FAO Plant Prot. Bull.*, 32: 147.
- Aycock, R. (1966). Stem rot and other diseases caused by *Sclerotium rolfsii*. *NC. Agric. Exp. Sta.Tech. Bull*, No. 174. 202pp.
- Domsh, K.H., W. Gams, and T.H. Anderson (1980). *Compendium of soil fungi*. Vol. 1. London: Academic Press.
- Harvey, J.V. (1925). A study of the water molds and pythiums occurring in the soil of Chapel Hill. *J. Elisha Mitchell Scient. Soc.*, 41: 151-164.
- Hashem-al-Shaikh and H.M.A. Abdelzaher (2010). Isolation of *Aspergillus sulfureus*, *Penicillium islandicum* and *Paecilomyces variotii* from agricultural soil and their biological activity against *Pythium spinosum*, the damping-off organism of soybean. *Journal of Biological sciences*, 10: 178-189.
- Lodhi, A.M, M.A. Khanzada, S. Shahzad and A. Ghaffar (2013). Prevalence of *Pythium aphanidermatum* in agro-ecosystem of Sindh province of Pakistan. *Pak. J. Bot.*, 45(2): 635-642, 2013.
- Mansoor, F., V. Sultana and S. Ehteshamul-Haque (2007). Enhancement of biocontrol potential of *Pseudomonas aeruginosa* and *Paecilomyces lilacinus* against root rot of mungbean by a medicinal plant *Launaea nudicaulis* L. *Pak. J. Bot.*, 39(6): 2113-2119.
- Muhammad, S. and N.A. Amusa (2003). *In vitro* inhibition of some seedling blight inducing pathogens by compost inhabiting microbes. *African J. of Biotechnology*, 2(6):161-164.
- Parveen, S. and A. Ghaffar (1998). Use of *Paecilomyces lilacinus* in the control of *Fusarium oxysporum* root rot and *Meloidogyne javanica* root knot infection on tomato. *Pak. J. Nematol.*, 16: 71-75.
- Perveen, Z. (2015). *Use of Paecilomyces species against root rot fungi and root knot nematodes*. Ph.D Thesis. Department of Agriculture and Agribusiness Management, University of Karachi, Karachi, Pakistan.
- Plaats-Niterink, A.J. Vander (1981). *Monograph of the genus Pythium*. Studies in Mycology. No. 21. Centraalbureau voorSchimmelcultures, Baarn. 242 pp.
- Qureshi, S.A., Ruqia, V. Sultana, J. Ara and S. EhteshamulHaque (2012). Nematicidal potential of culture filtrates of soil fungi associated with rhizosphere and rhizoplane of cultivated and wild plants. *Pak. J. Bot.*, 44: 1041-1046.
- Ruqia, B. (2001). *Studies on seed-borne mycoflora of sugarbeet (Beta vulgaris)*. M.Sc. Thesis, Department of Botany, University of Karachi, Karachi.30 pp.
- Shahzad, S. and A. Ghaffar (1987). Field application of *Paecilomyces lilacinus* and furadan for the control of root-knot disease of okra and mung. *Int. Nematol. Network Newsl.*, 4: 33-34.
- Shahzad, S. and A. Ghaffar (1989). Use of *Paecilomyces lilacinus* in the control of root rot and root- knot disease complex of okra and mungbean. *Pak. J. Nematol.*, 7: 47-53.
- Shahzad, S. and A. Ghaffar (1992). Root-knot and root-rot disease complex and their control. pp. 169-180. In: *Recent Advances in Nematology*. K. Dwiwedi (Ed.). Bioved Research Society, Allahabad, India.

- Shahzad, S. and A. Ghaffar (1995). New host records of root infecting fungi in Pakistan. *Pak. J. Bot.*, 27: 209-216.
- Siddiqui, I.A., S.A. Qureshi, V. Sultana, S. Ehteshamul-Haque and A. Ghaffar (2000). Biological control of root rot-root knot disease complex of tomato. *Plant and Soil*, 227: 163–169.
- Waksman, S.A. and E.B. Fred (1922). A tentative outline of the plate method for determining the number of microorganisms in the soil. *Soil Sci.*, 14: 27-28.
- We, W.S., S.D. Liu and S. Tschen (1986). Hyperparasitic relationship between antagonists and *Rhizoctonia solani*. *Plant Prot. Bull.*, 28: 91-100.
- Yaqub, F. and S. Shahzad (2005). Pathogenicity of *Sclerotium rolfsii* on different crops and effect of inoculum density on colonization of mungbean and sunflower. *Pak. J. Bot.*, 37: 175-180.

(Accepted for publication April 2015)