



Exogenous application of ascorbic acid stimulates growth and photosynthesis of wheat (*Triticum aestivum* L.) under drought

Samina Malik* and Muhammad Ashraf

Department of Botany, University of Agriculture, Faisalabad, Pakistan

Abstract

Drought causes considerable reduction in plant growth. A hydroponic experiment was conducted to appraise the potential role of exogenously applied ascorbic acid in alleviating the effect of drought on wheat. Two contrasting wheat genotypes, a drought tolerant cultivar Chakwal-86 and a drought sensitive strain 6544-6 were used in the study. Drought was induced by dissolving 20% Polyethylene glycol (PEG₈₀₀₀) in the nutrient solution producing -0.6MPa osmotic stress. Drought caused a significant decrease in chlorophyll pigments and net photosynthesis resulting in growth reduction of both wheat genotypes. However, this decrease was more severe in the genotype 6544-6 compared to Chakwal-86. Ascorbic acid (AsA) was applied through rooting medium, as a foliar spray and seed soaking treatment. Ascorbic acid treated seedlings of both genotypes maintained higher chlorophyll contents, net photosynthesis and growth compared to the non-treated plants. Of the three different modes of ascorbic acid application, rooting medium was more effective in alleviating the adversities of drought in wheat.

Key words: Wheat, drought, ascorbic acid, net photosynthesis, growth, chlorophyll

Introduction

Limited availability of water is a serious constraint to agricultural production of major crops (Shao *et al.*, 2009) because water is vital factor in plant development. Reduced supply of water is known to hamper important physiological and biochemical mechanisms leading to reduction in plant growth. Substantial yield losses have been observed in different crops due to reduced supply of water even for a short period of time (Pinheiro *et al.*, 2005). Drought stress causes reduced stomatal conductance resulting in decreased net photosynthetic rate. Chlorophyll degradation due to drought stress also inhibits photosynthetic rate in wheat (Moaveni, 2011). Drought stress deteriorates membranes which adversely affects a number of metabolic reactions occurring within the cell (Ashraf, 2009). Membrane stability is adversely affected due to oxidative damage caused by higher cellular concentrations of reactive oxygen species (ROS) like hydrogen peroxide, superoxide, singlet oxygen (Navarri-Izzo *et al.*, 1994) produced as a result of drought stress. Reactive oxygen species due to their reactive nature negatively interact with DNA, proteins, lipids and pigments causing huge cellular damage (Ashraf, 2009).

A number of enzymatic and non-enzymatic antioxidants are produced in plants in response to abiotic stresses which save plant from oxidative damage caused by ROS (Ashraf, 2009). Major enzymatic antioxidants reported

are superoxide dismutase (SOD), catalase (CAT), peroxidase (POD), ascorbate peroxidase (APX) whereas, ascorbic acid (vitamin C) and tocopherols are the main non-enzymatic antioxidants exploited by plants under stressful conditions to ameliorate the adverse effects imposed by ROS (Mittler, 2002). A few studies report that exogenously applied ascorbic acid ameliorates adverse effects of drought (Singh *et al.*, 2001; Amin *et al.*, 2009; Dolatabadian *et al.*, 2009a; Dolatabadian *et al.*, 2010; Khalil *et al.*, 2010). However, they have applied ascorbic acid either as a foliar application or in the rooting medium under soil conditions. In the present study, ascorbic acid was applied by three different modes; pre-sowing seed treatment, rooting medium and foliar spray to wheat plants under drought stress. Objective of the study was to investigate the comparative effect of ascorbic acid in three different modes.

Materials and Methods

Two wheat genotypes, a drought tolerant cultivar, Chakwal-86 and a drought sensitive genotype 6544-6 were used in the study. Ten seeds of each genotype were sown on wet filter paper in Petri plates kept in a growth chamber. The temperature and humidity was not controlled and varied between 15-20°C and 30-60%, respectively. The light intensity was maintained at 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 10 hours daily using fluorescent tubes. A week after germination, seedlings of both genotypes were transplanted

*Email: saminatmalik@yahoo.com

into plastic tubs containing Hoagland's nutrient solution (Epstein, 1972). The tubs were placed in a covered wirehouse to protect against rainfall. Aeration was maintained in the hydroponic system with the help of an electric pump. The environmental conditions during the course of experiment were 20.50-13.75°C average day and night temperature and 41.71-76.28% average relative humidity. A week after transplantation when the seedlings were 2 weeks old, PEG₈₀₀₀ was dissolved in the hydroponic culture in three equal doses with an interval of one day until the final concentration of 20% to develop -0.6 MPa drought stress. Optimized concentrations of ascorbic acid evaluated from the preliminary trials (data not mentioned) were exogenously applied as a foliar spray, seed priming and rooting medium treatments soon after dissolving PEG in the nutrient solution. Twenty four plastic tubs were used. There were twenty plants (10 plants of each genotype) in a tub. There were 3 tubs for each treatment. The experimental plan was a completely randomized design with three replications. The treatment breakup of the 24 tubs of hydroponic system was as below:

- Hoagland's nutrient solution
- Hoagland's nutrient solution + PEG
- Hoagland's nutrient solution + 0.5mM AsA in rooting medium
- Hoagland's nutrient solution + PEG and 0.5mM AsA in rooting medium
- Hoagland's nutrient solution + 1mM AsA foliar spray
- Hoagland's nutrient solution + PEG and 1mM AsA foliar spray
- Hoagland's nutrient solution + 1mM AsA applied as seed priming
- Hoagland's nutrient solution + PEG and 1mM of AsA as seed priming

For rooting medium treatment, 0.5mM AsA was added to nutrient solution of the plants. Seeds were soaked for 10 h in a solution of 1mM AsA for seed priming. Foliar spray was prepared by dissolving 1mM AsA in distilled water with the addition of 0.1% tween-20 as a surfactant to increase the penetration of ascorbic acid solution into the leaves. Foliar spray was applied twice, once on 2 weeks old seedlings and then on 3 weeks old seedlings. The data for gas exchange parameters were recorded on 3rd leaf of each plant using an open system portable infra red gas analyzer (Analytical Development Company, Hoddeson, England) after five weeks of transplantation. The plants were then harvested and fresh biomass was recorded with the help of an electronic balance. Harvested fresh leaves were used for the quantification of photosynthetic pigments (Chlorophyll "a" and "b") following the procedure described by Arnon

(1949). The data were subjected to analysis of variance using COSTAT software. LSD was calculated to see the differences among the means (Steel *et al.*, 1997).

Results and Discussion

A marked reduction in fresh biomass was recorded when plants of both wheat genotypes were exposed to PEG-induced drought stress. However, cultivar Chakwal-86 exhibited relatively higher shoot and root fresh biomass compared to the genotype 6544-6 (Figure 1; Table 1). Reduction in plant biomass due to water stress has been widely reported (Kamara *et al.*, 2003; Kusaka *et al.*, 2005; Amin *et al.* 2009). In the present investigation, ascorbic acid applied via different methods improved growth of the treated plants compared to non-treated ones which indicated that AsA helped plants to mitigate adverse effects of drought stress. The results of the present study on the role of ascorbic acid in circumventing the adverse effects of drought on plant fresh biomass are in line with some earlier reports. For example, Shalata and Neuman (2001) reported the ameliorative effect of ascorbic acid on fresh biomass of tomato seedlings when exposed to PEG or NaCl-induced osmotic stress. They found 0.5 mM as the most effective concentration of ascorbic acid in the rooting medium. The present study also revealed that 0.5 mM AsA concentration was the most effective in rooting medium. Of different modes of AsA application in the present study, rooting medium application was more effective compared to that of other modes. Athar *et al.* (2009) also applied ascorbic acid through different modes to counteract the adverse effects of salinity on the growth of wheat plants and found that rooting medium was the most effective mode of ascorbic acid application.

The chloroplast pigments, chlorophylls "a" and "b" play an important role in photochemical reactions (Taiz and Zieger, 2006). The present study showed that drought stress significantly reduced the leaf chlorophyll "a" and "b" content of drought sensitive as well as drought tolerant genotype. This is in line with what has been earlier reported in sunflower (Zhang and Kirkham, 1996; Manivannan *et al.*, 2007), rice (Pattanagul, 2011), barley (Havaux, 1998), maize Dolatabadian *et al.* 2009a), okra (Amin *et al.* 2009) and wheat (Moaveni, 2011). The decrease of chlorophyll content under water limited conditions is reported to take place because of its photo-oxidation and degradation under drought (Anjum *et al.*, 2011). Under drought stress, degradation of chlorophyll takes place due to the increased activity of chlorophyllase enzyme (Mihailovic *et al.*, 1997). Non-stomatal decrease in photosynthesis due to drought stress in the present study may have been as a result of chlorophyll degradation (Guo and Li, 1996; Sairam *et al.*, 1998; Anjum *et al.*, 2011).




Figure 1: Comparison of shoot fresh biomass, root fresh biomass, chlorophyll a (Chl a) and chlorophyll b (Chl b) contents of drought stressed and non-stressed 6 week old plants of two wheat genotypes, Chakwal-86 (Ch-86) and 6544-6 with ascorbic acid application through different modes (mean $\pm$ S.E.)

The cultivar Chakwal-86 maintained higher chlorophyll “a” and “b” content compared to that of genotype 6544-6 under drought stress. Earlier studies have also reported that the chlorophyll content of drought resistant genotypes of barley, wheat and maize were higher compared to the sensitive genotypes in drought stress conditions (Pastori and Trippi, 1992; Rong-Hua *et al.*, 2006). The present study revealed that exogenous application of ascorbic acid via rooting medium helped plants maintaining the chlorophyll pigments and hence mitigated the adverse effects of drought stress. These findings are in line with some earlier reports

on *Cassia* (Singh *et al.*, 2001), okra (Amin *et al.*, 2009), wheat (Azzedine *et al.*, 2011), and maize (Dolatabadian *et al.*, 2009a). Reactive oxygen species produced under stress conditions have been reported to cause pigment degradation (Sairam and Saxena, 2000; Anjum *et al.*, 2011). However, ascorbic acid being an antioxidant actively scavenges these ROS, thereby reducing the chlorophyll degradation under stress (Ashraf, 2009).




Figure 2: Comparison of net photosynthetic rate (P_n), transpiration rate (E), stomatal conductance (g_s), sub-stomatal CO_2 concentration (C_i) of drought stressed and non-stressed 6 week old plants of two wheat genotypes, Chakwal-86 (Ch-86) and 6544-6 with ascorbic acid application in different modes (mean $\pm$ S.E.)

Drought stress is also known to cause a significant reduction in the gas exchange attributes; net photosynthesis, transpiration rate and stomatal conductance of plants (El-hafid *et al.*, 1998; Shah and Paulsen, 2003; Flexas *et al.*, 2004; Ali and Ashraf, 2011). In the present investigation, drought stress caused a marked reduction in net photosynthesis, transpiration rate and stomatal conductance in both wheat genotypes. However, the drought tolerant cv. Chakwal-86 was superior to the drought sensitive genotype 6544-6 with respect to these gas exchange attributes (Figure 2; Table 2). Drought-induced reduction in photosynthesis and transpiration rate has been reported earlier in a number of crops including wheat (El Hafid *et al.*, 1998), maize (Ali and Ashraf, 2011), and sorghum (Kreig and Hutmacher, 1983). In the present study, exogenous application of ascorbic acid mitigated the adverse effects of drought on photosynthesis in both wheat genotypes by increasing stomatal conductance. This could have also been due to the fact that ascorbic acid as an antioxidant has the ability to mitigate the negative effects of stress on plants by

neutralizing harmful oxidants which have been reported to damage plant membranes such as the thylakoid membranes of chloroplasts (Miguel *et al.*, 2006; Dolatabadian *et al.*, 2009b). In the present investigation, ascorbic acid applied through the rooting medium was found to be more effective compared to the other modes of application in alleviating the adverse effects of drought on different gas exchange attributes of both wheat cultivars.

In the present study, ascorbic acid increased transpiration rate of wheat under drought stress compared to non-treated plants. Similar results were reported in okra plants under drought using ascorbic acid as a foliar spray (Amin *et al.*, 2009). It has been reported that maintenance of water status is regulated by stomatal conductance and rate of transpiration (Ashraf, 2009). In the present study, there was a significant decline in stomatal conductance of drought tolerant as well as drought sensitive genotype under drought. Application of ascorbic acid improved the stomatal conductance of plants compared to non-treated plants.

Table 1: Mean squares from analysis of variance of data for shoot fresh weight, root fresh weight, chlorophyll a (Chl a) and chlorophyll b (Chl b) contents of drought stressed and non-stressed 6 week old plants of two wheat genotypes (Chakwal-86 and 6544-6) with ascorbic acid application in different modes

Source	df	Shoot fresh wt	Root fresh wt	Chl a	Chl b
Main effects					
Genotype	1	3.44***	0.55***	0.62***	0.12*
Drought	1	7.83***	0.97***	0.82***	0.007ns
AsA	3	0.06ns	0.06ns	0.04***	0.01ns
Interaction					
Genotype x drought	1	0.20*	0.03ns	0.17***	0.04ns
Genotype x AsA	1	0.25***	0.04ns	0.04***	0.02ns
Drought x AsA	3	0.73***	0.26***	0.07***	0.09**
Genotype x drought x AsA	3	0.17**	0.03ns	0.02**	0.00655ns

Table 2: Mean squares from analysis of variance of data for net photosynthetic rate (P_n), transpiration rate (E), stomatal conductance (g_s), sub-stomatal CO_2 concentration (C_i) of drought stressed and non-stressed 6 week old plants of two wheat genotypes (Chakwal-86 and 6544-6) with ascorbic acid application in different modes

Source	df	P_n	E	g_s	C_i
Main effects					
Genotype	1	14.73*	0.47*	0.0007ns	43.12ns
Drought	1	23.98***	2.46***	0.067***	134.29**
AsA	3	4.72***	0.50**	0.0004ns	282.11***
Interaction					
Genotype x drought	1	0.12ns	0.34ns	0.0003ns	47.16ns
Genotype x AsA	1	0.36ns	0.18ns	0.0008ns	78.47ns
Drought x AsA	3	1.98*	0.26ns	0.0001ns	243.36***
Genotype x drought x AsA	3	0.01ns	0.17ns	0.0009ns	139.26**

*, **, *** = significant at 0.05, 0.01 and 0.001 levels, respectively, ns = non-significant

Although the ascorbic acid application was effective through all three modes, the effect of ascorbic acid through the rooting medium was relatively higher. This may be expected as in rooting medium, a continuous supply of ascorbic acid is available to the plant compared to the other two modes.

Conclusion

Overall, ascorbic acid application through rooting medium was comparatively more effective in overcoming the adverse effects of drought stress in wheat compared to that of foliar spray or seed priming treatment in terms of photosynthetic rate and plant growth. Ascorbic acid treated plants showed higher net photosynthetic rate, transpiration and stomatal conductance compared to non-treated ones. Moreover, relatively less stress-induced degradation of chlorophyll pigments took place in plants supplied with ascorbic acid. Thus, it may be concluded that exogenously applied ascorbic acid is effective in ameliorating the adverse effects of drought stress.

References

- Ali, Q. and M. Ashraf. 2011. Induction of drought tolerance in maize (*Zea mays* L.) due to exogenous application of trehalose: growth, photosynthesis, water relations and oxidative defence mechanism. *Agronomy and Crop Science* 1: 1-14.
- Amin, B., G. Mahlegah, H.M.R. Mahmood and M. Hossein. 2009. Evaluation of interaction effect of drought stress with ascorbate and salicylic acid on some of physiological and biochemical parameters in okra (*Hibiscus esculentus* L.). *Research Journal of Biological Sciences* 4: 380-387.
- Anjum, S., X.Y. Xie, L.C. Wang, M.F. Saleem, C. Man and L. Wang. 2011. Morphological, physiological and biochemical responses of plants to drought stress. *African Journal of Agricultural Research* 6: 2026-2032.
- Arnon, D.I. 1949. Copper enzymes in isolated chloroplasts, polyphenoloxidase in *Beta vulgaris*. *Plant Physiology* 24: 1-15.
- Ashraf, M. 2009. Biotechnological approach of improving plant salt tolerance using antioxidants as markers. *Biotechnology Advances* 27: 84-93.
- Athar, H.R., A. Khan and M. Ashraf. 2009. Inducing salt tolerance in wheat by exogenously applied ascorbic acid through different modes. *Journal of Plant Nutrition* 32: 1799-1817.
- Azzedine, F.H., Gheroucha and M. Baka. 2011. Improvement of salt tolerance in durum wheat by ascorbic acid application. *Journal of Stress Physiology and Biochemistry* 7: 27-37.
- Dolatabadian A., S.A.M. Modarressanavy and K.S. Asilan. 2010. Effect of ascorbic acid foliar application on yield, yield component and several morphological traits of grain corn under water deficit stress conditions. *Notulae Scientia Biologicae* 2: 45-50.
- Dolatabadian, A., A.M. Sanavi and M. Sharifi. 2009a. Alleviation of water deficit stress effects by foliar application of ascorbic acid on *Zea mays* L. *Journal of Agronomy and Crop Science* 195: 347-355.
- Dolatabadian, A., A.M. Sanavi and M.S. Harifi. 2009b. Effect of ascorbic acid (Vitamin C) leaf feeding on antioxidant enzymes activity, proline accumulation and lipid peroxidation of canola (*Brassica napus* L.) under salt stress condition. *Journal of Science and Technology of Agriculture and Natural Resources* 13: 611-620.
- El-Hafid, R., D.H. Smith, M. Karrou and K. Samir. 1998. Physiological responses of spring durum wheat cultivars to early-season drought in a mediterranean environment. *Annals of Botany* 81: 363-370.
- Epstein, E. 1972. Mineral nutrition of plants: principles and perspectives. John Wiley and Sons, New York.
- Flexas, J., J. Bota, J.F. Loreto, G. Cornic and T.D. Sharkey. 2004. Diffusive and metabolic limitations to photosynthesis under drought and salinity in C_3 plants. *Plant Biology* 6: 1-11.
- Guo, P. and M. Li. 1996. Studies on photosynthetic characteristics in rice hybrid progenies and their parents. I. chlorophyll content, chlorophyll-protein complex and chlorophyll fluorescence kinetics. *J. Tropical and Subtropical Botany* 4: 60-65.
- Havaux, M. 1998. Carotenoids as membrane stabilizers in chloroplasts. *Trends in Plant Science* 3: 147-151.
- Kamara, A.Y., A. Menkir, B.B. Apraku and O. Ibikunle. 2003. The influence of drought stress on growth, yield and yield components of selected maize genotypes. *J. Agricultural Science* 141: 43-50.
- Khalil, S.E., G. Nahed, A. Aziz and L.A.H. Bedour. 2010. Effect of water stress and ascorbic acid on some morphological and biochemical composition of *Ocimum basilicum* plant. *Journal of American Science* 6: 33-46.
- Krieg, D.R. and R.B. Hutmacher. 1983. Photosynthetic rate control in sorghum: Stomatal and nonstomatal factors. *Crop Science* 712: 789-796.
- Kusaka M., A.G. Lalusin and T. Fujimura, 2005. The maintenance of growth and turgor in pearl millet (*Pennisetum glaucum* (L.) Leeke) cultivars with different root structures and osmo-regulation under drought stress. *Plant Science* 168:1-14.
- Manivannan, P., C.A. Jaleel, B. Sankar, A. Kishorekumar, R. Somasundaram, G.M. Alagu Lakshmanan and R. Panneerselvam. 2007. Growth, biochemical modifications and proline metabolism in *Helianthus*

- annuus* L. as induced by drought stress. *Colloids Surfaces and Biointerfaces* 59: 141-149.
- Miguel, A., J. Rosales, M. Ruiz, J. Hernandez, T. Soriano, N. Castilla and L. Romero. 2006. Antioxidant content and ascorbate metabolism in cherry tomato exocarp in relation to temperature and solar radiation. *Journal of the Science of Food and Agriculture* 86: 1525-1551.
- Mihailović, N., M. Lazarević and Z. Dželetović. 1997. Chlorophyllase activity in wheat, *Triticum aestivum* L. leaves during drought and its dependence on the nitrogen ion form applied. *Plant Science* 129: 141-6.
- Mittler, R. 2002. Oxidative stress, antioxidant and stress tolerance. *Trends in Plant Science* 7: 405-410.
- Moaveni, P. 2011. Effect of water deficit stress on some physiological traits of wheat (*Triticum aestivum*). *Agricultural Science Research Journal* 1:64-68.
- Navarri-Izzo, F., C. Pinzino, M.F. Quartacci and C.L.M. Sgherri. 1994. Intracellular membranes: kinetics of superoxide production and changes in thylakoids of resurrection plants upon dehydration and rehydration. *Proc. Royal Soc. Edinburgh, UK*, 102: 187-191.
- Pastori, G.M. and V.S. Trippi. 1992. Oxidative stress induces high rate of glutathione reductase synthesis in a drought resistant maize strain. *Plant Cell Physiology* 33: 957-961.
- Pattanagul, W. 2011. Exogenous abscisic acid enhances sugar accumulation in rice (*Oriza sativa* L.) under drought stress. *Asian Journal of Plant Science* 10: 212-219.
- Pinheiro, H.A., F.M. DaMalta, R. Agnaldo, M. Chaves, M.E. Loureiro and C. Ducatta. 2005. Drought tolerance is associated with rooting depth and stomatal control of water use in clones of *Coffea canephora*. *Annals of Botany* 96: 101-108.
- Rong-hua, L.I., G.U.O.P. Guo, M. Baum, S. Grando and S. Ceccarelli. 2006. Evaluation of chlorophyll content and fluorescence parameters as indicators of drought tolerance in barley. *Agriculture Science (China)* 5: 751-757.
- Sairam, R.K. and D.C. Saxena. 2000. Oxidative stress and antioxidants in wheat genotypes: Possible mechanism of water stress tolerance. *Journal of Agronomy and Crop Science* 184: 55-61.
- Sairam, R.K., P.S. Deshmukh and D.C. Saxena. 1998. Role of antioxidant systems in wheat genotypes tolerance to water stress. *Biologia Plantarum* 41: 387-394.
- Shah, N.H. and G.M. Paulsen. 2003. Interaction of drought and high temperature on photosynthesis and grain-filling of wheat. *Plant and Soil* 257: 219-226.
- Shalata, A. and P.M. Neumann. 2001. Exogenous ascorbic acid (vitamin C) increases resistance to salt stress and reduces lipid peroxidation. *Journal of Experimental Botany* 52: 2207-2211.
- Shao, H.B., L.Y. Chu, C.A. Jaleel, P. Manivannan, R. Panneerselvam and M.A. Shao. 2009. Understanding water deficit stress-induced changes in the basic metabolism of higher plants-biotechnologically and sustainably improving agriculture and the ecoenvironment in arid regions of the globe. *Critical Review in Biotechnology* 29: 131-151.
- Singh, D.V. G.C. Srivastava and M.Z. Abdin. 2001. Amelioration of negative effect of water stress in *Cassia angustifolia* by benzyladenine and/or ascorbic acid. *Biologia Plantarum* 44: 141-143.
- Steele, R.G.D., J.H. Torrie and D.A. Dickey. 1997. Principles and Procedures of Statistics: A Biometrical Approach. McGraw-Hill, New York, USA.
- Taiz, L. and E. Zeiger. 2006. Plant Physiology, 4th Ed. Sinauer Associates Inc. Publishers, Sunderland, Massachusetts, USA.
- Zhang, J.X. and M.B. Kirkham. 1996. Lipid peroxidation in sorghum and sunflower seedlings as affected by ascorbic acid, benzoic acid and propyl gallate. *Journal of Plant Physiology* 149: 489-493.