Soil & Environ. 28(2): 119-123, 2009 www.se.org.pk ISSN: 2074-9546 (Print) ISSN: 2075-1141 (Online)



Siderophore production by mycorrhizal sorghum roots under micronutrient deficient condition

N. Aliasgharzad*, E. Shirmohamadi and S. Oustan

Department of Soil Science, Faculty of Agriculture, University of Tabriz, Tabriz, I.R. of Iran

Abstract

It has widely been accepted that mycorrhizal symbiosis improves micronutrients uptake by most of the plants. In this study, sorghum (Sorghum bicolor L.) plants were grown in sterile perlite and were inoculated with either Glomus etunicatum (GE) or G.intraradices (GI), while the control set was left un-inoculated. Rorison's nutrient solution with three levels of 0, half and full strength (C_0 , $C_{0.5}$ and C_1 , respectively) of Fe, Cu, Zn and Mn was applied to the pots during 85 days of growth period. Chrome azurol-S assay was used for determination of siderophores in root leachates on 45, 65 and 85 days after sowing (DAS). Siderophore production per unit volume of root was higher in mycorrhizal than non-mycorrhizal plants. Both GE and GI were efficient fungi in this respect. Siderophore production was significantly induced at C_0 level of the micronutrients. Amount of siderophores produced on 45 and 85 DAS was more than 65 DAS. Mycorrhizal root colonization by GE or GI was not significantly affected by micronutrient levels.

Keywords: Arbuscular mycorrhizal fungi, siderophore, micronutrient deficiency, sorghum

Introduction

Arbuscular mycorrhizal fungi (AMF) enhance nutrient uptake by establishing a hyphal net inside and around the roots (Lee and George, 2005). Increment of Fe, Zn and Cu uptake by mycorrhizal plants have been reported world wide (Clark and Zeto, 1996; Caris et al., 1998; Liu et al., 2000; Lee and George, 2005; Silvia et al., 2005; Schreiner, 2007). The mechanism(s) involved are not well documented, although some of these have been suggested as follows: (1) siderophore production by both fungi and roots (Caris et al., 1998), (2) reduction of Fe³⁺ and Mn⁴⁺ to Fe²⁺ and Mn²⁺ by mycorrhizal roots, which led to their higher solubility (Cairney and Ashford, 1989) and (3) organic acid and H⁺ production (Cairney and Ashford, 1989; Angle et al., 1989). Siderophores and phytosiderophores are low molecular chelators which are produced by microbes and plant roots, respectively and have high affinity to Fe³⁺, but they can also make complexes with other metal ions such as Zn, Cu, Mn, Cd, Cr, Ni and Al (Romheld, 1987; Buyer et al., 1993; Bakkaus et al., 2006). Siderophore is a metabolic product of a fungus (or other organism) which binds iron and facilitates its transport from the environment into the microbial cell. In fact, siderophores and phytosiderophores are secondary metabolites from inducible genes which are generally expressed by Fe³⁺ and to some extent by Zn, Cu and Mn deficiencies (Cakmak et al., 1994; Zhang et al., 1991).

Several plant species are able to produce phytosiderophore but they can also use microbial siderophores as metal ion transporters (Romheld, 1987; Leyval and Reid, 1991). Some fungal species produce more siderophore than bacteria (Milagres *et al.*, 1999). Ericoid mycorrhizal fungi release ferricrocin or fusigen as the main siderophores. Ferricrocin was also shown to be produced by the ectomycorrhizal fungi, *Cenococcum geophilum* and *Hebeloma crustuliniforme*. Arbuscular mycorrhizal fungi are reported to enhance Fe-uptake rates of associated host plants which can be taken as an indication that mycorrhizal siderophores of a yet unknown structure may be involved (Lee and George, 2005; Schreiner, 2007).

Mycorrhizal *Hilaria janesii* grass, which showed greater Fe uptake than non-mycorrhizal controls, tested positively when bioassayed for hydroxamate siderophores (Cress *et al.*, 1986). However, it has to be stressed that these tests were carried out under non-axenic conditions; hence these results cannot be taken as proof that *Glomus* species may produce hydroxamate siderophore. In non-setile soil, it may well be that the hydroxamate siderophores detected, were produced by microorganisms other than arbuscular mycorrhizal fungi. Sound evidence for the nature of siderophore released by AMF can only be obtained when the fungi together with suitable host plants are grown under pure culture conditions.

^{*}Email: n_aliaasgharzad@yahoo.com

In this study, siderophore production by mycorrhizal sorghum roots inoculated with either *G.etunicatum* or *G.intraradices* in axenic condition, were quantitatively determined using chrome azurol-S assay (Schwyn and Neilands, 1984) by inducing micronutrients (Fe, Zn, Cu and Mn) deficiencies.

Materials and Methods

Mycorrhizal inoculum production

Two species of arbuscular mycorrhizal fungi, Glomus & Gerdemann) etunicatum (Becker (GE) and G.intraradices (Schenck & Smith) (GI) were propagated with sorghum plants in 7 liter pots containing sterile sandy loam soil (Aliasgharzad et al., 2001). Rorison's nutrient solution, 20 mM Ca(NO₃)₂.4H₂O; 10 mM MgSO₄.7H₂O; 10 mM K₂HPO₄.3H₂O; 0.5 mM FeNaEDTA; 0.1 mM MnSO₄.4H₂O; 0.5 mM H₃BO₃; 0.01 mM (NH₄)₆ Mo₇O₂₄.4H₂O; 0.02 mM ZnSO₄.7H₂O and 0.015 mM CuSO₄.5H₂O in deionized water (Merryweather and Fitter, 1991) with 1/2 strength of phosphorus was added to the pots twice a week to bring the soil moisture to field capacity. Pots were kept in growth room with 28/20 ±2°C day/night temperatures and 16h photoperiod. After four months, top plants were cut off and pot materials containing soil, mycorrhizal roots, hyphae and spores were throughly mixed and used as fungal inoculum. Root colonization percentage (Giovanetti and Mosse, 1980) and number of spores per 10 g soil (Gerdemann and Nicolson, 1963) were assessed to determine inoculum potential. Both inocula had an average of 65% root colonization and ~150 spores per 10 g soil.

Plant culture

Sorghum (Sorghum bicolor L.) seeds were surface sterilized with 0.5% sodium hypochlorite for 15 min, and 10 seeds were sown in pots containing 2.8 liter acid washed and sterilized perlite. Fungal inocula were rinsed three times with distilled water to minimize their micronutrients content. Each pot received 60 g mycorrhizal inoculum as a layer of 0.5 cm thickness, 5 cm below the seeds. Control pots (non-mycorrhizal) received 60 g autoclaved inoculum. Two weeks after sowing, these were thinned to 3 plants per pot. Rorison's nutrient solution with three levels of 0, half and full strength (C₀, C_{0.5}, C₁, respectively) of Fe, Zn, Cu and Mn was applied to the pots twice a week during total growth period of 85 days. Pots were kept in growth room with $28/20 \pm 2^{\circ}$ C day/night temperatures and 16h photoperiod.

Root leachate collection

On 45, 65 and 85 days after sowing (DAS), pot materials were eluted with two pore volume (2.5 liter) of

distilled water (DW) to remove nutrients from root zone. Three hours after subjecting root to the DW, root leachates were collected by adding 500mL DW to each pot (Koa *et al.*, 2001).

Siderophores assessment

Siderophore concentrations were immediately measured in the collected leachates as described below. Chrome azurol-s (CAS) solution was prepared as described by Schwyn and Neilands (1984). Five mL of CAS solution and one mL of root leachate were mixed in a test tube and absorbance was measured at 698 nm after one hour. Siderophore concentration was calculated using a DTPA (diethylenetriaminepentaacetic acid) standard calibration curve. Standard solutions containing 0, 25, 75, 100, 125 and 150 µmol DTPA were mixed in 1/5 ratio with CAS before subjecting to spectrophotometry. The amount of siderophore was expressed in µmol per total volume of leachate in one hour (µmol pot⁻¹ h⁻¹) or µmol per unit volume of root in one hour (μ mol cm⁻³ h⁻¹). The latter unit was applied just for data obtained at harvest time (85 DAS). Root volume was determined by suspending total root system in water and re-measuring water height in cylinder.

Root colonization

Eighty five days after sowing, plants were harvested and whole root system was washed, while removing perlite particles gently and its total volume was measured as described above. Fine feeding roots (0.5 g fresh weight) were sub-sampled, cleared in 10% KOH and stained with trypan blue. Root mycorrhizal colonization percentage was determined by gridline intersect method (Giovanetti and Mosse, 1980).

Statistical analysis

A factorial completely randomized design was used with three factors of (1) mycorrhizal fungi with three variations (*G.etunicatum*, *G.intraradices* and nonmycorrhizal), (2) nutrient solution with three concentrations of micronutrients (C_0 , $C_{0.5}$ and C_1), and (3) time of leachate collection, 45, 65 and 85 DAS with three replications per treatment. Analysis of variance and mean comparison by Duncan's Multiple Range Test were carried out using MSTATC software.

Results and Discussion

Mycorrhizal roots produced considerably higher (p<0.05) siderophore (13.7 μ mol cm⁻³ h⁻¹) than nonmycorrhizal roots (2.0 μ mol cm⁻³ h⁻¹). Sorghum plants inoculated with either GE or GI were not different in siderophore production (Table 1). Obviously, nonmycorrhizal roots secrete only phytosiderophore but mycorrhizal roots can produce both siderophore and phytosiderophore. Enhancement of siderophore and/or phytosiderophore per unit volume of root in mycorrhizal plants suggests that AM fungi may secrete siderophore by themselves and/or induce plant root to produce more phytosiderophore. Arbuscular mycorrhizal fungi are reported to enhance Fe-uptake rates of associated host plants, which can be taken as an indication that mycorrhizal siderophores of a yet unknown structure may be involved (Haselwandter, 2008). An arbuscular mycorrhizal grass species, which showed greater Fe uptake than nonmycorrhizal controls, tested positively when bioassayed for hydroxymate siderophores (Haselwandter, 1995).

 Table 1. Main effect of fungal species on siderophore production and root colonization (RC)

Mycorrhiza	Sidero	% D C	
	µmol pot ⁻¹ h ⁻¹	µmol cm ⁻³ h ⁻¹	70 KC
NM	39.0a*	02.0c	00b
GE	39.7a	13.7b	43a
GI	39.7a	13.7b	43a
NTN C	1. 1 OF CI		

NM, non-mycorrhizal; GE, Glomus etunicatum; GI, Glomus intraradices.

*Means in each column followed by same letter are not significantly different (p<0.05)

Siderophore production per total volume of leachate obtained from each pot $(39-39.7 \ \mu mol \ pot^{-1} \ h^{-1})$ was not significantly different in mycorrhizal and non-mycorrhizal plants (Table 1). It was due to lower root volume in mycorrhizal plants (data not shown). Higher efficiency of mycorrhizal roots in nutrients uptake (Liao *et al.*, 2003; Aliasgharzad *et al.*, 2009) and carbon consumption by the symbiotic fungi (Landis and Fraser, 2008) may lead to restricted root development in AMF inoculated plants. Mycorrhizal colonization percentage was not significantly different between GE and GI inoculated roots (Table 1). This means that the two fungal species have nearly the same potential in producing siderophore or inducing plant root to enhance phytosiderophore production.

Siderophore and/or phytosiderophore concentrations $(11.3-25.0 \ \mu\text{mol}\ \text{cm}^{-3}\ \text{h}^{-1})$ have also been adversely affected by the micronutrient levels. The highest concentration was achieved at the C₀ level. There was no significant difference between C_{0.5} and C₁ in this respect (Table 2). Severe micronutrient starvation in C₀ has induced both root and AMF to secrete more siderophore. Sigel and Sigel (1998) pointed out that genes coding for siderophore in microorganisms, plants and animals are inducible and are usually expressed in response to Fe deficiency.

Table	2.	Main	effect	of	micronutri	ents	levels	on
		sidero	phore	I	production	ar	ıd	root
		coloni	zation ((RC))			

Mycorrhiza	Sidero	9/ DC	
	µmol pot ⁻¹ h ⁻¹	µmol cm ⁻³ h ⁻¹	- 70 KC
C ₀	40.3a*	25.0a	29a
C _{0.5}	38.0a	15.7b	25a
C1	40.3a	11.3b	26a

NM, non-mycorrhizal; GE, Glomus etunicatum; GI, Glomus intraradices.

*Means in each column followed by same letter are not significantly different (p<0.05).

Root colonization percentage was not affected (p<0.05) by micronutrients levels (Table 2). The maximum level of micronutrients used in this experiment, was the full strength of Rorison's nutrient solution. It has been reported that enough or higher levels of these nutrients may restrict root mycorrhizal colonization (Khaliel and Sohaibani, 1993; Liao *et al.*, 2003).

Interaction effect of micronutrient levels \times mycorrhizal inoculation on siderophore production indicates that in nonmycorrhizal roots the highest siderophore concentration occurred at C₀ level, but there was no difference between C_{0.5} and C₁ levels. In mycorrhizal roots there was no significant difference between micronutrient levels for siderophore production (Figure 1). Mycorrhizal fungi may alleviate nutrient deficiency in plant by increasing absorbing surfaces and therefore poor induction of siderophore genes.

With an exception for GE, siderophore production was significantly reduced from 45 to 65 DAS and thereafter was markedly increased toward 85 DAS in both mycorrhizal and non-mycorrhizal plants (Figure 2). Higher demand of plant for nutrients including Fe, in flowering stage (85 DAS) may be a reason for this phenomenon, which would lead to higher siderophore production. However, additional mechanisms may be involved in GE inoculated plants for this reduction.

Conclusion

Considering μ mol cm⁻³ h⁻¹ as a reasonable unit for siderophore production, which was obtained at harvest time (85 DAS), it can be concluded that mycorrhizal fungi enhance siderophore production by themselves and/or by inducing plant roots. Thus, micronutrients deficiencies (especially Fe) can stimulate siderophore production by both AM fungi and plant root. Overall, it seems that inoculation of sorghum plants with either GE or GI in micronutrient deficient condition, would help this plant to absorb enough micronutrients through chelate formation with siderophores. In case of fungi as symbionts of plants. the role of fungal capacity to synthesize siderophores, is not clear with regard to the establishment of the symbiosis. On the other hand, it is evident that such a fungal capacity may affect the Fe nutrition of associated host plants. Further study is necessary to distinguish between fungal derived siderophores and root derived phytosiderophores. A new approach involving a siderophore deficient mutant of a mycorrhizal fungus, which is already known to produce siderophore could be applied for an evaluation of the siderophore biosynthesis importance of for the establishment of the symbiosis with host plant.



Figure 1. Effect of mycorrhizal fungi on siderophore production in variable regimes of micronutrients





Acknowledgment

The financial support provided by the Faculty of Agriculture, University of Tabriz is gratefully acknowledged.

References

- Aliasgharzad, N., N. Saleh Rastin, H. Towfighi and A. Alizadeh. 2001. Occurrence of arbuscular mycorrhizal fungi in saline soils of the Tabriz Plain of Iran in relation to some physical and chemical properties of soil. *Mycorrhiza* 11: 119-122.
- Aliasgharzad, N., S.A. Bolandnazar, M.R. Neyshabouri and N. Chaparzadeh. 2009. Impact of soil sterilization and irrigation intervals on P and K acquisition by mycorrhizal onion (*Allium cepa L.*). *Biologia* 64: 512-515.
- Angle, J.S., A. Heggo and R.L. Chaney. 1989. Soil pH, rhizobia, vesicular-arbuscular mycorrhiza inoculation effect on growth and heavy metal uptake of alfalfa (*Medicago Sativa* L.). *Biology and Fertility of Soils* 8: 61-65.
- Bakkaus, E., R.N. Collins, J.L. Morel and B. Gouget. 2006. Anion exchange liquid chromatography inductively coupled plasma- mass spectrometry of the Co²⁺, Cu²⁺, Fe³⁺ and Ni²⁺ complexes of mugineic and deoxymugineic acid. *Journal of Chromatography* 1129: 208 -215.
- Buyer, J. S., M.G. Kratzke and L.J. Sikora. 1993. A method for detection of pseudobactin, the siderophore prodaced by a plant-growth-promoting *Psedomonas* strain, in the barley rhizosphere. *Applied Environmental Microbiology* 59: 677-681.
- Cairney. J. W. G. and A.E. Ashford. 1989. Reducing activity at the root surface in *Eucaliptus Pilularris-Pisolithus tinctorius* ectomycorrhizas. *Plant Phytopathology* 16: 99-105.
- Cakmak, I., K.Y. Gülüt, H. Marschner and R.D. Graham. 1994. Effect of zinc and iron deficiency on phytosiderophore release in wheat genotypes differing in zinc efficiency. *Journal of Plant Nutrition* 17:1-17.
- Caris, C., W. Hördt, H.J. Hawkins, V. Römheld and E. George. 1998. Studies of iron transport by arbuscular mycorrhizal hyphae from soil to peanut and sorghum plants. *Mycorrhiza* 8: 35-39.
- Clark, R. B. and S.K. Zeto. 1996. Mineral acquisition by mycorrhizal maize grown on acid and alkaline Soil. *Soil Biology and Biochemistry* 28: 1495-1503.
- Cress, W.A., G.V. Johnson and L.L. Barton. 1986. The role of endomycorrhizal fungi in iron uptake by *Hilaria jamesii* L. *Journal of Plant Nutrition* 9: 547-556.
- Gerdemann, J.W. and T.H. Nicolson. 1963. Spores of edomycorrhizal *Endogone* species extracted from soil by wet sieving and decanting. *Transactions of British Mycological Society* 46: 235-244.
- Giovanetti, M. and B. Mosse. 1980. An evaluation of techniques for measuring vesicular-arbuscular infections in roots. *New Phytologist* 84: 489-500.

- Haselwandter, K. 1995. Mycorrhizal fungi: Siderophore production. *Critical Reviews in Biotechnology* 15: 287-291.
- Haselwandter, K. 2008. Structure and function of siderophores produced by mycorrhizal fungi. *Mineralogical Magazine* 72:61-64.
- Khaliel A.S. and S.A. Sohaibani. 1993. Mycorrhizal corn growth in response to different iron levels. *Journal of King Saudi University* 6:5-11.
- Koa, B. J., A.C. Chang, A.L. Page and D.C. Adriano. 2001. Organic acids in root exudates and their effects on plant uptake of biosolids-borne metals. Savanah River Ecology Lab, University of Georgia Aiken, Sc 29802 Department of Environmental Sciences, University of California, Riverside, CA 92521.
- Landis F.C. and L.H. Fraser. 2008. A new model of carbon and phosphorus transfers in arbuscular mycorrhizas. *New Phytologist* 177: 466–479.
- Lee, Y.J. and E. George. 2005. Contributions of mycorrhizal hyphae to the uptake of metal cations by cucumber plants at two levels of phosphorus supply. *Plant and Soil* 278, 361-370.
- Leyval, C. and C.P.P. Reid. 1991. Utilization of microbial sideropheres by mycorrhizal and non-mycorrhizal pine roots. *New Phytologist* 119: 93-98.
- Liao, J.P., X.G. Lin, Z.H. Cao, Y.Q. Shi and M.H. Wong. 2003. Interaction between arbuscular mycorrhiza and heavy metals under sand culture experiment. *Chemosphere* 50, 847-853.

- Liu, A., C. Hamel, R.I. Hamilton, B.L. Ma and D.L. Smith. 2000. Acquisition of Cu, Zn, Mn and Fe by mycorrhizal maize (*Zea mays L.*) growing in soil at different P and micronutrient levels. *Mycorrhiza* 9: 331-336.
- Merryweather, J.W. and A.H. Fitter. 1991. Techniques in Arbuscular Mycorrhiza Research. York Mycorrhiza Research Group. UK.
- Milagres, A.M.F., A. Machuca and D. Napoleao. 1999. Detection of siderophore production from several fungi and bacteria by a modification of chrome azurol S (CAS) agar plate assay. *Journal of Microbiological Methods* 37: 1-6.
- Romheld, V.1987. Different strategies for iron acquisition in higher plants. *Physiological Plantarum* 90:231-234
- Schreiner, P.R., 2007. Effect of native and non native arbuscular mycorrhizal fungi on growth and nutrient uptake of 'Pinot noir' (*Vitis vinifera* L.) in two soils with contrasting levels of phosphorus. *Applied Soil Ecology* 36, 205-215.
- Schwyn, B. and I.B. Neilands. 1984. Universal chemical assay for the detection and determination of siderophores. *Analytical Biochemistry* 160: 47-56.
- Sigel A. and H. Sigel. 1998. Iron Transport and Storage in Microorganisms, Plants, and Animals. Vol.35, Marcel Dekker: New York.
- Silvia, G.A., S.F.B. Trufem, J.O.J. Saggin and L.C. Maia. 2005. Arbuscular mycorrhizal fungi in a semi-arid copper mining area in Brazil. *Mycorrhiza* 15, 47-53.
- Zhang, F., V. Römheld and H. Marschner. 1991. Diurnal rhythm of release of phytosiderophores and uptake rate of zinc in iron deficient wheat. *Soil Science and Plant Nutrition* 37:671-678.