

INDIGENOUS MEDICINAL PLANTS USED BY LOCAL PEOPLE OF SHAHI, LOWER DIR, (KHYBER PAKHTOON KHWA) SOUTHERN HIMALAYAN REGIONS OF PAKISTAN

Gul Jan¹, Mir Ajab Khan¹, Habib Ahmad² and Farzana Gul¹

¹Department of Plant Sciences Quaid-i-Azam University, Islamabad.

²Botany Department, Hazara University, Mansehra, Pakistan

Correspondence: drguljan@yahoo.com

ABSTRACT

The present work is based on the results of research conducted on traditional uses of some important plants by the local people of Shahi, Lower Dir in southern Himalayan Mountains, Pakistan. The locals of the area have been using the medicinal plants for many day to day uses for various ailments and are dependent on the plants in their surroundings for food, health, medication and various cultural purposes. A total of 50 important plant species belonging to 35 families of wild herbs, shrubs and trees, were found to be used as medicinal plants.

Key Words: Indigenous medicinal plants, local uses, Southern Himalayan regions of Pakistan

INTRODUCTION

Prior to the merger with Pakistan in 1969, whole of the district Dir was a princely state ruled by Nawab Shah Jehan Khan and his ancestors. Lower Dir was declared as full fledged district of Malakand division on 13.08.1996. It is composed of two sub divisions i.e., Timergara and Jandool. Sub-division have been further divided into six tehsils; Samarbagh, Munda, Lal Qillah, Balambat, Timergara and Adezai. Miskaini and Shahi are the two villages of tehsils Samarbagh.

It is located from latitude 34°- 37' to 35°-07' north longitudes and from 71°-31' to 72°-14' east longitudes. It is bounded on the north by Upper Dir district, on the east by Upper Dir and Swat districts, on the south by Malakand protected area and on the west by Bajur agency and Afghanistan. The population of lower Dir district is 717.65 thousand in 1998. The total area of the district is 1583 square kilometers.

The topography of the district is dominated by the mountains and hills which are part of the branches of the southern Hindukush. The mountains ranges, in general run from north to south, with highest peaks in the upper northern part of the district where they reach more than 3000 meters in height. In the central part the height varies between 1800 to 2000 meters. In the south the height shows a rapid decrease where it is around 600 meters. The villagers devoted to agriculture, cattle-breeding and wood related activities have halved in number during past 50 years. The absence of industrial and tourist facilities, due to the geographical position and road conditions, prevented in fact the development of the area.

Pakistan has rich history on the folk use of plants. Afridi (1986) listed 67 medicinal plants from Khyber agency. Haq & Hussain (1993) reported local medicinal and other traditional uses of plants of Mansehra. Medicinal plants of Rawalpindi (Arshad and Akram, 1999), Kalat (Durrani and Hussain, 2003), Kurram (Gillani *et al.*, 2003) and Margalla (Shinwari and Khan, 1998) have also been investigated. Ethnobotanical studies have also been made in the various parts of Dir Kohistan valleys (Jan *et al.*, 2010).


The present study was planned keeping in view the following objectives.

- To establish a database of the plants of Southern Himalayan regions with special reference to their ethnobotanical uses.
- To collect valuable information which has been transmitted orally from ancestors and knowledgeable persons to record these information for wider circulation.
- To investigate the local medicinal uses of plants for further investigation by plant scientists, pharmacologists and phytochemists.
- To aware the local communities about the conservation strategies of these valuable natural resources.

MATERIALS AND METHODS

This research was conducted during 2008-2009 in Shahi valley. The plants were collected, dried and preserved for identification. They were identified with the help of available literature (Stewart 1972, Nasir and Ali, 1971-95).

The information about the medicinal uses of the plants was obtained from the local people through questionnaire. The outcome of the results were rechecked and compared with literature like that of Rubina (1998), Ali and Fefevre (1996) and Khalid (1995). The data was analyzed and indigenous knowledge was documented.



Map showing Shahi from where plants were collected.

RESULTS AND DISCUSSION

The present study provides information on the indigenous uses of 42 important ethno-botanically important plants belonging to 30 families. These ethno-botanically valued plants with their respective families, botanical name, local name, habit, part used and local uses were documented (Table 1).

Local people of Shahi are using the plants for various purposes i.e., medication, food, cosmetics, and fodder for the cattle. They have faith in these plants. The ratio of the women using allopathic medicines is negligible because they are directly dependent on plants for medication and other basic needs. The ethno-botanically important and other beneficial plants are quite useful for the basic health and hygiene of the local people. Local people are directly dependent on these plants for cure of different diseases, food, skin care, cosmetics and fodder for the cattle. About 80% population of the world depends on the traditional system of health care. These medicines have less side effects and man can get the herbs easily from nature (Ahmad, 2005). Most of the plants used by the local people are not conserved but are over exploited. It is therefore necessary to find the ways of promoting the local people towards conservation. The investigated area has a rich diversity of medicinal plants and provides a conducive habitat and ideal climatic conditions for their growth. It is suggested that local community should receive education on the identification of their indigenous medicinal plants.

Table 1. Common medicinal plants of Shahi valley, lower Dir, South Himalaya, Pakistan.

S.No.	Botanical Name	Family	Part use	Habit	Medicinal uses
1	<i>Acorus calamus</i> L.	Aceraceae	Rhizome	H	Dysnetery and chronic diarrhea
2	<i>Artemesia maritime</i> L.	Asteraceae	Leaves	S	Intestinal worm killer
3	<i>Achyranthus aspera</i> L.	Amaranthaceae	Roots	H	Stomachache, urine formation and cough
4	<i>Astragalus macropterus</i> DC	Fabaceae	Leaves	H	Stomachic
5	<i>Aquilegia pubiflora</i> Wall. ex Royle	Ranunculaceae	Seed	H	Astringents and helpful to women in child birth

S.No.	Botanical Name	Family	Part use	Habit	Medicinal uses
6	<i>Atriplex crossifolia</i> C.A. Mey	Chenopodiaceae	Leaves	S	Throat infection and yellow jaundice
7	<i>Berberis lycium</i> Royle	Berberidaceae	Roots	S	Jaundice powder for rheumatism & backache
8	<i>Bergenia himalaica</i> Boiss	Saxifragaceae	Root	H	Headache and wound healing
9	<i>Betula utilis</i> D. Don	Betulaceae	Bark	T	Tonic, leprosy and earache
10	<i>Buxus wallichiana</i> Bill	Buxaceae	Leaves	S	Antirheumatic, Diaphoretic, purgative, poisonous and febrifuge
11	<i>Bunium persicum</i> B. Fedtsch	Apiaceae	Fruit	H	Carminative, stomachic and stimulant
12	<i>Capparus spinosa</i> L.	Capparidaceae	Floral buds	S	Improve taste powder
13	<i>Calotropis procera</i> (Wild).	Asclepiadaceae	Leaves	H	Asthma
14	<i>Chenopodium ambrosioides</i> L.	Chenopodiaceae	Whole plant	H	Intestinal worm killer
15	<i>Corydalis adiantifolia</i> H. & T.	Fumiraceae	Root	H	Eye diseases and improve eye sight
16	<i>Datura stramonium</i> L.	Solanaceae	Seeds	H	Anticholinergic and sedative
17	<i>Ephedra gerardiana</i> Wall ex Stapf	Ephedraceae	Dried twigs	S	Asthma and cough
18	<i>Ficus carica</i> Forssk.	Moraceae	Fruit	T	Laxative, demulcent used in constipation, piles and urinary bladder problems
19	<i>Fragaria vesca</i> Lindle. ex HK	Rosaceae	Leaves	H	Mildly astringent and diuretic
20	<i>Galium boreale</i> L.	Rubiaceae	Flower	H	Blood purifier and diuretic
21	<i>Hedera nepalensis</i> K. Koch.	Araliaceae	Leaves	H	Antidiabetic
22	<i>Hyoscyamus niger</i> L.	Solanaceae	Seeds	H	Seeds are chewed for relieving toothache
23	<i>Hypericum perforatum</i> L.	Hypericaceae	Leaves	H	Diuretic
24	<i>Inula racemosa</i> HK	Asteraceae	Root	H	Tonic, stomachic and carminative
25	<i>Lonicera perichyenum</i> L.	Caprifoliaceae	Leaves and flower	S	Wound healing, laxative and cough
26	<i>Lactuca decipiens</i> (H.& T) Clarke	Asteraceae	Seeds	H	Headache and tonic
27	<i>Malva neglecta</i> Wall.	Malvaceae	Leaves	H	Leaf decoction is antispasmodic
28	<i>Mentha sylvestris</i> L.	Lamiaceae	Leaves	H	Carminative, stimulant and rheumatism
29	<i>Morus alba</i> L.	Moraceae	Fruit	T	Fruit is laxative and edible
30	<i>Onosma bracteatum</i> L.	Boraginaceae	Root	H	Fever, heart trouble and tonic
31	<i>Origanum vulgare</i> L.	Lamiaceae	Leaves	H	Toothache and earache
32	<i>Prunus persica</i> (L.) Stokes	Rosaceae	Fruit	T	Jaundice, sedative, expectorant and tonic
33	<i>Plantago major</i> L.	Plantaginaceae	Seed	H	Demulcent, diuretic, constipation and piles
34	<i>Pistacio integerrima</i> J.L. Stewart ex Brand.	Anacardiaceae	Fruit	T	Jaundice
35	<i>Ranunculus aquatilis</i> L.	Ranunculaceae	Whole plant	H	Asthma and periodic fever
36	<i>Sonchus asper</i> (L.) Hill	Asteraceae	Whole plant	H	Tonic

S.No.	Botanical Name	Family	Part use	Habit	Medicinal uses
37	<i>Sassurea lappa</i> (Dene.) Sch.	Asteraceae	Root	H	Bronchial asthma, stimulant, cough and dyspepsia
38	<i>Salix acmophylla</i> Boiss.	Salicaceae	Bark	T	Tonic and febrifuge
39	<i>Solanum surratense</i> Burm.f.	Solanaceae	Whole plant	H	Expectorant, antiasthmatic, antigonorrhoea stomachache, cough and fever
40	<i>Taraxicum ooficinale</i> L.	Asteraceae	Root	H	Laxative
41	<i>Thymus linearis</i> L.	Lamiaceae	Dried leaves	H	Whooping cough, asthma, expel round worm and antiseptic
42	<i>Teucrium stocksianum</i> Boiss	Lamiaceae	Whole plant		Cooling agent, tonic and against hepatitis
43	<i>Trachyspermum ammi</i> (L.) Sprague	Apiaceae	Seeds	H	Laxative, aphrodisiac, stimulant and carminative
44	<i>Urtica dioica</i> L.	Urticaceae	Leaves and stem	H	Anthelmintic, diuretic and jaundice
45	<i>Verbascum Thapsus</i> L.	Scrophulariaceae	Leaves and Flower	H	Fever, astringent, bleeding of lungs and narcotic
46	<i>Verbena officinale</i> L.	Valerianaceae	Roots	H	Jaundice and kills and expels worms in intestine
47	<i>Veronica ciboria</i> L.	Scrophulariaceae	Roots	H	Expectorant
48	<i>Woodfordia fruticosa</i> (L.) Kurz.	Lythraceae	Flower	S	Dried flowers are used as astringent, stimulant and used in liver complains
49	<i>Xanthium strumarium</i> L.	Asteraceae	Leaves	S	Skin diseases
50	<i>Ziziphus jujube</i> Mill.	Rhamnaceae	Fruits	T	Expectorant, emollient and blood purifier

REFERENCES

- Afridi, S.K. (1986). *Medicinal plants of Khyber Agency*. M.Sc. Thesis. Department of Botany, University of Peshawar.
- Ahmad H (2005). *Issues Regarding Medicinal Plants of Pakistan*. Udyana Today, 6(3): pp 6-7.
- Ali, A. and J.L. Fefevre (1996). "Indigenous knowledge of plants-A case study in Chitral"-*Proceedings of Ethnobotany workshop*, September 16-24, NARC Islamabad, pp. 136-151.
- Arshad, M and S. Akram (1999). Medicinal plants of University of Arid Agriculture, Rawalpindi *Hamdard Medicus*, 40: 46-50.
- Durrani, M.J. and F. Hussain (2003). Ethnoecological profile of plants of Harboi rangeland, Kalat, Pakistan. *Int. J. Biol. and Biotech.*, 2: 15-22
- Gilani, S. S., S. Q. Abbas. Z. K. Shinwari, F. Hussain, and K. Nargis (2003). Ethnobotanical studies of Kurram Agency, Pakistan through rural community participation. *Pak. J. Biol. Sci.*, 6: 1368-1375
- Haq, I.U and M. Hussain (1993). Medicinal plants of Mansehra District. *Hamdard Medicus*, 34: 63-99.
- Jan, G., M.A. Khan, F. Gull, M. Ahmad, M. Jan and M. Zafar (2010). Ethnobotanical studies of common weeds of Dir, Kohistan valley, Khyber Pakhtoonkhwa, Pakistan. *Pak. J. Weed Sci. Res.* 16 (1): 81 -88.
- Khalid, S. (1995). "Plant in danger", *Fifth National Conference of Plant Scientists*, March 28-30, NARC Islamabad.
- Nasir, E. and S.I. Ali (1971-95). "*Flora of west Pakistan*", Department of Botany, University of Karachi, Karachi.
- Rubina, A.R. (1998). "*Need of conservation for endangered Medicinal Plants*", *Proceedings of wild medicinal plant resources of Northern Pakistan*, May 11-12, PFI, Peshawar, Pakistan.
- Shinwari, M. I. and M. A. Khan (1998). *Ethnobotany of Margalla Hills*. National Park, Islamabad. Dept. of Biological Sciences, Q.A.U.
- Stewart, R.R. (1972). An Annotated Catalogue of the Vascular Plants of West Pakistan and Kashmir. Fakhri Printing Press, Karachi. XVIII + 1028 pp.

(Accepted for publication October 2010)