

## IN VITRO MAST CELL STABILIZATION ACTIVITY OF *VIOLA ODORATA* L.

G.P. Choudhary, S.C. Chaturvedi and S. Bharti

School of Pharmacy, Ring road, Devi Ahilya University, Indore-452017, INDIA

Email address: choudhary.gp@gmail.com

---

### ABSTRACT

The influence of the ethanolic extract of the flower of *Viola odorata* L. on degranulation of rat peritoneal mast cell induced by compound 48/80 and Egg albumin was studied. The inhibitory effect of the extract was significant in immunologically induced degranulation of mast cells.

**Keywords:** Flavonoids, Kitotifen, Mast cell degranulation, *Viola odorata*.

---

### INTRODUCTION

Mast cells and basophils play a central role in inflammatory and immediate allergic reactions. On stimulation, they are able to release potent inflammatory mediators, such as histamine, proteases, chemotactic factors, cytokines and metabolites of arachidonic acid that act on the vasculature, smooth muscle, connective tissue, mucous glands and inflammatory cells. Mast cells settle in connective tissues and usually do not circulate in the blood stream. Basophils are the smallest circulating granulocytes with relatively the least known function. They arise in the bone marrow, and following maturation and differentiation, are released into the blood circulation. Adequately stimulated basophils may settle in the tissues. There are two categories of inflammatory (anaphylactic) mediators in mast cells and basophils. Preformed mediators, stored in secretory granules and secreted upon cell activation, include a biogenic amine, typically histamine, proteoglycans, either heparin, over-sulphate chondroitin sulphates or both, and a spectrum of neutral proteases. Released histamine acts at H1, H2 and H3 receptors on cells and tissues, and is rapidly metabolized extracellularly. The proteoglycan imparts the metachromatic staining characteristic of mast cells when exposed to certain basic dyes such as toluidine blue. It has two functions, (a) may package histamine and basic proteins into secretory granules, and in mast cells and (b) appears to regulate the stability of the protease called tryptase. Neutral proteases, which account for the vast majority of the granule protein, serve as markers of mast cells found in serosal, mucosal and brain region. Newly generated mediators, often absent in the resting mast cells, are typically produced during IgE-mediated activation, and consist of arachidonic acid metabolites, principally leukotriene C4 (LTC4) and prostaglandin D2 (PGD2) and cytokines. Of particular interest in humans is the production of tumour necrosis factor, IL-4, IL-5 and IL-6. In the cytoplasm of both mastocytes and macrophages are special lipid bodies, where metabolism of arachidonic acid occur and their products, including leukotrienes, may be stored (Forman, 1993).

*Viola odorata* (Violaceae), It was well known herb in India for its medicinal virtues and has been in use since ancient times. It was used in the treatment of several diseases both in the Ayurvedic and the unani systems of medicines. It was found to be effective against induced inflammation, similar to ipecac extract. The flowers are useful in the treatment of cold and cough. The flowers and leaves of plant are made into a syrup used for respiratory ailments associated with congestion, coughing and sore throat. Tea made from the entire plant is used to treat digestive disorders. It acts as a medicinal remedy for headache, body pains and as a sedative. It is used externally, the fresh crushed leaves reduce swelling and soothe irritation. The whole plant is anti-inflammatory, diaphoretic, diuretic, emollient, expectorant and laxative. It is taken internally in the treatment of bronchitis, respiratory catarrh, coughs, asthma and cancer of the breast, lungs or digestive tract. The herb shows antibacterial activity and is effective in treatment of eczema. Flowers contain traces of volatile oil, several peculiar colouring matter and violaquercitrin a flavonoid; the drug also contains glucoside, methyl salicylic ester. Violin is an emetic principle present in all parts of the plant (Pillai *et al.*, 1992, Nadkarni and Nadkarni, 1982).

### MATERIALS AND METHODS

#### *Plant material*

The dried flower powder of *Viola odorata* was purchased from Dravid Herbs World, Pondicherry, India. Egg albumin was purchased from Hi-Media Lab., Mumbai and compound 48/80 was purchased from sigma chemicals company, USA. Kitotifen (Ketovent) was purchased from Intas Pharm. Ltd., India.

### Extraction

*Viola odorata* flower powder extracted with 95% ethanol in a soxhlet extractor. The extract was concentrated under reduced pressure at a temperature below 50°C to yield a syrupy mass (Yield -8.33%), which was used for the present investigation

### Preliminary phytochemical investigation

Preliminary phytochemical analysis shows the presence of glycosides and flavonoids (Kokate, 1991).

### Animals

Animals-Male Wister rats (200-250g) were obtained from the experimental animal house, School of life science, Devi Ahilya University, Indore. They were maintained under standard housing condition. The animals were given standard laboratory feed and water ad libitum. The study was cleared by Animal ethics committee. All the animals received humane care according to criteria outlined in the guide for the care and use of laboratory animals prepared by the national academy of the sciences and published by national institute of health.

### Degranulation studies

Sensitized mast cells were obtained from animals sensitized with egg albumin. The doses being given on the 1<sup>st</sup>, 3<sup>rd</sup> and 5<sup>th</sup> day. The sensitized mast cells were degranulated using egg albumin (1mg/ml) on the 10<sup>th</sup> day of sensitization. The normal mast cells were degranulated using compound 48/80 (100mcg/ml). The cell suspension of mast cells was treated as follows.

To 0.1 ml of the peritoneal mast cell suspension, 0.1ml of the test agent in the saline was added and incubated in a constant temperature water bath (37°C) for 15 minutes. Then 0.1 ml of degranulating agent (Egg albumin 1 mg/ml or compound 48/80 100mcg/ml) was added and further incubated for a period of 10 minutes. The cell were then stained with 0.1% toluidine blue for 5-10 min and the tissue was then washed in acetone and then xylene (2 changes each) for 5 min each wash.

The stained cells were viewed through a digital light microscope at 100x magnification and 100 mast cells were counted. The number of intact and fragmented or disrupted mast cells was noted. A mast cell was considered disrupted if four or five granules were found around the mast cells. The numbers of fragmented or disrupted mast cells as well as of the intact mast cells were counted (Nair *et al.*, 1994, Ghosai and Bhide, 1981, Norton, 1954, Peachell, 2005).

### Analysis

Values were expressed as mean  $\pm$  SE. The values were statistically analyzed using one-way Analysis of Variance (ANOVA) followed by Tukey's multiple comparison test. The analysis was carried out using Graph Pad Prism software V.4.

**Table I. Effect of *Viola odorata* extract on egg albumin induced mast cell degranulation in rats.**

S.No.	Treatment	Dose mcg/ml	Number of mast cell	Percent Inhibition
1.	Control	-	7 $\pm$ 2	-
2.	Ketotifen	10	82 $\pm$ 4	78.22*
3.	VO extract	05	31 $\pm$ 3	18.85*
4.	VO extract	10	47 $\pm$ 5	38.96*
5.	VO extract	20	60 $\pm$ 6	59.97*
6.	VO extract	40	75 $\pm$ 4	72.65*

Values are mean $\pm$ S.E., \*P<0.001 when compared with control.

## RESULTS

### Egg albumin induced degranulation studies

Kitotifen as a reference standard was found to inhibit degranulation to an extent of 78.22. *Viola odorata* extract at concentration 5,10,20 and 40 mcg/ml produced dose related inhibition of 18.85,38.96,59.97 and 72.65 respectively (Table 1).

**Compound 48/80 induced degranulation studies**

Ketotifen at a concentration of 10mcg/ml was found to inhibit degranulation to an extent of 77.52. *Viola odorata* extract at concentration of 5,10,20 and 40 mcg/ml showed reduction in degranulation of mast cell to 19.36,35.08,58.52 and 40.28 respectively (Table 2).

**Table 2. Effect of *Viola odorata* extract on compound 48/80 induced mast cell degranulation in rats.**

S.No.	Treatment	Dose mcg/ml	Number of mast cell	Percent Inhibition
1.	Control	-	8±2	-
2.	Ketotifen	10	80±4	77.52*
3.	VO extract	05	32±4	19.36*
4.	VO extract	10	44±3	35.08*
5.	VO extract	20	60±4	58.52*
6.	VO extract	40	44±3	40.28*

Values are mean ± S.E., \*P < 0.001 when compared with control.

**DISCUSSION**

The mast cells have a crucial role in the development of many physiological changes during anaphylactic and allergic responses. Immunoglobulin- E antibodies bind to receptors on the surface of mast cell. Allergen-IgE interaction on mast cell leads to the release of histamine, heparin, proteases and other mediators and the synthesis and secretion of leukotrienes and prostaglandins. These products result in bronchoconstriction, changes in blood vessel tone, increased vascular permeability and myriad other proinflammatory effects (Subramoniam *et al.*, 1999). The functions of mast cells can be manipulated for therapeutic ends by regulating their function with appropriate drugs. Plant origin constituents may influence differentiation into mast cells, chemical composition and or architecture of mast cell surface membrane. It may influence the synthesis of IgE molecules or binding of IgE on mast cell surface. It is also possible, that the plant drug may reduce the life span of mast cells.

Extract of *Viola odorata* markedly protected the sensitized mast cells. However, the effect was less than that observed with the standard drug (Ketotifen) used. The pathological mechanism involved in Type-I allergy has been explained as the degranulation of mast cells and basophils, followed by the release of mediators such as histamine, leukotrienes and prostaglandins from these cells (Crossland, 1980). The degranulation of mast cells occurs in response to the immunological stimuli in which the antigen-antibody interaction on the cell surface predominates. The present investigation indicates that the extract of *Viola odorata* is active against Type-I allergic conditions because of their ability to inhibit the release of mediators from mast cells and basophils and thus influences the course of the disease. The preliminary phytochemical tests showed the presence of flavonoids in the *Viola odorata* ethanolic extract.

Mast cells after degranulation shows demonstrated that transgranulation occurs between mast cells and fibroblasts with mast cells apparently transferring their granules to the cytoplasm of fibroblasts or to the mesothelium. It has been reported that mast cell granules are internalized in fibroblasts 1-3 h after C48/80 injection (Norrby and Enestrom, 1984). In the mast cell the extruded granules might be degraded by the extracellular, as the initial compact morphological appearance of the discharged granules is gradually lost, and the granule contents are discharged (Takayama *et al.*, 1994). Mast cells are well known for their close appositions to the nervous system, such as to the enteric nerves of the intestine, vagus nerves of the mesentery in the rat, and trigeminal sensory fibers in the rat dura mater. Electrical trigeminal stimulation promotes mast cell secretion and degranulation in the dura mater and tongue, and this activation of mast cells by neurogenic mechanisms appears to be important in the development of neurogenic inflammation (Dimitriadou *et al.*, 1991).

The present investigation indicates the ethanolic extract of *Viola odorata* is active in the Type-I allergic conditions because of their ability to inhibit the release of mediators from mast cells and thus influence the course of the disease by preventing the harmful effects of the released mediators. The preliminary phytochemical tests showed the presence of flavonoids in the ethanolic extract. Plant flavonoids are known to inhibit basophil histamine release and neutrophil beta-glucuronidase release, and thereby possess in-vivo antiallergic activity (Pathak *et al.*, 1991). The flavonoids also inhibited the histamine release induced by 48/80. Plants containing flavonoids have been reported to possess antihistaminic, antiallergic and mast cell degranulation properties (Tripathi *et al.*, 1979, Havsteen, 1983).

Based on these results, it could be suggested that *Viola odorata* stabilizes mast cells in the rat. As mast cells play a major role in Type I hypersensitivity-mediated diseases like allergic asthma and rhinitis, studies are under

way to evaluate the efficacy of *Viola odorata* due to its mast stabilization property in these animal allergic models.

## REFERENCES

- Crossland J. (1980). *Pharmacological Aspects of Allergy*. Lewis Pharmacology. Churchill Livingstone, New York, p. 393.
- Dimitriadou, V., M.G. Buzzi, M.A. Moskowitz and T.C. Theoharides (1991). Trigeminal sensory fiber stimulation induces morphological changes reflecting secretion in rat dura mater mast cells. *Neuroscience*, 44: 97-112.
- Foreman, J.C. (1993). *Non-immunological stimuli of mast cells and basophil leucocytes*. *Immunopharmacology of Mast cells and Basophils*. London: Academic Press. p.57-69.
- Ghoosi, R.B. and M.B. Bhide (1981). ,*Asp.Allergy Appl.Immunol*;1981,15,53.
- Havsteen, B. (1983). Flavonoids: a class of natural products of high pharmacological potency. *Biochem. Pharmacol.*, 32: 1141–1148.
- Kokate, C.K., (1991). *Practical Pharmacognosy*, 4<sup>th</sup> edn., Vallabh Prakashan, New Delhi, p.110.
- Nadkarni, A.K. and K.M. Nadkarni (1982). *Indian Materia Medica*, Vol.I, Popular Prakashan, Bombay, p.1274-1275.
- Nair, A.M., C.P. Tamhankar and M.N. Saraf (1994). *Indian drugs*, 32(6),1994,277-282.
- Norrby K, and S Enestrom (1984). Cellular and extracellular changes following mast-cell secretion in avascular rat mesentery. *Cell Tissue Res.*, 235: 339-345.
- Norton S. (1954). Quantitative determination of mast cell fragmentation by compound 48/80. *Br. J. Pharmacol.*, 9: 494-497.
- Pathak, D., K. Pathak, A. K.. Singla. (1991). ,Flavonoids as medicinal agents — recent advances. *Fitoterapia*, 62: 371–389.
- Peachell P. (2005). Targeting the mast cell in asthma. *Curr. Opin. Pharmacol.*, 5: 251-256.
- Pillai et al. (1992). *Indian Journal of Natural Prod.*, 8(1): 16.
- Subramoniam, A., D.A. Evans, R. Valsaraj, S. Rajasekharan and P. Pushpangadan (1999). Inhibition of antigen-induced degranulation of sensitized mast cells by *Trichopus zeylanicus* in mice and rats. *J. Ethnopharmacol.* 68: 137–143.
- Takayama, I., Y. Fujii, S. Ohno and M.A. Fujino (1994). Ultrastructural study of mast cells stimulated with C48/80 as revealed by quick-freezing method. *Virchows Arch.*, 424: 287-294.
- Tripathi, R.M., P.C. Sen and P.K. Das (1979). Studies on the mechanism of action of *Albizia lebbeck*, an Indian indigenous drug used in the treatment of atopic allergy. *J. Ethnopharmacol.*, 1: 385–396.

(Accepted for publication September 2008)