

EFFECT OF STAGE OF MATURITY AND CULTIVARS ON THE DIGESTIBILITY OF MAIZE FODDER

Rafia Firdous, Abrar H. Gilani; A.R. Barque & Muhammad Akram

Departments of Animal Nutrition & Poultry Husbandry, University of Agriculture, Faisalabad

A study was conducted on four maize cultivars to determine the dry matter and fibre digestibility as influenced by advancing plant age. Samples of maize cultivars Akbar, Naeelum, U.M-81 and I.Z-31 were harvested at weekly intervals/ various growth stages. The samples of morphological fractions such as leaf and stem were also collected at various growth stages. Whole mixed fodder and different fractions of maize plant were analyzed for their chemical composition and in vitro digestibility. The results showed that in vitro dry matter digestibility of whole maize plant, leaf and stem decreased significantly with advancing stage of maturity. Digestibility of neutral detergent fibre, acid detergent fibre, hemicellulose and cellulose decreased significantly in all plant parts with advancing plant age/growth stages. Maximum values for the digestibility of dry matter and various cell wall constituents were observed in leaf, followed by whole plant and stem fractions. Cultivars were observed to have significant effect on dry matter digestibility and digestibility of neutral detergent fibre, acid detergent fibre and cellulose in all plant fractions. The results indicated that digestibility of maize fodder was affected by both stage of maturity as well as cultivars, however, maturity had a greater effect on digestibility in all plant fractions than did cultivars. Dry matter contents were found to be significantly and negatively correlated with dry matter digestibility of whole plant and its leaf and stem fractions. Based on correlations, regression equations were computed to predict in vitro dry matter digestibility.

Key words: cultivars, dry matter digestibility, maize, morphological fractions, stage of maturity

INTRODUCTION

The value of a good quality fodder for animal production depends upon the nutrient concentration as well as fodder intake by animals. The quality of fodder is evaluated in terms of palatability, voluntary intake, digestibility and utilization of various nutrients. Poor digestibility and lower voluntary intake are always associated with relatively higher lignin contents. Lignin is a chemical fraction of the cell wall most frequently associated with digestibility of forages by ruminants (Van Soest, 1987). Various mechanisms have been suggested by which lignin inhibits cell wall digestion e.g. encrustation, toxicity to digesting microbes and lignin-polysaccharide complexes. Plant polysaccharides are not simply encrusted by lignin but are probably covalently bonded to it (Bolker, 1963). Maize has for centuries been used as a forage crop in Pakistan. Generally, the maize fodder along with cobs at milking stage is cut and fed to animals. It provides adequate energy and protein for growth and milk production (Chaudhry, 1983). Maize has always higher in vitro dry matter digestibility as compared to other fodder crops and is considered as the most suitable summer fodder crop (Lloveras, 1990). Most of the work conducted on forage evaluation is on forages grown in temperate regions and this information as such may not be applicable to fodders grown in tropical or sub-tropical climate. The tropical forages are generally higher in lignin contents than those in the temperate regions. Animal production in tropical countries is, therefore, handicapped by the low quality of the forages (Van Soest, 1987). Comprehensive nutritional evaluation of maize fodder has not been undertaken in Pakistan. An attempt was therefore, made to determine the

nutritive value of local maize fodder with particular reference to morphological fractions, cultivar differences and the effect of advancing harvest stages.

MATERIALS AND METHODS

The study was conducted on four approved maize cultivars, Akbar, Neelum, U.M-81 and I.Z-31. All the maize cultivars were sown from March to June, 1991 in the experimental fields of the University of Agriculture, Faisalabad. The crop was sown in 3 replicates using randomized complete block design. Urea at the rate of 125 kg/hectare was applied as fertilizer. Six irrigations of canal water were given during the 14 weeks experimental period. The fodder was harvested for drawing samples at weekly intervals. The morphological fractions of the plant such as leaf and stem were also collected at various growth stages. The leaves (blade + sheath) were separated manually from the stem. The whole maize plant and its leaf and stem fractions were chopped into 2 to 3 cm pieces and dried at 60°C (AOAC, 1984). Dried samples were ground, passed through 1 mm screen and saved in labeled air tight containers for analysis. The structural components such as neutral detergent fibre (NDF), acid detergent fibre (ADF), hemicellulose, cellulose, lignin, cutin and silica were determined by using detergent system (Van Soest and Robertson, 1985).

The in vitro digestion technique developed by Troelsen (1971) was used for measuring the digestibility of dry matter and various fibrous fractions such as NDF, ADF, hemicellulose and cellulose of the fodder samples. Fodder samples (3g) were weighed in 250 ml Erlenmeyer flasks fitted with rubber

stoppers. The flasks and samples were placed in an incubator at $39 \pm 1^\circ\text{C}$ and 120 ml inoculum was added to each flask. The inoculum was prepared by mixing the strained rumen fluid with artificial sheep saliva (McDougall, 1947) as modified by Baumgardt *et al.* (1962). The artificial sheep saliva was heated and held at 39°C and charged with stream of carbon dioxide before the rumen fluid was added. The digestion time was 48 ± 1 hours. The digested samples were filtered and washed with boiling water. The residue was dried at 105°C and weighed. The above mentioned procedure was followed upto stage one only.

Statistical Analysis: The data on various parameters were subjected to statistical analysis using analysis of variance technique and Duncan's multiple range test was applied to compare treatment means (Steel and Torrie, 1984).

RESULTS AND DISCUSSION

In Vitro Digestibility: Average in vitro digestibility coefficients of DM and various structural components such as NDF, ADF, hemicellulose and cellulose of different fractions of maize plant at various growth stages are presented in Table 1. Table 2 shows the average digestibility coefficients of DM and various cell wall components of various cultivars of maize plant and its morphological fractions.

In Vitro Dry Matter Digestibility (IVDMD): The DM digestibility declined significantly with advancing age in whole maize plant and its leaf and stem fractions. The highest DM digestibility coefficients were recorded in case of leaf fraction (68.21 ± 0.59 to 77.04 ± 0.68). The DM digestibility of whole plant ranged from 65.69 ± 0.44 to 75.82 ± 0.73 , being higher than that of stem fraction of the plant (58.52 ± 0.32 to 69.35 ± 0.35). Azim *et al.* (1989) also reported a decline in dry matter digestibility (DMD) of whole maize plant and its fractions at two vegetative stages. They further reported that maximum DMD was found in leaves followed by whole mixed plant, middle and bottom portions of the stem. However, the values reported by them were lower than those observed in the present study.

Variations in DM digestibility due to cultivars were significant in whole plant and its leaf and stem fractions. Neelum cultivar had significantly higher DM digestibility in whole plant, leaf and stem fractions when compared to other cultivars, whereas significantly lower digestibility was observed in case of U.M-81 cultivar (Table 2). The higher DM digestibility of Neelum, as compared to other cultivars could be due to lower NDF, ADF and lignin concentrations in it. Hunt *et al.* (1993) also reported that rumen in situ (24 hr) and 2-stage in vitro DM digestibility of whole maize plant and stover samples was greater ($P < 0.01$) for maize hybrid 3377 than for 3389.

NDF: Significant differences existed among in vitro NDF

digestibility coefficients of whole maize plant and its morphological fractions with advancing growth stages. NDF digestibility coefficients of leaf fraction ranged from 70.24 ± 0.90 to 76.70 ± 0.78 , being higher than those of whole plant (64.54 ± 0.47 to 72.45 ± 0.65). However, the stem fraction had the lowest NDF digestibility (54.40 ± 0.67 to 62.34 ± 0.95). A similar trend was reported by Vona *et al.* (1984). They observed that digestibility of NDF by cattle and sheep declined ($P < 0.05$) with advancing maturity of the grasses. Cultivar effects on NDF digestibility varied in whole plant and its leaf and stem fractions. Neelum cultivar had significantly higher NDF digestibility in all plant fractions, whereas it was the lowest in U.M-81 (Table 2). It was probably due to the higher structural components of U.M-81 than that of Neelum cultivar. Hunt *et al.* (1993) also reported significant differences in digestibility of NDF between maize hybrids.

ADF: In vitro ADF digestibility declined significantly in whole maize plant and its leaf and stem fractions with advancing stages of growth. The leaf fraction of the plant was found to have the highest ADF digestibility (61.86 ± 0.95 to 67.71 ± 0.35) followed by whole plant (57.40 ± 0.79 to 67.93 ± 0.88). However, the lowest ADF digestibility (42.73 ± 1.33 to 54.11 ± 2.49) was observed in case of stem fraction. This variation was probably due to higher hemicellulose and cell contents and lower lignin concentration in the leaf fraction of the plant.

Significant variations also existed in the in vitro ADF digestibility of whole maize plant and its leaf and stem fractions with respect to various cultivars. ADF digestibility was found to be higher for Neelum cultivar in whole plant, leaf and stem fractions (Table 2). However, ADF digestibility was almost similar for Akbar, U.M-81 and I.Z-31 cultivars. It may be due to lower lignin concentration in Neelum, as compared to other cultivars. Digestibility of ADF was reported to be higher in maize hybrid pioneer 3377 than in 3389 (Hunt *et al.*, 1993).

Hemicellulose: In vitro hemicellulose digestibility declined significantly with advancing plant maturity in whole maize plant and its morphological fractions. In vitro hemicellulose digestibility coefficients were higher in leaves (82.27 ± 0.95 to 88.42 ± 0.66) than those in whole plant (73.26 ± 0.72 to 86.27 ± 0.48). Hemicellulose digestibility coefficients ranged from 67.71 ± 0.23 to 76.03 ± 0.55 , being the lowest in the stem fraction of the plant. The digestibility of hemicellulose has been reported to decline non-linearly with advancing maturity stage (Sanderson *et al.*, 1989). Hemicellulose was digested 80% faster than cellulose. Singh and Narang (1990) also noted a slightly higher hemicellulose digestibility than that of cellulose in different forages. Findings of this study are in line with those of the above reported work.

Variations in hemicellulose digestibility due to various cultivars were observed only in case of leaf fraction of the plant.

Digestibility of maize fodder

Table 1. Average in vitro digestibility coefficients of OM, NDF, ADF, hemicellulose and cellulose of whole maize plant and its morphological fractions at different growth stages

| Description | Growth stages (age in weeks) | | | | |
|---------------|------------------------------|-----------------------|--------------------|----------------------|------------------|
| | Seedling (4th) | Early growth (5th) | Flowering (9th) | Milk/dough (11th) | Mature (14th) |
| | Whole plant | | | | |
| IVDMD | 75.82±0.73a | 70.34±0.49b | 70.30±0.40c | 67.35±0.48d | 65.69±0.44e |
| NDF | 69.45±0.65a | 70.82±0.51b | 69.01 ±0.53c | 66.n±0.73d | 65.54 ±0.47e |
| ADF | 67.93±0.88a | 64.11±1.15b | 63.06±0.94c | 58.18±0.69d | 57.40±0.79d |
| Hemicellulose | 86.27 ±0.48a | 81.64±0.82b | 77.21 ±0.65c | 75.93±0.64c | 73.26±0.nd |
| Cellulose | 83.47±0.60a | 79.12±0.65b | n.94±0.73c | 65.77 ±0.67d | 63.01 ±0.71e |
| | Leaf | | | | |
| IVDMD | 77.04±0.68a | 75.75 ±0.46b | 73.98 ±0.44c | 69.94±0.68d | 68.21 ±0.5ge |
| NDF | 76.70±0.78a | 76.29±0.67a | 74.49±0.68b | 71.83±0.74c | 70.24±0.90d |
| ADF | 69.71 ±0.35a | 69.06±0.22a | 67.63±0.51b | 62.78± 1.05c | 61.86±0.95c |
| Hemicellulose | 55.42 ±0.66a | 86.34±0.57b | 86.18 ±0.47b | 85.13 ±0.25b | 82.27 ±0.95c |
| Cellulose | 87.74±0.47a | 85.88±0.56b | 83.8/ ±0.82c | 78.// ±0.93d | 75./2±0.7/c |
| | Stem | | | | |
| IVDMD | 69.51±0.5a | 66.82±0.17b | 62.02±0.56c | 59.45±0.45d | 58.52±0.32d |
| NDF | 62.34±0.95a | 59.48± 0.22b | 58.67 ± 1.04b | 54.98±0.77c | 54.40±0.67c |
| ADF | 54.11 ±2.49a | 50.91 ±3.10ab | 49.61 ±2.32b | 43.84± 1.28c | 42.n± 1.33c |
| Hemicellulose | 76.03±0.55a | 73.11±1.04b | 70.48±0.42c | 68.81 ±0.13cd | 67.71 ±0.23d |
| Cellulose | 74.44±0.45a | 67.61 ± 1.62b | 60. II ± 1.92c | 51.35 ± 1.07d | 49.39+ 1.25d |

Mean bearing different letters in a row show significant (P < 0.0 1) differences.

Table 2. Digestibility coefficient of OM, NDF, ADF, hemicellulose and cellulose of different cultivars of whole maize plant and its morphological fractions

| Cultivar | Structural components | | | | |
|----------|-----------------------|---------------|-----------------|---------------|---------------|
| | OM | NDF | ADF | Hemicellulose | Cellulose |
| | Whole plant | | | | |
| Akbar | 70.84± 1.84b | 69.08± 1.39b | 62.29±2.04h | 79.15±2.69 | n.16±3.69b |
| Neelum | 71.39 ±2.00a | 69.95 ± 1.42a | 64.51 ±2.00a | 80.02±2.27 | 74.55±3.92a |
| U.M - n | 69.13 ± UOd | 67.38±1.35c | 60.23 ± U8c | 78.01 ±2.25 | 71.69±3.91b |
| I.Z - 31 | 69.85 ± 1.80c | 68.44± 1.55c | 61.57 ±2.06bc | 78.26±2.14 | n.35±3.75b |
| | Leaf | | | | |
| Akbar | 73.45 ± 1.56b | 74.03±1.33b | 67.71 ± 1.12a | 86.82 ±0.88a | 80.63 ±2.1td |
| Neelum | 74.22 ± 1.67a | 75.88±1.14a | 65.15 ± 1.~Ob | 84.37 ± 1.08b | 80.63 ±2.35c |
| U.M - n | 71.96± 1.80c | n.40± 1.12c | (J5,44 ± 1.91 b | 85.76± 1.30ab | 62.35±2.54b |
| I.Z - 3 | n.29± 1.76c | 73.32± 1.38b | | | |
| | Stem | | | | |
| Akbar | 63.22±2.11ab | 57.71±1.85b | 48.44±2.74b | 71.55±1.88 | 59.42 ±4.82bc |
| Neelum | 63.86±2.16a | 60.50±1.58a | 53.75±2.80a | 70.51±1.13. | 63.10±4.25a |
| U.M 81 | 62.88±2.14b | 56.86±1.32b | 45.66±2.23b | 70.98± 1.52 | 58.07±5.11c |
| I.Z 31 | 63.39±2.92 | 56.83±1.22b | 45.11±1.12b | 70.80+ 1.67 | 61.73±4.96ab |

Means bearing different letters in a row show significant (P < 0.0 1) differences.

Table I. Prediction equations for IVDMO (Y) from OM contents of various cultivars of maize (whole plant) and its morphological fractions.

| Varieties | Regression equations | Correlation (r) | S.E. |
|-------------|-----------------------|-----------------|--------|
| Whole plant | | | |
| Akhar | $Y = 79Y - 04t5 OM$ | - 0.930** | ± 0.38 |
| Ncclum | $Y = X0.3 - 0421 OM$ | - 0.906** | ± 043 |
| UM-XI | $Y = n.0 - 0426 OM$ | - 0.940** | ± 036 |
| I.Z - 11 | $Y = 82Y - 0463 OM$ | - 0.946** | ± 0.33 |
| Leaf | | | |
| Akhar | $Y = 82.9 - 0356 OM$ | - 0.972** | ± 021 |
| Ncclum | $Y = 83.5 - 0359 OM$ | - 0.966** | ± 0.24 |
| UM-81 | $Y = 82.1 - 0.406 OM$ | - 0.964** | ± 0.27 |
| I.Z - 31 | $Y = 82.5 - 0.406 OM$ | - 0.959** | ± 0.28 |
| Stem | | | |
| Akbar | $Y = 73.6 - 0.576 OM$ | - 0.940** | ± 043 |
| Ncclum | $Y = 76.3 - 0.751 OM$ | - 0.939** | ± 0.42 |
| CM-XI | $Y = 71.7 - 0.524 OM$ | - 0.862** | ± 0.61 |
| U - 1 | $Y = 72.4 - 0.525 OM$ | - 0.862** | ± 0.50 |

Y = IVDMO; ** = Significant (P < 0.01).




Figure 1. Relationship between In vitro Digestible Organic Matter in the Dry Matter (IVDOM) and Organic Matter (OM) content of whole plant, leaf and stem fractions of maize cultivars.

Ncclum cultivar had higher hemicellulose digestibility than that of other cultivars. The higher hemicellulose digestibility of leaves of Ncclum could be due to its higher hemicellulose content and lower indigestible fibre fractions.

Cellulose: In vitro cellulose digestibility of whole plant and its leaf and stem fractions continued to decrease significantly with advancing stages of growth. Cellulose digestibility coefficients of the leaf fraction of the plant ranged from 75.12±0.71 to

Digestibility of maize fodder

87.74±0.47, being higher than that of whole plant (63.01 ±0.71 to 83.47±0.60), whereas the stem fraction had the lowest (49.39± 1.25 to 74.44±0.45) cellulose digestibility.

It was probably due to lower cell wall and lignin concentration in leaves than stem. Rocha and Vera (1981) reported a decrease in cellulose digestibility by 0.39±0.14% daily in some tropical grasses.

Cellulose digestibility was also affected by different cultivars in whole maize plant, leaf and stem. Cellulose digestibility of Neelum cultivar was found to be significantly higher in all cases (Table 2). However, U.M-81 cultivar had the lowest digestibility of various plant fractions. Neelum cultivar had lower concentration of indigestible fibre fractions, whereas U.M - 81 cultivar had higher lignin concentration, which may be related to lower cellulose digestibility in this variety.

Relationship Between OM Contents and IVOMO:

Correlation was computed between DM contents and IVDMD of different plant fractions. Correlation values (r) were -0.897, -0.118 and -0.892 for whole plant, leaf and stem, respectively. Based on the correlations, regression/prediction equations were computed to work out percent IVDMD (Y) of whole plant, leaf and stem. The prediction equations are given below:

| | |
|--------------|--|
| Whole plant: | $Y = 79.3 - 0.423 \text{ DM (SE, 0.22)}$ |
| Leaf: | $Y = 82.6 - 0.375 \text{ DM (SE, 0.37)}$ |
| Stem: | $Y = 73.3 - 0.578 \text{ DM (SE, 0.51)}$ |

Regression lines showing the relationship between DM contents and IVDMD of whole plant (a), Leaf (b), and stem (c), are presented in Figure 1.

Correlation coefficients were also calculated between DM contents and IVDMD of different cultivars of whole maize plant and its morphological fractions. DM contents were found to be significantly ($P < 0.01$) and negatively correlated with IVDMD of various cultivars of whole plant, leaf and stem. Regression equations along with correlation coefficients (r) and standard error (S.E.) of different varieties of maize plant and its morphological fractions are given in Table 3.

The results showed that as the DM concentration increased there was a decrease in IVDMD of maize (whole plant) and its morphological fractions. This might have negative effect on the palatability and digestibility. It is therefore, apparent, that the plant maturity have adverse effect on the quality of fodder.

Conclusions and Recommendations

- i) Among the maize cultivars, Neelum proved to be the best. This cultivar had higher digestibility and DM contents and lower concentration of undigestible fibre fractions.
- ii) Maize fodder should be cut preferably between 8th - 9th week of age (flowering stage) to obtain fodder having more nutrients and better digestibility for livestock feeding.
- iii) Regression equations should be of great help for quick estimation of nutritive value of maize fodder by scientists working in the field of forage evaluation.

REFERENCES

- AOAC. 1984. Official Methods of Analysis, Association of Official Analytical Chemists, 14th ed. Arlington, Virginia, USA.
- Azim, A., Z. Naseer and A. Ali. 1989. Nutritional evaluation of maize fodder at two different vegetative stages. *AJAS*. 2 (1): 27-34.
- Baumgardt, B.R., J.L. Cason and M.W. Taylor. 1962. Evaluation of forages in the laboratory. I. Comparative accuracy of several methods. *J. Dairy Sci.* 45: 59-61.
- Bolker, H.I. 1963. A lignin-carbohydrate bond as revealed by infrared spectroscopy. *Nature*. 197: 489.
- Choudhry, A.R. 1983. Maize in Pakistan. Punjab Agri. Res. Coordination Board, Univ. Agri. Faisalabad.
- Hunt, e.W., W. Kezar, D.D. Hinman and J.J. Combs. 1993. Effects of hybrids and ensiling with or without microbial inoculant on the nutritional characteristics of whole corn plant. *J. Anim. Sci.* 71 (1): 38-43 (Maize Abst. 9 (6): 3785, 1993).
- Lloveras, J. 1990. Dry matter yield and nutritive value of four summer annual crops in northwest Spain (Galicia). *Grass Forage Sci.* 45 (3): 243-248 (Nutr. Abst. Rev. 60: 6319, 1990).
- McDougall, E.I. 1947. Studies on ruminant saliva. I. The composition and output of sheep's saliva. *Bio. Chem. J.* 43: 99.
- Rocha, G.P. and R.R. Vera. 1981. Structural carbohydrates, protein and in vitro digestibility, of eight tropical grasses. *Turrialba*, 31 (to: 15-20 (Nutr. Abst. Rev. 52 (9): 4334, 1982).
- Sanderson, M.A., J.S. Hornstein and W.F. Wedin. 1989. Alfalfa morphological stage and its relation to in situ digestibility of detergent fibre fractions of stem. *Crop Sci.* 29 (5): 1315-1319 (Nutr. Abst. Rev. 60 (J): 1677, 1990).
- Singh, B. and M.P. Narang. 1990. Nutritive evaluation of *Seteria anceps* cv. Kazungula, *S. anceps* cv. Narok, *Pennisetum clandestinum* and *Avena sativa* forages. *Indian J. Anim. Sci.* 62 (1): 63-68.
- Steel, R.G.D. and J.H. Torrie. 1984. Principles and Procedures of International Student. Ed. McGraw Hill Book Co. Inc., New York.
- Troelsen, J.E. 1971. Consumption of digestible energy by sheep from the concentration of in vitro digestible energy, cell wall constituents, and crude fibre in coarse roughages. *J. Anim. Sci.* 51: 433-438.
- Van Soest, P.J. and J.B. Robertson. 1985. Analysis of forage and fibrous feed: A Laboratory Manual for Animal Science 613, Cornell University, New York.
- Van Soest P.J. 1987. Nutritional Ecology of the Ruminants. Comstock Publishing Associates: A Division of Cornell University Press, Cornell, New York.
- Vona, L.e., G.A. Jung, R.L., Reid and W.e., Sharp. 1984. Nutritive value of warm season grass hays for beef cattle and sheep, digestibility, intake and mineral utilization. *J. Anim. Sci.* 59 (6): 1582-1595 (Nutr. Abst. Rev. 55: 2727, 1985).