

EFFECT OF SPACINGS ON THE GROWTH OF *EUCALYPTUS CAMALDULENSIS* UNDER AGRO-FORESTRY SYSTEMS

Abdul Khaliq Chauhdry and M. Jahangir Ghauri

Senior Research Officer and Senior Research Officer (Biometrics)

Punjab Forestry research Institute, Faisalabad,

INTRODUCTION

Amongst the various species In Agroforestry programme, Eucalyptus has established it self as a tree most suitable for Agroforestry, as it can grow on a variety of soils under various climatic conditions. It has desirable qualities, such as fast rate of growth, clean straight bole, thin crown, ornamental value and less shade casting. Besides that, it has multiple uses i.e. poles, pulp, fuel, charcoal, fiber board, chipboard etc. and above all it gives high and early return.

Due to these characters *Eucalyptus camaldulensis* has been liked very much by farmers. Planting of Eucalyptus is being done by the farmers on field boundaries, in rows along water channel, on farm approach road, as woodlots and in compact blocks in fields with agricultural crops at variable spacings,

In this paper results of effect of various spacings on the growth of *Eucalyptus camaldulensis* under agroforestry systems have been presented.

LAYOUT OF THE STUDY

In order to find out the effect of spacing on the growth of *Eucalyptus camaldulensis* in conjunction with agricultural crops, a study was laid out in Research Garden, Punjab Forestry Research Institute, Faisalabad during May, 1990. Six months old Eucalyptus seedling raised in poly bags were planted at three spacings viz. 1.9 x 1.9 meters, 1.9 x 3.2 meters and 1.9 x 3.8 meter with 126, 70, and

56 plants in each plot of 28 x 17.2 meters dimension. Randomized Complete Block Design was adopted for planting and was replicated three times.

A variety of agricultural crops, one after the other were raised in between the lines by the farmers. These include Mash beans, Bersim and maize fodder. Irrigation with canal water was given according to the requirements of agricultural crops.

RESULTS AND DISCUSSION

Data for height and diameter growth were recorded during December, 1992 at the age of about 3 years and are compared for various spacings in table- I.

Statistical analysis of the above data indicated that the spacings have significant effect on diameter growth while it is non-significant in case of height growth at 95% confidence level. LSD (Least significant Difference) test showed 1.9 x 3.8 meter spacing at 5% confidence level to be the best, while spacing of 1.9 x 3.2 m took intermediate position and 1.9 x 1.9 m spacing gave minimum diameter growth which is logical also.

In order to compare wood production under different spacings, volume per hectare was obtained by using volume table prepared from the material removed in thinning. Bole volume in meter cubes of 89 felled trees were calculated covering diameter range of 4 to 16 cm. The local column table so obtained is given below.

Table 1, Height (m) and Diameter (cm) under different spacings

Replication	Spacings					
	1.9 x 1.9 m		1.9 x 3.2 m		1.9 x 3.8 m	
	Ht.	Dia.	Ht.	Dia.	Ht.	Dia.
R ₁	9.0	8.1	8.8	8.9	9.0	8.9
R ₂	10.2	9.4	8.4	9.4	9.3	10.2
R ₃	9.6	9.1	9.5	9.9	10.3	10.2
Total	28.8	26.6	26.7	28.2	28.6	29.3
Average	9.6	8.9	8.9	9.4	9.5	9.8

Table 2: Wood Production~ per hectare (Cu. m.) under different spacings

Replication	Spacings (Treatment) in meters		
	1.9 x 1.9 m	1.9 x 3.2 m	1.9 x 3.8 m
R ₁	1882.9	1129.8	941.3
R ₂	4393.4	2636.0	2196.6
R ₃	4393.4	2636.0	2296.6
Total	10669.7	6401.8	5334.5
Mean	3256.6	2133.9	1778.2

D.B.H (cms)	Tree Volume (m ³)
4.1 - 6.5	0.0088
6.6 - 9.0	0.0178
9.1-11.5	0.0416
11.6 - 14.0	0.0702
14.1 - 16.5	0.1022

The volume per hectare under various treatments (spacings) for the three replications is given in table 2.

The statistical analysis (F-test) indicated that spacings had significant effect on wood production (volume per hectare) at 95% confidence level. The LSD test showed a significant difference of volume per hectare between spacings of 1.9 x 1.9 m and 1.9 x 3.2 m and also between 1.9 x 1.9 m and 1.9 x 3.8 m. However, there is no significant change in volume when the spacing is increased from 1.9 x 3.2 m to 1.9 x 3.8 m.

After taking the growth data in December, 1992 thinning was done as the tree crop

became congested and farmers were reluctant to put the area under agri-crops cultivation. To widen the spacing, alternate rows of trees were removed keeping into consideration that more space be made available for smooth agri-crop cultivation by easy soil working in between tree rows.

The felled trees were measured and percentage of trees in various dia classes was determined under different spacings and are given in Table-3.

The above table indicates that if the spacing between tree rows is wider, more trees fall in bigger diameter classes over a period of time. The percentage of trees fell down from 68 to 44% in case of smaller diameter class (upto 10 cms) and increased from 32 to 54% in case of higher diameter classes as the spacings increased from 1.9 x 1.9 m to 1.9 x 3.8 m.

Even under wider spacing of 1.9 x 3.8 m, 2% trees entered in the biggest diameter class of 15.1 cms and above where no trees approached to this class under closer spacings.

Table 3: Frequency distribution (in percent) for felled trees under various diameter classes.

Spacing (meters)	No. of trees measured	Diameter Classes (cms)		
		5.1-10.0	10.1-15.0	15.1 and above
1.9 x 1.9	57	68	32	-
1.9 x 3.2	50	60	40	-
1.9 x 3.8	61	44	54	2

CONCLUSION

From the above results and discussion it is concluded that wider spacing between tree rows has a positive and significant effect on tree diameter growth. The spacing of 1.9 x 3.8 meter is more useful for Eucalyptus plantations which in addition to giving increased number of agriculture crop rotations, also enhances tree volume per unit area. Moreover, wider spacing provides better opportunity to the farmer for soil working and increased utility of his land resources by harvesting crops for longer periods of time.

REFERENCES

- Sheikh M.I. and Raza-ul-Haq, 1982. Performance of Poplars and other species in conjunction with agricultural crops. Pak. Jour., For., Vol. 32 (2).
- Sheikh, M. I. and Raza-ul-Haq, 1982. Effect of spacing on the growth of *Dalbergia Sissoo* (Shisham). Pak. Jour. for. Vol. 32 (2).
- Sheikh, M. I. 1981. Management study of *Eucalyptus camaldulensis*. Final technical report, P.F.I, Peshawar.